

Michael Crabb and Andrew Ranicki

The geometric Hopf invariant and surgery theory

March 1, 2016



For Nico Marcel Vallauri

Preface

This joint project is an outcome of two separate projects started by us in the 1970's, which independently dealt with double points of maps of manifolds in geometry and quadratic structures in homotopy theory and algebra, generalizing the Hopf invariant. However, the first-named author was more concerned with \mathbb{Z}_2 -equivariant homotopy theory and simply-connected manifolds, while the second-named author was more concerned with chain complexes and the surgery theory of non-simply-connected manifolds.

The *geometric Hopf invariant* of a stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ for pointed spaces X, Y is the stable \mathbb{Z}_2 -equivariant map

$$h(F) = (F \wedge F)\Delta_X - \Delta_Y F : \Sigma^\infty X \rightarrow \Sigma^\infty(Y \wedge Y)$$

measuring the failure of F to preserve the diagonal maps of X and Y . Here \mathbb{Z}_2 acts on $Y \wedge Y$ by transposition

$$T : Y \wedge Y \rightarrow Y \wedge Y ; (y_1, y_2) \mapsto (y_2, y_1) .$$

The stable \mathbb{Z}_2 -equivariant homotopy class of $h(F)$ is the primary obstruction to desuspending F , i.e. to F being stable homotopic to an unstable map.

The original Hopf invariant $H(F) \in \mathbb{Z}$ of a map $F : S^3 \rightarrow S^2$ ([26]) was defined geometrically to be the linking number of

$$F^{-1}(\{x, y\}) = S^1 \cup S^1 \subset S^3$$

for generic $x \neq y \in S^2$ with $F^{-1}(x), F^{-1}(y) \subset S^3$ unions of disjoint circles. In the 1930's the Hopf invariant map was proved to be an isomorphism

$$H : \pi_3(S^2) \rightarrow \mathbb{Z}_2 ; F \mapsto H(F) ,$$

with the projection of the Hopf bundle

$$S^1 \longrightarrow S^3 \xrightarrow{\eta} S^2$$

such that the fibres $\eta^{-1}(x), \eta^{-1}(y) \subset S^3$ are disjoint circles with linking number $H(\eta) = 1$. Samelson [55] described the historical antecedents (Clifford, Klein) of the Hopf bundle¹. The original definition of the Hopf invariant $H(F) \in \mathbb{Z}$ applied to any map $F : S^{2n-1} \rightarrow S^n$, with $H : \pi_{2n-1}(S^n) \rightarrow \mathbb{Z}$ an isomorphism for $n = 1, 2, 4, 8$.

In the 1940's Steenrod [62] used algebraic topology to interpret the Hopf invariant of a map $F : S^{2n-1} \rightarrow S^n$ as the evaluation of the cup product in the mapping cone $\mathcal{C}(F) = S^n \cup_F D^{2n}$. The mod 2 Hopf invariant $H_2(F) \in \mathbb{Z}_2$ of a map $F : S^j \rightarrow S^k$ was then defined using the mod 2 Steenrod square Sq^{j-k+1} in $\mathcal{C}(F) = S^k \cup_F D^{j+1}$, or equivalently the functional mod 2 Steenrod square Sq_F^{j-k+1} .

For a pointed space X write the loop space and the suspension as usual

$$\Omega X = \text{map}_*(S^1, X), \quad \Sigma X = S^1 \wedge X.$$

In the 1950's and 1960's it was realized that the Hopf invariant was the first example of the primary obstruction to desuspending a stable maps $F : \Sigma^k X \rightarrow \Sigma^k Y$ for any pointed spaces X, Y (not necessarily spheres), with $1 \leq k \leq \infty$. The desuspension of a k -stable map $F : \Sigma^k X \rightarrow \Sigma^k Y$ is equivalent to compressing the adjoint map

$$X \rightarrow \Omega^k \Sigma^k Y ; x \mapsto (s \mapsto F(s, x)) \quad (s \in S^k)$$

into

$$Y \subset \Omega^k \Sigma^k Y ; y \mapsto (s \mapsto (s, y)) .$$

The homotopy groups of $\Omega \Sigma Y$ are

$$\pi_n(\Omega \Sigma Y) = \pi_{n+1}(\Sigma Y) .$$

The desuspension of a map $F : \Sigma X \rightarrow \Sigma Y$ is equivalent to compressing the adjoint map

$$X \rightarrow \Omega \Sigma Y ; x \mapsto (s \mapsto F(s, x)) \quad (s \in S^1)$$

into

$$Y \subset \Omega \Sigma Y ; y \mapsto (s \mapsto (s, y)) .$$

¹ Samelson's paper includes the letter sent by Hopf to Freudenthal giving a first-hand account of the Hopf invariant. The letter was sent on 17th August 1928 from 30 Murray Place, Princeton. By a coincidence, this was the house occupied 50 years later by one of the authors (A.R.)

James [28] showed that for a connected Y there is a homology equivalence

$$\Omega \Sigma Y \simeq \bigvee_{j=1}^{\infty} \bigwedge_j Y .$$

The adjoint of a 1-stable map $F : \Sigma X \rightarrow \Sigma Y$ is a map $\text{adj}(F) : X \rightarrow \Omega \Sigma Y$. The primary obstruction to desuspending F is the composite

$$H_*(X) \xrightarrow{\text{adj}(F)_*} H_*(\Omega \Sigma Y) = \bigoplus_{j=1}^{\infty} H_*(\bigwedge_j Y) \rightarrow H_*(Y \wedge Y) .$$

For $F : S^3 = \Sigma(S^2) \rightarrow S^2 = \Sigma(S^1)$ this is just the Hopf invariant

$$H_2(S^2) = \mathbb{Z} \xrightarrow{\text{adj}(F)_*} H_2(\Omega S^2) = \bigoplus_{j=1}^{\infty} H_2(S^j) = H_2(S^2) = \mathbb{Z} ; 1 \mapsto H(F) .$$

The homotopy groups of the space

$$\Omega^{\infty} \Sigma^{\infty} Y = \varinjlim_k \Omega^k \Sigma^k Y$$

are the stable homotopy groups of Y

$$\pi_n(\Omega^{\infty} \Sigma^{\infty} Y) = \varinjlim_k \pi_{n+k}(\Sigma^k Y) .$$

For connected Y the space $\Omega^{\infty} \Sigma^{\infty} Y$ has a filtration in stable homotopy theory with successive quotients

$$(E \Sigma_k)^+ \wedge_{\Sigma_k} \bigwedge_k Y \quad (k \geq 1) .$$

Here, Σ_k is the permutation group on k letters acting by

$$\Sigma_k \times \bigwedge_k Y \rightarrow \bigwedge_k Y ; (\sigma, (y_1, y_2, \dots, y_k)) \mapsto (y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k)})$$

and $E \Sigma_k$ a contractible space with a free Σ_k -action. (There is also a version for disconnected Y). The k th filtration quotient correspond to the k -tuple points of maps of manifolds. We shall only be concerned with the second filtration quotient, for $k = 2$, as we are particularly interested in double points. We write

$$E \Sigma_2 = S(\infty) , \Sigma_2 = \mathbb{Z}_2 .$$

For any pointed space Y define the *quadratic construction* pointed space

$$Q(Y) = S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y) .$$

Such quadratic spaces were introduced in homotopy theory by Toda, in the 1950's. Following Crabb [12] and Crabb and James [13] the primary obstruction to desuspending a stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ is identified here both with the stable \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant $h(F)$ and with the stable homotopy class of a stable map

$$h'(F) : \Sigma^\infty X \rightarrow \Sigma^\infty Q(Y) .$$

The algebraic theory of surgery of Ranicki [49, 50] was based on the *quadratic construction* on a chain complex C , the chain complex defined by

$$Q(C) = W \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_{\mathbb{Z}} C)$$

with $W = C(S(\infty))$ a free $\mathbb{Z}[\mathbb{Z}_2]$ -module resolution of \mathbb{Z} . A stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ was shown to induce a natural class of chain maps

$$\psi_F : \dot{C}(X) \rightarrow Q(\dot{C}(Y)) = \dot{C}(Q(Y))$$

called the *quadratic construction* on F .

The homotopy groups of the space

$$\Omega^\infty \Sigma^\infty Y = \varinjlim_k \Omega^k \Sigma^k Y$$

are the stable homotopy groups of Y

$$\pi_n(\Omega^\infty \Sigma^\infty Y) = \varinjlim_k \pi_{n+k}(\Sigma^k Y) .$$

The desuspension of F is equivalent to compressing the adjoint map

$$X \rightarrow \Omega^\infty \Sigma^\infty Y ; x \mapsto (s \mapsto F(s, x)) \ (s \in S^\infty)$$

into

$$Y \subset \Omega^\infty \Sigma^\infty Y ; y \mapsto (s \mapsto (s, y)) .$$

For connected Y the space $\Omega^\infty \Sigma^\infty Y$ has a filtration in stable homotopy theory with successive quotients

$$(E\Sigma_k)^+ \wedge_{\Sigma_k} \bigwedge_k Y \ (k \geq 1) .$$

Here, Σ_k is the permutation group on k letters acting by

$$\Sigma_k \times \bigwedge_k Y \rightarrow \bigwedge_k Y ; (\sigma, (y_1, y_2, \dots, y_k)) \mapsto (y_{\sigma(1)}, y_{\sigma(2)}, \dots, y_{\sigma(k)})$$

and $E\Sigma_k$ a contractible space with a free Σ_k -action. (There is also a version for disconnected Y). The k th filtration quotient correspond to the k -tuple points of maps of manifolds. We shall only be concerned with the first filtration quotient, for $k = 2$, as we are particularly interested in double points. We write

$$E\Sigma_2 = S(\infty), \Sigma_2 = \mathbb{Z}_2.$$

For any pointed space Y define the *quadratic construction* pointed space

$$Q(Y) = S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y).$$

Such quadratic spaces were introduced in homotopy theory by Toda, in the 1950's. Following Crabb [12] and Crabb and James [13] the primary obstruction to desuspending a stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ is identified here both with the stable \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant $h(F)$ and with the stable homotopy class of a stable map

$$h'(F) : \Sigma^\infty X \rightarrow \Sigma^\infty Q(Y).$$

The algebraic theory of surgery of Ranicki [49, 50] was based on the *quadratic construction* on a chain complex C , the chain complex defined by

$$Q(C) = W \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_{\mathbb{Z}} C)$$

with $W = C(S(\infty))$ a free $\mathbb{Z}[\mathbb{Z}_2]$ -module resolution of \mathbb{Z} . A stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ was shown to induce a natural class of chain maps

$$\psi_F : \dot{C}(X) \rightarrow Q(\dot{C}(Y)) = \dot{C}(Q(Y))$$

called the *quadratic construction* on F .

In May 1999 it became clear to us that the quadratic construction ψ_F of [49, 50] was induced by the geometric Hopf invariant $h(F)$ of [12, 13]. In this book we unite the two approaches into a single theory, together with the applications to the double points of immersions and to various constructions in the algebraic theory of surgery. In the applications it is necessary to consider π -equivariant, fibrewise and local versions of the quadratic construction/geometric Hopf invariant.

The double point set of a map $f : M \rightarrow N$ is the free \mathbb{Z}_2 -space

$$D_2(f) = \{(x, y) \in M \times M \mid x \neq y \in M, f(x) = f(y) \in N\}$$

with $T \in \mathbb{Z}_2$ acting by

$$T : D_2(f) \rightarrow D_2(f) ; (x, y) \mapsto (y, x).$$

An immersion of manifolds $f : M^m \looparrowright N^n$ has a normal $(n-m)$ -plane bundle ν_f , with a Thom space $T(\nu_f)$. There exists a map $e : M \rightarrow \mathbb{R}^k$ (k large) such that

$$M \hookrightarrow \mathbb{R}^k \times N ; x \mapsto (e(x), f(x))$$

is an embedding, and the Pontrjagin-Thom construction gives a stable Umkehr map

$$F : \Sigma^\infty N^+ \rightarrow \Sigma^\infty T(\nu_f) .$$

f is an embedding if and only if $D_2(f) = \emptyset$, in which case F is unstable. Our main result is the Double Point Theorem 6.19, which stably factors the geometric Hopf invariant $h(F)$ for a generic immersion f through the double point set

$$h(F) : N^+ \rightarrow D_2(f) \rightarrow Q(T(\nu_f)) .$$

The stable homotopy class of $h(F)$ depends only on the regular homotopy class of f . If f is regular homotopic to an embedding then $h(F)$ is stably null-homotopic. In Crabb and Ranicki [14] we interpreted the Smale-Hirsch-Haefliger regular homotopy classification of immersions f in the metastable dimension range $3m < 2n - 1$ (when a generic f has no triple points) in terms of the geometric Hopf invariant $h(F)$. In fact, [14] is reprinted here in Appendix C, and may be regarded as an extended introduction to this book.

We use the geometric Hopf invariant to provide the homotopy theory underpinnings of the following constructions of surgery theory:

1. The double point invariant of Wall [68] for a generic immersion $f : S^m \looparrowright N^{2m}$

$$\mu(f) \in \mathbb{Z}[\pi_1(N)] / \{x - (-)^m \bar{x} \mid x \in \mathbb{Z}[\pi_1(N)]\} \quad (\bar{g} = g^{-1} \in \pi_1(N))$$

such that $\mu(f) = 0$ if (and, for $m > 2$, only if) f is regular homotopic to an embedding.

2. The Wall surgery obstruction $\sigma_*(f, b) \in L_n(\mathbb{Z}[\pi_1(X)])$ of an n -dimensional normal map $(f, b) : M \rightarrow X$ was expressed as the cobordism class of an n -dimensional $\mathbb{Z}[\pi_1(X)]$ -module chain complex C with a quadratic Poincaré duality determined via the quadratic construction ψ_F of a $\pi_1(X)$ -equivariant stable Umkehr map $F : \Sigma^\infty(\tilde{X}^+) \rightarrow \Sigma^\infty(\tilde{M}^+)$. Here \tilde{X} is the universal cover of X , $\tilde{M} = f^* \tilde{X}$ is the pullback cover of M , and

$$H_*(C) = \ker(\tilde{f}_* : H_*(\tilde{M}) \rightarrow H_*(\tilde{X})) .$$

3. The total surgery obstruction $s(X) \in \mathbb{S}_n(X)$ of an n -dimensional geometric Poincaré complex X is expressed the cobordism class of the $\mathbb{Z}[\pi_1(X)]$ -contractible \mathbb{Z} -module chain complex

$$C = \mathcal{C}([X] \cap - : C(X)^{n-*} \rightarrow C(X))_{*+1}$$

with an X -local $(n-1)$ -dimensional quadratic Poincaré duality determined via the spectral quadratic construction $s\psi_F$ of a $\pi_1(X)$ -equivariant semistable homotopy equivalence $F : \Sigma^\infty T(\nu_{\tilde{X}})^* \rightarrow \Sigma^\infty(\tilde{X}^+)$ inducing the chain equivalence $[X] \cap -$.

Chapters 1-6 develop the relationship between the geometric Hopf invariant and double points in such a natural way that the π -equivariant constructions fall out without extra work. Along the way, we point out the many particular instances of this relationship in the literature, and explain how they fit into our development. The applications to non-simply-connected surgery theory are only considered in Chapter 7.

In principle, it is possible to develop surgery theory in the context of the fibrewise homotopy theory of Crabb and James [13], relating it to the controlled and bounded surgery theories. However, we shall not do this here, restricting ourselves to the application of \mathbb{Z}_2 -equivariant fibrewise homotopy theory in Appendix A to obtain the π -equivariant quadratic construction used in the conventional non-simply-connected surgery theory.

The treatment of \mathbb{Z}_2 -equivariant bordism theory in Chapter 4 is somewhat elaborate, although appropriate to the application to double point sets. Appendix B provides a much more succinct treatment of \mathbb{Z}_2 -equivariant bordism theory, including the extension to G -equivariant bordism for any finite group G .

In 2010 we published a joint paper *The geometric Hopf invariant and double points* ([14]) which analyzes double points using a combination of stable fibrewise homotopy and immersion theories, obtaining the double point theorem of Chapter 6 from a somewhat different viewpoint. The paper is reprinted here in Appendix C.

Michael Crabb, University of Aberdeen, mccrabb@abdn.ac.uk

Andrew Ranicki, University of Edinburgh, a.ranicki@ed.ac.uk

February 2016

Contents

1	The difference construction	1
1.1	Joins, cones and suspensions	1
1.2	The difference construction $\delta(f, g)$	4
1.3	Chain complexes	11
2	Umkehr maps and inner product spaces	15
2.1	Adjunction and compactification	15
2.2	Inner product spaces	18
2.3	The addition and subtraction of maps	22
3	Stable homotopy theory	39
3.1	Stable maps	39
3.2	Vector bundles	47
3.3	Bordism	59
3.4	S -duality	63
3.5	The stable cohomotopy Thom and Euler classes	68

4	\mathbb{Z}_2-equivariant homotopy and bordism theory	77
4.1	π -equivariant homotopy theory	77
4.2	The bi-degree	85
4.3	Stable \mathbb{Z}_2 -equivariant homotopy theory	93
4.4	\mathbb{Z}_2 -equivariant bundles	113
4.5	\mathbb{Z}_2 -equivariant S -duality	137
4.6	\mathbb{Z}_2 -equivariant bordism	142
5	The geometric Hopf invariant	161
5.1	The Q -groups	161
5.2	The symmetric construction $\phi_V(X)$	177
5.3	The geometric Hopf invariant $h_V(F)$	189
5.4	The stable geometric Hopf invariant $h'_V(F)$	204
5.5	The quadratic construction $\psi_V(F)$	207
5.6	The ultraquadratic construction $\hat{\psi}(F)$	213
5.7	The spectral quadratic construction $s\psi_V(F)$	221
5.8	Stably trivialized vector bundles	226
6	The double point theorem	245
6.1	Framed manifolds	245
6.2	Double points	263
6.3	Positive embeddings	273
6.4	Finite covers	284
6.5	Function spaces	290

Contents	13
6.6 Embeddings and immersions	302
6.7 Linking and self-linking	316
6.8 Intersections and self-intersections for $M^m \rightarrow N^{2m}$	328
6.9 The simply-connected surgery obstruction	332
7 The π-equivariant geometric Hopf invariant	347
7.1 π -equivariant S -duality	347
7.2 The π -equivariant constructions	348
7.3 Non-simply-connected surgery obstruction theory	354
A The fibrewise Umkehr map	357
B Notes on \mathbb{Z}_2-equivariant bordism	365
B.1 Introduction	365
B.2 Framed bordism	365
B.3 Unoriented bordism	375
B.4 Finite groups	380
C Reprint of “The geometric Hopf invariant and double points” Journal of Fixed Point Theory and Applications 7, 325–350 (2010)	
(Including the correction of a misprint in Proposition 3.1)	383
References	409
Index	413

Chapter 1

The difference construction

Chapter 1 describes the difference construction in both homotopy and chain homotopy theory. The homotopy version will be used in Chapter 2 to define the subtraction of maps.

We shall only be concerned with locally compact Hausdorff spaces X .

1.1 Joins, cones and suspensions

We shall be dealing with both pointed and unpointed spaces, and likewise for maps and homotopies. For a pointed space X we shall denote the base point by $*$ $\in X$.

The *one-point adjunction* of an unpointed space X is the pointed space

$$X^+ = X \sqcup \{*\}$$

obtained by adjoining a base point $*$, with open subsets $U, U \sqcup \{*\} \subseteq X^+$ for open subsets $U \subseteq X$.

The *smash product* of pointed spaces X, Y is the pointed space

$$X \wedge Y = X \times Y / (X \times \{*\} \cup \{*\} \times Y) .$$

In particular, for unpointed spaces A, B

$$A^+ \wedge B^+ = (A \times B)^+ .$$

The *join* of spaces X, Y is the space defined as usual by

$$X * Y = I \times X \times Y / \{(0, x, y_1) \sim (0, x, y_2), (1, x_1, y) \sim (1, x_2, y)\} ,$$

so that

$$X = [0 \times X \times Y] , Y = [1 \times X \times Y] \subset X * Y .$$

The *cone* on a space X is the space

$$cX = X * \{\text{pt.}\} = I \times X / \{1\} \times X = I \wedge X^+$$

with I pointed at 1. Identify $X = \{0\} \times X \subset cX$. A map $cX \rightarrow Y$ is a map $X \rightarrow Y$ with a homotopy to a constant map. The *reduced cone* on a pointed space X is the pointed space

$$CX = cX / c\{*\} = I \times X / (\{1\} \times X \cup I \times \{*\}) = I \wedge X .$$

A pointed homotopy $h : f \simeq g : X \rightarrow Y$ is a pointed map $h : I^+ \wedge X \rightarrow Y$ with

$$h(0, x) = f(x) , h(1, x) = g(x) \in Y \quad (x \in X) .$$

A *null-homotopy* of a pointed map $f : X \rightarrow Y$ is a pointed homotopy $f \simeq * : X \rightarrow Y$, which is the same as an extension of f to a pointed map $\delta f : CX \rightarrow Y$.

The *suspension* of a space X is the space

$$sX = S^0 * X = I \times X / \{(i, x) \sim (i, x') \mid i = 0, 1, x, x' \in X\} .$$

A map $sX \rightarrow Y$ is a map $X \rightarrow Y$ with two homotopies to constant maps. The *reduced suspension* of a pointed space X is the pointed space

$$\Sigma X = sX / s\{*\} = I \times X / \{0, 1\} \times X \cup I \times \{*\} = S^1 \wedge X .$$

A pointed map $\Sigma X \rightarrow Y$ is a pointed map $X \rightarrow Y$ with two null-homotopies.

The *mapping cylinder* of a map $F : X \rightarrow Y$ is the space

$$m(F) = I \times X \cup_F Y / \{(0, x) \sim F(x) \mid x \in X\} .$$

The *reduced mapping cylinder* of a pointed map $F : X \rightarrow Y$ is the pointed space

$$\mathcal{M}(F) = m(F) / I \times \{*\} = I^+ \wedge X \cup_F Y .$$

The *mapping cone* of a map $F : X \rightarrow Y$ is the space

$$c(F) = m(F) / \{1\} \times X = cX \cup_F Y .$$

The *reduced mapping cone* of a pointed map F is the pointed space

$$\mathcal{C}(F) = c(F)/\{1\} \times X = CX \cup_F Y .$$

Proposition 1.1. (i) For any map $F : X \rightarrow Y$ the maps

$$\begin{aligned} i & : X \rightarrow m(F) ; x \mapsto [1, x] , \\ j & : Y \rightarrow m(F) ; y \mapsto [y] , \\ k & : I \times X \rightarrow m(F) ; (t, x) \mapsto [t, x] \end{aligned}$$

are such that j is a homotopy equivalence, and $k : jF \simeq i : X \rightarrow m(F)$. Similarly for a pointed map F and the pointed mapping cylinder $\mathcal{M}(F)$.

(ii) For any pointed map $F : X \rightarrow Y$ the reduced mapping cone $Z = \mathcal{C}(F)$ fits into a cofibration sequence

$$X \xrightarrow{F} Y \xrightarrow{G} Z \xrightarrow{H} \Sigma X \xrightarrow{\Sigma F} \Sigma Y \longrightarrow \dots$$

with $G : Y \rightarrow Z$ the inclusion and $H : Z \rightarrow \Sigma X$ the projection. The inclusion $CX \rightarrow Z$ defines null-homotopy $GF \simeq * : X \rightarrow Z$.

□

Example 1.2. For an unpointed space X define the pointed map

$$F : X^+ \rightarrow S^0 ; x \mapsto -1 , * \mapsto +1 .$$

The reduced mapping cylinder and cone of F are the cylinder and cone on X

$$\mathcal{M}(F) = cX , \mathcal{C}(F) = sX .$$

By Proposition 1.1 there is defined a cofibration sequence of pointed spaces

$$X^+ \xrightarrow{F} S^0 \xrightarrow{G} sX \xrightarrow{H} \Sigma X^+ \xrightarrow{\Sigma F} S^1 \longrightarrow \dots$$

with

$$\begin{aligned} G & : S^0 \rightarrow sX ; -1 \mapsto (0, X) , 1 \mapsto (1, X) , \\ H & : sX \rightarrow sX/S^0 = \Sigma X^+ ; (t, x) \mapsto (t, x) , \\ \Sigma F & : \Sigma X^+ \rightarrow S^1 = I/(0=1) ; (t, x) \mapsto t . \end{aligned}$$

The reduced suspension ΣX^+ is obtained from sX by identifying the two suspension points $S^0 \subset sX$, with G the inclusion and H the projection. If X is non-empty then for any $x \in X$ then the pointed maps

$$\begin{aligned} E & : S^0 \rightarrow X^+ ; 1 \mapsto * , -1 \mapsto x , \\ \Sigma E & : \Sigma(S^0) = S^1 \rightarrow \Sigma X^+ ; t \mapsto (t, x) \end{aligned}$$

are such that $FE = 1 : S^0 \rightarrow S^0$ and

$$H \vee \Sigma E : sX \vee S^1 \rightarrow \Sigma X^+ .$$

is a homotopy equivalence. \square

1.2 The difference construction $\delta(f, g)$

The *sum* of homotopies

$$f : c \simeq d , g : d \simeq e : A \rightarrow B$$

is the concatenation homotopy

$$f + g : c \simeq e : A \rightarrow B$$

defined by

$$f + g : A \times I \rightarrow B ; (a, t) \mapsto \begin{cases} f(a, 2t) & \text{if } 0 \leq t \leq 1/2 \\ g(a, 2t - 1) & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

The *reverse* of a homotopy $f : c \simeq d : A \rightarrow B$ is the homotopy

$$-f : d \simeq c : A \rightarrow B$$

defined by

$$-f : A \times I \rightarrow B ; (a, t) \mapsto f(a, 1 - t) .$$

As usual, given pointed spaces A, B let $[A, B]$ be the set of pointed homotopy classes of pointed maps $A \rightarrow B$.

A pointed map $d : \Sigma A \rightarrow B$ is essentially the same as a pointed homotopy

$$d : \{*\} \simeq \{*\} : A \rightarrow B$$

and $[\Sigma A, B]$ is a group, with multiplication and inverses given by

$$\begin{aligned} [\Sigma A, B] \times [\Sigma A, B] &\rightarrow [\Sigma A, B] ; (f, g) \mapsto f + g , \\ [\Sigma A, B] &\rightarrow [\Sigma A, B] ; h \mapsto -h \end{aligned}$$

Definition 1.3. Let $e : A \rightarrow B$ be a pointed map, and let

$$f : e \simeq * , g : e \simeq * : A \rightarrow B$$

be two null-homotopies, corresponding to two extensions $f, g : CA \rightarrow B$ of e . The *rel A difference* of f, g is the pointed map

$$\delta(f, g) = f \cup -g : \Sigma A \rightarrow B ; (t, a) \mapsto \begin{cases} g(1 - 2t, a) & \text{if } 0 \leq t \leq 1/2 \\ f(2t - 1, a) & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

Equivalently

$$\delta(f, g) : \Sigma A \xrightarrow{-1 \cup 1} CA \cup_A CA \xrightarrow{g \cup f} B$$

with $-1 \cup 1$ the homeomorphism

$$-1 \cup 1 : \Sigma A \rightarrow CA \cup_A CA ; (t, a) \mapsto \begin{cases} (1 - 2t, a)_1 & \text{if } 0 \leq t \leq 1/2 \\ (2t - 1, a)_2 & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

□

Proposition 1.4. *Every pointed map $d : \Sigma A \rightarrow B$ is the rel A difference $\delta(f, g)$ of null-homotopies $f : e \simeq *$, $g : e \simeq *$ of a map $e : A \rightarrow B$.*

Proof. The pointed maps

$$\begin{aligned} e &: A \rightarrow B ; a \mapsto d(1/2, a) , \\ f &: CA \rightarrow B ; (t, a) \mapsto d((1 + t)/2, a) , \\ g &: CA \rightarrow B ; (t, a) \mapsto d((1 - t)/2, a) \end{aligned}$$

are such that

$$e(a) = f(0, a) = g(0, a) \in B \ (a \in A) , \ \delta(f, g) = d .$$

□

Remark 1.5. (i) In the special case $e = * : A \rightarrow B$ the null-homotopies $f, g : e \simeq \{*\}$ in Proposition 1.3 are just maps $f, g : \Sigma A \rightarrow B$. The sum is

$$f + g = f \cup g : \Sigma A = CA \cup_A CA \rightarrow B$$

and the rel A difference is just

$$\delta(f, g) = f - g = f \cup -g : \Sigma A = CA \cup_A CA \rightarrow B .$$

(ii) In the special case $A = S^n$ the rel A difference of maps $f, g : CA = D^{n+1} \rightarrow B$ such that $f|_S = g|_S : S^n \rightarrow B$ is the *separation element* $\delta(f, g) :$

$\Sigma A = S^{n+1} \rightarrow B$ of James [29, Appendix].

□

Proposition 1.6. (i) *A homotopy $h : f \simeq g : CA \rightarrow B$ which is constant on A determines a null-homotopy*

$$\delta(h) : \delta(f, g) \simeq * : \Sigma A \rightarrow B .$$

(ii) *For any map $f : CA \rightarrow B$ there is a canonical null-homotopy*

$$\delta(f, f) \simeq * : \Sigma A \rightarrow B .$$

(iii) *Let $i : X \rightarrow A$ be the inclusion of a subspace $X \subseteq A$ such that the projection $\mathcal{C}(i) \rightarrow A/X$ is a homotopy equivalence, and let $j : A \rightarrow A/X$ be the projection. If $f, g : CA \rightarrow B$ are maps which agree on $A \cup CX \subseteq CA$ the composite*

$$\delta(f, g)(\Sigma i) : \Sigma X \xrightarrow{\Sigma i} \Sigma A \xrightarrow{\delta(f, g)} B$$

*is equipped with a canonical null-homotopy $h : \delta(f, g)(\Sigma i) \simeq *$, giving rise to a rel $A \cup CX$ difference map*

$$\delta(f, g) : \Sigma(A/X) = CA/(A \cup CX) \simeq \mathcal{C}(\Sigma i) \xrightarrow{\delta(f, g) \cup h} B$$

such that the rel A difference map factors up to homotopy as

$$\delta(f, g) : \Sigma A \xrightarrow{\Sigma j} \Sigma(A/X) \xrightarrow{\delta(f, g)} B .$$

Proof. (i) The map defined by

$$\delta(h) : C(\Sigma A) \rightarrow B ;$$

$$(s, t, a) \mapsto \begin{cases} h(2s, 1 - 2t, a) & \text{if } 0 \leq s, t \leq 1/2 \\ f(2t - 1, a) & \text{if } 0 \leq s \leq 1/2 \leq t \leq 1 \\ f(1 + 4(s - 1)t, a) & \text{if } 0 \leq t \leq 1/2 \leq s \leq 1 \\ f(4s - 3 + 4(1 - s)t, a) & \text{if } 1/2 \leq s, t \leq 1 \end{cases}$$

is a null-homotopy of $\delta(f, g)$.

(ii) This is the special case $f = g$ of (i), with $h : f \simeq g$ the constant homotopy.

(iii) Let

$$e = f|_A = g|_A : A \cup CX \rightarrow B ,$$

so that

$$\delta(f, g)(\Sigma i) : \Sigma X \rightarrow B ; (t, x) \mapsto \begin{cases} e(1 - 2t, x) & \text{if } 0 \leq t \leq 1/2 \\ e(2t - 1, x) & \text{if } 1/2 \leq t \leq 1 \end{cases}$$

with a canonical null-homotopy $h : \delta(f, g)(\Sigma i) \simeq *$.

□

Thus if $f, g : CA \rightarrow B$ agree on $A \cup CX \subseteq CA$ the homotopy class of $\delta(f, g) : \Sigma(A/X) \rightarrow B$ is an obstruction to the existence of a homotopy $f \simeq g : CA \rightarrow B$ which is constant on $A \cup CX$.

If f, g agree on a neighbourhood of $A \subset CA$ and $f = *$ on the frontier and outside the neighbourhood, then $\delta(f, g)$ is homotopic to the restriction of g to the complement of the neighbourhood :

Proposition 1.7. (i) *If $f, g : CA \rightarrow B$ are maps such that for some neighbourhood $U \subseteq CA$ of $A \subseteq CA$*

$$g(t, a) = \begin{cases} f(t, a) & \text{if } (t, a) \in U \\ * & \text{if } (t, a) \in \overline{CA \setminus U} \end{cases}$$

then there exists a homotopy

$$\gamma(f, g) : \delta(f, g) \simeq f' : \Sigma A \rightarrow B$$

with f' defined by

$$f' : \Sigma A = CA/A \rightarrow B ; (t, a) \mapsto \begin{cases} * & \text{if } (t, a) \in U \\ f(t, a) & \text{if } (t, a) \in \overline{CA \setminus U} \end{cases} .$$

(ii) *The homotopy $\gamma(f, g)$ in (i) can be chosen to be natural, meaning that given commutative squares of maps*

$$\begin{array}{ccc} CA_1 & \xrightarrow{f_1} & B_1 \\ h \downarrow & & \downarrow k \\ CA_2 & \xrightarrow{f_2} & B_2 \end{array} , \quad \begin{array}{ccc} CA_1 & \xrightarrow{g_1} & B_1 \\ h \downarrow & & \downarrow k \\ CA_2 & \xrightarrow{g_2} & B_2 \end{array}$$

such that for neighbourhoods $U_i \subseteq CA_i$ of $A_i \subseteq CA_i$ with $U_2 = (1 \times h)(U_1)$

$$g_i(t, a) = \begin{cases} f_i(t, a) & \text{if } (t, a) \in U_i \\ * & \text{if } (t, a) \in \overline{CA_i \setminus U_i} \end{cases} \quad (i = 1, 2)$$

there is defined a commutative square

$$\begin{array}{ccc}
I \times \Sigma A_1 & \xrightarrow{\gamma(f_1, g_1)} & B_1 \\
1 \times \Sigma h \downarrow & & \downarrow k \\
I \times \Sigma A_2 & \xrightarrow{\gamma(f_2, g_2)} & B_2
\end{array}$$

Proof. (i) We have

$$\begin{aligned}
\delta(f, g)(t, a) &= \begin{cases} g(1 - 2t, a) & \text{if } 0 \leq t \leq 1/2 \\ f(2t - 1, a) & \text{if } 1/2 \leq t \leq 1 \end{cases} \\
&= \begin{cases} * & \text{if } 0 \leq t \leq 1/2 \text{ and } (1 - 2t, a) \in \overline{CA \setminus U} \\ g(1 - 2t, a) & \text{if } 0 \leq t \leq 1/2 \text{ and } (1 - 2t, a) \in U \\ g(2t - 1, a) & \text{if } 1/2 \leq t \leq 1 \text{ and } (2t - 1, a) \in U \\ f'(2t - 1, a) & \text{if } 1/2 \leq t \leq 1 \text{ and } (2t - 1, a) \in \overline{CA \setminus U} \end{cases} .
\end{aligned}$$

Define a homotopy $\gamma(f, g) : \delta(f, g) \simeq f'$ by

$$\gamma(f, g) : I \times \Sigma A \rightarrow X ; (s, t, a) \mapsto \begin{cases} g((1 - 2s)(1 - 2t) + 2s, a) & \text{if } 0 \leq s, t \leq 1/2 \text{ and } (1 - 2t, a) \in U \\ g((1 - 2s)(2t - 1) + 2s, a) & \text{if } 0 \leq s \leq 1/2 \leq t \leq 1 \text{ and } (2t - 1, a) \in U \\ f'(2t - 1, a) & \text{if } 0 \leq s \leq 1/2 \text{ and } (2t - 1, a) \in \overline{CA \setminus U} \\ f'((2 - 2s)(2t - 1) + (2s - 1)t, a) & \text{if } 1/2 \leq s, t \leq 1 \\ * & \text{otherwise} . \end{cases}$$

(ii) The homotopy $\gamma(f, g)$ constructed in (i) is natural.

□

Example 1.8. Here are two special cases of Proposition 1.7.

(i) For any map $f : CA \rightarrow B$ take $f = g$, $U = CA$ to obtain a homotopy

$$\gamma(f, f) : \delta(f, f) \simeq * : \Sigma A \rightarrow B .$$

(ii) For any map $g : CA \rightarrow B$ such that

$$g(0, a) = * \in B \quad (a \in A)$$

take $f = *$, $U = A$ to obtain a homotopy

$$\gamma(*, g) : \delta(*, g) \simeq [g] : \Sigma A \rightarrow B .$$

□

Proposition 1.9. *For any pointed space X and any map $F : A \rightarrow B$ there is induced a Barratt-Puppe exact sequence of groups and pointed sets*

$$\dots \longrightarrow [\Sigma B, X] \xrightarrow{\Sigma F} [\Sigma A, X] \xrightarrow{H} [\mathcal{C}(F), X] \xrightarrow{G} [B, X] \xrightarrow{F} [A, X] .$$

A map $q \cup r : \mathcal{C}(F) \rightarrow X$ is the same as a map $r : B \rightarrow X$ together with a null-homotopy $q : rF \simeq * : A \rightarrow X$. Use the projection

$$\begin{aligned} \nabla : \mathcal{C}(F) &\rightarrow \mathcal{C}(F) \vee \Sigma A ; \\ (t, x) &\mapsto \begin{cases} (2t, x) \in \mathcal{C}(F) & \text{if } 0 \leq t \leq 1/2 \\ (2t - 1, x) \in \Sigma A & \text{if } 1/2 \leq t \leq 1 \end{cases} \quad (x \in A) \\ y &\mapsto y \quad (y \in B) \end{aligned}$$

to define a pairing

$$[\mathcal{C}(F), X] \times [\Sigma A, X] \rightarrow [\mathcal{C}(F), X] ; (q \cup r, s) \mapsto ((q \cup r) \vee s) \nabla .$$

Proposition 1.10. *Suppose that $F : A \rightarrow B$, $r : B \rightarrow X$ are maps and there are given two null-homotopies $p, q : CA \rightarrow X$ of $rF : A \rightarrow X$*

$$p, q : rF \simeq * : A \rightarrow X .$$

The maps $p \cup r, q \cup r : \mathcal{C}(F) \rightarrow X$ are both extensions of r , with

$$(p \cup r)G = (q \cup r)G = r \in [B, X] ,$$

and the rel A difference

$$\delta(p, q) : \Sigma A \rightarrow X ; (t, x) \mapsto \begin{cases} q(1 - 2t, x) & \text{if } 0 \leq t \leq 1/2 \\ p(2t - 1, x) & \text{if } 1/2 \leq t \leq 1 \end{cases}$$

is such that there exists a homotopy

$$h : p \cup r \simeq ((q \cup r) \vee \delta(p, q)) \nabla : \mathcal{C}(F) \rightarrow X .$$

Thus if $[\mathcal{C}(F), X]$ has a group structure (e.g. if $\mathcal{C}(F)$ is a suspension or X is an H -space)

$$\begin{aligned} p \cup r - q \cup r &= H(\delta(p, q)) \\ &\in \ker(G : [\mathcal{C}(F), X] \rightarrow [B, X]) = \text{im}(H : [\Sigma A, X] \rightarrow [\mathcal{C}(F), X]) . \end{aligned}$$

Proof. By construction

$$\begin{aligned} & ((q \cup r) \vee \delta(p, q)) \nabla : \mathcal{C}(F) \rightarrow X ; \\ & (t, x) \mapsto \begin{cases} q(2t, x) & \text{if } 0 \leq t \leq 1/2 \\ q(3 - 4t, x) & \text{if } 1/2 \leq t \leq 3/4 \quad (x \in A) \\ p(4t - 3, x) & \text{if } 3/4 \leq t \leq 1 , \end{cases} \\ & y \mapsto y \quad (y \in B) . \end{aligned}$$

Define a homotopy $h : p \cup r \simeq ((q \cup r) \vee \delta(p, q)) \nabla$ by

$$\begin{aligned} & h : I \wedge \mathcal{C}(F) \rightarrow X ; \\ & (s, (t, x)) \mapsto \begin{cases} q(2st, x) & \text{if } 0 \leq t \leq s/2 \\ q(3s - 4t, x) & \text{if } s/2 \leq t \leq 3s/4 \quad (x \in A) \\ p((4t - 3s)/(4 - 3s), t) & \text{if } 3s/4 \leq t \leq 1 , \end{cases} \\ & y \mapsto y \quad (y \in B) . \end{aligned}$$

□

Proposition 1.11. (i) *A commutative diagram of spaces and maps*

$$\begin{array}{ccc} A & \xrightarrow{F} & B \\ a \downarrow & & \downarrow b \\ A' & \xrightarrow{F'} & B' \end{array}$$

induces a natural transformation of cofibration sequences

$$\begin{array}{ccccccccc} A & \xrightarrow{F} & B & \xrightarrow{G} & \mathcal{C}(F) & \xrightarrow{H} & \Sigma A & \xrightarrow{\Sigma F} & \dots \\ a \downarrow & & b \downarrow & & (a, b) \downarrow & & \Sigma a \downarrow & & \\ A' & \xrightarrow{F'} & B' & \xrightarrow{G'} & \mathcal{C}(F') & \xrightarrow{H'} & \Sigma A' & \xrightarrow{\Sigma F'} & \dots \end{array}$$

(ii) *Suppose given a commutative diagram as in (i), with the maps a, b null-homotopic. For any null-homotopies $\delta a : CA \rightarrow A'$, $\delta b : CB \rightarrow B'$ of a, b the map (a, b) is determined up to homotopy by the rel A difference of null-homotopies of $F'a = bF : A \rightarrow B'$,*

$$f = (\delta b)CF , \quad g = F'(\delta a) : CA \rightarrow B'$$

with

$$(a, b) \simeq G' \delta(f, g) H : \mathcal{C}(F) \rightarrow \mathcal{C}(F') .$$

(iii) Suppose given a commutative diagram as in (i), with b null-homotopic. For any null-homotopy $\delta b : CB \rightarrow B'$ of b the map (a, b) is homotopic to the rel A difference of the inclusion $i : CA \rightarrow \mathcal{C}(F')$ and the composite

$$j = G'(\delta b)CF : CA \rightarrow CB \rightarrow B' \rightarrow \mathcal{C}(F') ,$$

that is

$$(a, b) \simeq \delta(j, i) : \Sigma A \rightarrow \mathcal{C}(F') .$$

□

1.3 Chain complexes

The difference construction for maps of spaces induces the difference construction for chain maps of chain complexes.

Let A be a ring.

Definition 1.12. (i) The *suspension* of an A -module chain complex C is the A -module chain complex SC with

$$d_{SC} = d_C : (SC)_r = C_{r-1} \rightarrow (SC)_{r-1} = C_{r-2} .$$

(ii) Let $f : C \rightarrow D$ be an A -module chain map. The *relative difference* of chain homotopies

$$p : f \simeq 0 : C \rightarrow D , q : f \simeq 0 : C \rightarrow D$$

is the A -module chain map

$$\delta(p, q) = q - p : SC \rightarrow D .$$

(iii) The *algebraic mapping cone* of an A -module chain map $f : C \rightarrow D$ is the A -module chain complex $\mathcal{C}(f)$ with

$$d_{\mathcal{C}(f)} = \begin{pmatrix} d_D & (-)^r f \\ 0 & d_C \end{pmatrix} : \mathcal{C}(f)_r = D_r \oplus C_{r-1} \rightarrow \mathcal{C}(f)_{r-1} = D_{r-1} \oplus C_{r-2} .$$

The algebraic mapping cone fits into a short exact sequence

$$0 \longrightarrow D \xrightarrow{g} \mathcal{C}(f) \xrightarrow{h} SC \longrightarrow 0$$

with

$$\begin{aligned} g &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} : D_r \rightarrow \mathcal{C}(f)_r = D_r \oplus C_{r-1} , \\ h &= (0 \ 1) : \mathcal{C}(f)_r = D_r \oplus C_{r-1} \rightarrow SC_r = C_{r-1} . \end{aligned}$$

□

Given a space X let $C(X)$ denote the singular \mathbb{Z} -module chain complex, and for a pointed space let

$$\dot{C}(X) = C(X, \{*\})$$

be the reduced singular chain complex. The Eilenberg-Zilber theorem gives a natural chain equivalence

$$S\dot{C}(X) \simeq \dot{C}(\Sigma X) .$$

Proposition 1.13. (i) *A map of spaces $F : X \rightarrow Y$ induces a chain map $f : C(X) \rightarrow C(Y)$.*

(ii) *A homotopy $H : F \simeq G : X \rightarrow Y$ induces a chain homotopy*

$$h : f \simeq g : C(X) \rightarrow C(Y) .$$

(iii) *The reverse homotopy $-H : G \simeq F : X \rightarrow Y$ induces the reverse chain homotopy $-h : g \simeq f : C(X) \rightarrow C(Y)$ (up to a natural higher chain homotopy).*

(iv) *The sum $H_1 + H_2 : F_1 \simeq F_3$ of homotopies*

$$H_1 : F_1 \simeq F_2 , H_2 : F_2 \simeq F_3 : X \rightarrow Y$$

induces the sum chain homotopy

$$h_1 + h_2 : f_1 \simeq f_3 : C(X) \rightarrow C(Y)$$

(up to a natural higher chain homotopy).

(v) *A pointed map $F : X \rightarrow Y$ induces a chain map $f : \dot{C}(X) \rightarrow \dot{C}(Y)$, and similarly for chain homotopies. The reduced singular chain complex of the mapping cone $\mathcal{C}(F)$ is chain equivalent to the algebraic mapping cone of f*

$$\dot{C}(\mathcal{C}(F)) \simeq \mathcal{C}(f) ,$$

and the cofibration sequence of pointed spaces (1.1)

$$X \xrightarrow{F} Y \xrightarrow{G} \mathcal{C}(F) \xrightarrow{H} \Sigma X \xrightarrow{\Sigma F} \Sigma Y \longrightarrow \dots$$

induces the sequence of chain complexes

$$\dot{C}(X) \xrightarrow{f} \dot{C}(Y) \xrightarrow{g} \mathcal{C}(f) \xrightarrow{h} S\dot{C}(X) \xrightarrow{Sf} S\dot{C}(Y) \longrightarrow \dots$$

with g, h as in 1.12.

(vi) The relative difference $\delta(P, Q) : \Sigma X \rightarrow Y$ of null-homotopies

$$P : F \simeq \{*\} , Q : F \simeq \{*\} : X \rightarrow Y$$

of a pointed map $F : X \rightarrow Y$ induces the relative difference of the induced null-chain homotopies

$$p : f \simeq 0 , q : f \simeq 0 : \dot{C}(X) \rightarrow \dot{C}(Y)$$

i.e. there is defined a commutative diagram

$$\begin{array}{ccc} S\dot{C}(X) & \xrightarrow{\delta(p, q) = q - p} & \dot{C}(Y) \\ & \searrow \simeq & \nearrow \delta(P, Q) \\ & \dot{C}(\Sigma X) & \end{array}$$

□

Chapter 2

Umkehr maps and inner product spaces

This Chapter develops a coordinate-free approach to stable homotopy theory, using inner product spaces.

2.1 Adjunction and compactification

As in §1 we shall only be concerned with locally compact Hausdorff spaces X .

Definition 2.1. The *one-point compactification* X^∞ of a space X is the space $X \cup \{\infty\}$ with open sets $U, (X \setminus K) \cup \{\infty\} \subseteq X$ for open subsets $U \subseteq X$ and compact subsets $K \subseteq X$.

□

The canonical map $X^\infty \rightarrow X^+$ from the one-point compactification to the one-point adjunction is a continuous bijection which is a homeomorphism if and only if X is compact. For any spaces X, Y

$$(X \times Y)^\infty = X^\infty \wedge Y^\infty .$$

For any pair of spaces $(A, B \subseteq A)$ the quotient A/B is a pointed space with base point $* = [B] \in A/B$. The quotient space A/B understood to be A^+ if $B = \emptyset$. The projection $\pi^+ : A^+ \rightarrow A/B$ is a map such that a subspace $U \subseteq A/B$ is open if and only if $(\pi^+)^{-1}(U) \subseteq A^+$ is open. The canonical map $(A \setminus B)^+ \rightarrow A/B$ is a continuous bijection which is a homeomorphism if and only if B is a component of A . The canonical map $(A \setminus B)^\infty \rightarrow A/B$ is a continuous bijection which is a homeomorphism if and only if A is compact.

Definition 2.2. The *adjunction Umkehr map* of an injective map $f : X \hookrightarrow Y$ with respect to a subspace $Y_0 \subseteq Y \setminus f(X)$ is the projection

$$F^+ : Y/Y_0 \rightarrow Y/(Y \setminus f(X)) .$$

□

The quotient pointed space $Y/(Y \setminus f(X))$ has open sets $f(U)$, $[V]$ with $U \subseteq X$, $Y \setminus f(X) \subseteq V \subseteq Y$ such that $f(U)$, $V \subseteq Y$ are open. The function

$$Y/(Y \setminus f(X)) \rightarrow X^+ ; f(x) \mapsto x , [Y \setminus f(X)] \mapsto *$$

is a continuous bijection.

Definition 2.3. (i) An *open embedding* $f : X \hookrightarrow Y$ is an injective map such that a subset $U \subseteq X$ is open if and only if $f(U) \subseteq Y$ is open. It follows that $f| : K \rightarrow f(K)$ is a homeomorphism for any subspace $K \subseteq X$, giving $f(K) \subseteq Y$ the subspace topology.
(ii) The *compactification Umkehr map* of an open embedding f as in (i) is the surjective function

$$F^\infty : Y^\infty \rightarrow X^\infty ; y \mapsto \begin{cases} x & \text{if } y = f(x) \in f(X) \\ \infty & \text{otherwise} \end{cases}$$

which preserves the base points and is continuous: for any open subset $U \subseteq X$ and any closed compact subset $K \subseteq X$

$$\begin{aligned} (F^\infty)^{-1}(U) &= f(U) \subset Y^\infty , \\ (F^\infty)^{-1}((X \setminus K) \cup \{\infty\}) &= (Y \setminus f(K)) \cup \{\infty\} \subseteq Y^\infty \end{aligned}$$

with $f(U) \subseteq Y$ open and $f(K) \subseteq Y$ closed compact. □

Example 2.4. Let $(X, \partial X)$, $(Y, \partial Y)$ be n -dimensional manifolds with boundary, and let $f : X \hookrightarrow Y$ is an injective map such that

$$f(\partial X) \cap \partial Y = \emptyset .$$

(i) The map

$$X/\partial X \rightarrow Y/(Y \setminus f(X)) ; [x] \mapsto [f(x)]$$

is a homeomorphism. The adjunction Umkehr map of f with respect to $Y_0 = \partial Y \subset Y$ is the Pontryagin-Thom map

$$F^+ : Y/\partial Y \rightarrow Y/(Y \setminus f(X)) = X/\partial X .$$

(ii) If X, Y are compact the compactification Umkehr of the open embedding defined by the restriction to the interiors

$$g = f| : X \setminus \partial X \hookrightarrow Y \setminus \partial Y ,$$

is the adjunction Umkehr F^+ of (i)

$$G^\infty = F^+ : (Y \setminus \partial Y)^\infty = Y/\partial Y \rightarrow (X \setminus \partial X)^\infty = X/\partial X .$$

□

The Umkehr maps of isotopic embeddings are homotopic.

Definition 2.5. (i) An *isotopy* is a homotopy

$$h : I \times X \rightarrow Y ; (t, x) \mapsto h(t, x)$$

such that

$$I \times X \rightarrow I \times Y ; (t, x) \mapsto (t, h(t, x))$$

is an embedding.

(ii) An isotopy h is *open* if $I \times X \hookrightarrow I \times Y$ is an open embedding. □

Proposition 2.6. Let $h : I \times X \rightarrow Y$ be an isotopy, so that for each $t \in I$ there is defined an embedding

$$f_t : X \hookrightarrow Y ; x \mapsto h(t, x) .$$

(i) If $Y_0 \subseteq Y$ is a subspace such that $Y_0 \subseteq Y \setminus h(I \times X)$ the adjunction Umkehr maps

$$F_t^+ : Y/Y_0 \rightarrow Y/(Y \setminus h(I \times X)) \quad (t \in I)$$

are homotopic.

(ii) If h is an open isotopy then each f_t is an open embedding, and the compactification Umkehr maps

$$F_t^\infty : Y^\infty \rightarrow X^\infty \quad (t \in I)$$

are homotopic.

Proof. (i) The adjunction Umkehr maps F_t^+ define a homotopy

$$H^+ : I \times Y/Y_0 \rightarrow Y/h(X \times I) ; (t, y) \mapsto F_t^+(y)$$

(ii) The compactification Umkehr maps F_t^∞ define a homotopy

$$H^\infty : I \times Y^\infty \rightarrow X^\infty ; (t, y) \mapsto F_t^\infty(y) = \begin{cases} x & \text{if } y = f_t(x) \in f_t(X) \\ \infty & \text{otherwise} . \end{cases}$$

□

2.2 Inner product spaces

Inner product spaces are given the metric topology.

Terminology 2.7 Given an inner product space V write the unit disc, unit sphere and projective space of V as

$$\begin{aligned} D(V) &= \{v \in V \mid \|v\| \leq 1\} , \\ S(V) &= \{v \in V \mid \|v\| = 1\} , \\ P(V) &= S(V)/\{v \sim -v\} . \end{aligned}$$

For finite-dimensional V the spaces $D(V), S(V), P(V)$ are compact.

□

Proposition 2.8. *Let V be an inner product space.*

(i) *The maps*

$$\begin{cases} D(V) \setminus S(V) \rightarrow V ; v \mapsto \frac{v}{1 - \|v\|} \\ V \rightarrow D(V) \setminus S(V) ; v \mapsto \frac{v}{1 + \|v\|} \end{cases}$$

are inverse homeomorphisms.

(ii) *The maps*

$$\begin{cases} D(V)/S(V) \rightarrow V^\infty ; v \mapsto \frac{v}{1 - \|v\|} \\ V^\infty \rightarrow D(V)/S(V) ; v \mapsto \begin{cases} \frac{v}{1 + \|v\|} & \text{for } v \in V \\ [S(V)] & \text{for } v = \infty \end{cases} \end{cases}$$

are inverse bijections which are inverse homeomorphisms for finite-dimensional V .

(iii) *The maps*

$$\left\{ \begin{array}{ll} (0, S(V)) & \text{for } v = 0 \in V \\ V^\infty \rightarrow sS(V) ; v \mapsto \left(\frac{\|v\|}{1 + \|v\|}, \frac{v}{\|v\|} \right) & \text{for } v \in V \setminus \{0\} \\ (1, S(V)) & \text{for } v = \infty \in V^\infty \\ sS(V) \rightarrow V^\infty ; (t, u) \mapsto [t, u] = \frac{tu}{1-t} \end{array} \right.$$

are inverse bijections which are inverse homeomorphisms for finite-dimensional V .

(iv) The map

$$S(V) \times \mathbb{R} \rightarrow V \setminus \{0\} ; (v, x) \mapsto ve^x$$

is a homeomorphism, inducing a homeomorphism

$$(S(V) \times \mathbb{R})^\infty = \Sigma S(V)^\infty \rightarrow (V \setminus \{0\})^\infty .$$

Thus for finite-dimensional V

$$(V \setminus \{0\})^\infty = \Sigma S(V)^+ .$$

□

For $V = \mathbb{R}^k$ there are identifications

$$\begin{aligned} D(\mathbb{R}^k) &= D^k, S(\mathbb{R}^k) = S^{k-1}, \\ (\mathbb{R}^k)^\infty &= D^k/S^{k-1} = S^k, P(\mathbb{R}^k) = \mathbb{RP}^{k-1} \end{aligned}$$

and for any space X

$$\begin{aligned} (\mathbb{R}^k \times X)^\infty &= S^k \wedge X^\infty = \Sigma^k X^\infty, \\ (D^k \times X)/(S^{k-1} \times X) &= (D^k/S^{k-1}) \wedge X^+ = \Sigma^k X^+ . \end{aligned}$$

Proposition 2.9. *Let V be an inner product space.*

(i) The map

$$S(V) \times [0, 1) \rightarrow V ; (t, u) \mapsto [t, u] = \frac{tu}{1-t}$$

is a surjection, which induces a homeomorphism

$$(S(V) \times [0, 1))/(S(V) \times \{0\}) = S(V)^+ \wedge [0, 1) \rightarrow V$$

with $[0, 1)$ based at 0.

(ii) For finite-dimensional V there is defined a pushout diagram

$$\begin{array}{ccc}
S(V)^+ & \xrightarrow{s_V} & 0^+ \\
\downarrow & & \downarrow 0_V \\
C(S(V)^+) & \longrightarrow & V^\infty
\end{array}$$

with

$$\begin{aligned}
s_V &= \text{projection} : S(V)^+ \rightarrow S(V)^+/S(V) = S^0, \\
C(S(V)^+) &= S(V)^+ \wedge [0, 1] \quad ([0, 1] \text{ based at } 1), \\
S(V)^+ &\rightarrow C(S(V)^+) ; u \mapsto (0, u), \\
C(S(V)^+) &\rightarrow V^\infty ; (t, u) \mapsto [t, u].
\end{aligned}$$

Proof. By construction. \square

In view of the pushout square of Proposition 2.9 (ii) for finite-dimensional V a pointed map $p : V^\infty \wedge X \rightarrow Y$ can be viewed as a map $r : X \rightarrow Y$ together with a null-homotopy

$$q : r(s_V \wedge 1_X) \simeq \{*\} : S(V)^+ \wedge X \rightarrow Y.$$

More precisely:

Proposition 2.10. *For any pointed spaces X, Y and a finite-dimensional inner product space V there are natural one-one correspondences between*

- (a) maps $p : V^\infty \wedge X \rightarrow Y$,
- (b) maps $q : CS(V)^+ \wedge X \rightarrow Y$ such that

$$q(0, u, x) = q(0, v, x) \in Y \quad (u, v \in S(V), x \in X),$$

- (c) maps $r : X \rightarrow Y$ together with a null-homotopy

$$\delta r : CS(V)^+ \wedge X \rightarrow Y$$

of

$$r(s_V \wedge 1_X) : S(V)^+ \wedge X \rightarrow Y ; (u, x) \mapsto r(x).$$

Proof. (a) \implies (b) Given a map $p : V^\infty \wedge X \rightarrow Y$ define a null-homotopy

$$\delta p : CS(V)^+ \wedge X \rightarrow Y ; (t, u, x) \mapsto p([t, u], x)$$

of

$$(p|)(s_V \wedge 1_X) : S(V)^+ \wedge X \rightarrow Y ; (u, x) \mapsto p(0, x) .$$

(b) \implies (a) Given a map $q : CS(V)^+ \wedge X \rightarrow Y$ such that

$$q(0, u, x) = q(0, v, x) \in Y \quad (u, v \in S(V), x \in X)$$

define a map

$$p : V^\infty \wedge X \rightarrow Y ; ([t, u], x) \mapsto q([t, u], x)$$

with $q = \delta p$.

(b) \implies (c) Given q define

$$r : X \rightarrow Y ; x \mapsto q([0, u], x) \text{ (for any } u \in S(V) \text{)} .$$

(c) \implies (b) Given $r, \delta r$ define $q = \delta r$. □

Proposition 2.11. (i) *For any inner product spaces U, V*

$$D(U \oplus V) = D(U) \times D(V) ,$$

$$S(U \oplus V) = S(U) \times D(V) \cup D(U) \times S(V)$$

and there is defined a homeomorphism

$$\lambda_{U,V} : S(U) * S(V) \rightarrow S(U \oplus V) ; (t, u, v) \mapsto (u \cos(\pi t/2), v \sin(\pi t/2))$$

with inverse

$$\lambda_{U,V}^{-1} : S(U \oplus V) \rightarrow S(U) * S(V) ; (x, y) \mapsto \left(\frac{2}{\pi} \tan^{-1} \left(\frac{\|y\|}{\|x\|} \right), \frac{x}{\|x\|}, \frac{y}{\|y\|} \right) .$$

(ii) *For finite-dimensional U, V there is a cofibration sequence of pointed spaces*

$$S(V)^+ \xrightarrow{i_{U,V}} S(U \oplus V)^+ \xrightarrow{j_{U,V}} S(U)^+ \wedge V^\infty \xrightarrow{k_{U,V}} \Sigma S(V)^+$$

and a pushout diagram of pointed spaces

$$\begin{array}{ccc} S(U \oplus V)^+ & \xrightarrow{j_{U,V}} & S(U)^+ \wedge V^\infty \\ \downarrow & & \downarrow k_{U,V} \\ CS(U \oplus V)^+ & \xrightarrow{\delta j_{U,V}} & \Sigma S(V)^+ \end{array}$$

with

$$\begin{aligned}
i_{U,V} &= \text{inclusion} : S(V)^+ \rightarrow S(U \oplus V) ; v \mapsto (0, v) , \\
j_{U,V} &= \text{projection} : S(U \oplus V)^+ = (S(U) * S(V))^+ \\
&\rightarrow S(U \oplus V)/(S(V) \times D(U)) = (D(V) \times S(U)/S(V) \times S(U)) \\
&= S(U)^+ \wedge V^\infty ; (t, u, v) \mapsto (u, [t, v]) , \\
\delta j_{U,V} &: CS(U \oplus V)^+ \rightarrow \Sigma S(V)^+ ; (s, (t, u, v)) \mapsto (s + (1-s)t, v) , \\
k_{U,V} &: S(U)^+ \wedge V^\infty \rightarrow \Sigma S(V)^+ ; (u, [t, v]) \mapsto (t, v) .
\end{aligned}$$

(iii) For finite-dimensional V there is a natural one-one correspondence between the homotopy classes of pointed maps $f : \Sigma S(V)^+ \wedge X \rightarrow Y$ and the homotopy classes of pairs (g, h) defined by a pointed map $g : S(U)^+ \wedge V^\infty \wedge X \rightarrow Y$ and a null-homotopy

$$h : g(j_{U,V} \wedge 1_X) \simeq \{*\} : S(U \oplus V)^+ \wedge X \rightarrow Y .$$

Proof. (i)+(ii) By construction.

(iii) By (ii) there is defined a pushout square

$$\begin{array}{ccc}
S(U \oplus V)^+ \wedge X & \xrightarrow{j_{U,V} \wedge 1_X} & S(U)^+ \wedge V^\infty \wedge X \\
\downarrow & & \downarrow k_{U,V} \wedge 1_X \\
CS(U \oplus V)^+ \wedge X & \xrightarrow{\delta j_{U,V} \wedge 1_X} & \Sigma S(V)^+ \wedge X
\end{array}$$

□

2.3 The addition and subtraction of maps

The compactification Umkehr construction is now used to define the addition and subtraction of maps $F : V^\infty \wedge X \rightarrow Y$ for a non-zero inner product space V .

In dealing with open embeddings $e : V \hookrightarrow V$ we shall only be concerned with ones (e.g. smooth) such that the compactification Umkehr map is a homotopy equivalence $F : V^\infty \rightarrow V^\infty$.

Definition 2.12. (i) A homotopy equivalence $F : V^\infty \rightarrow V^\infty$ is

$$\begin{cases} \text{orientation-preserving} \\ \text{orientation-reversing} \end{cases} \quad \text{if}$$

$$\text{degree}(F) = \begin{cases} +1 \\ -1 \end{cases}.$$

(ii) An embedding $e : V \hookrightarrow V$ is $\begin{cases} \text{orientation-preserving} \\ \text{orientation-reversing} \end{cases}$ if the compact-

ification Umkehr map $F^\infty : V^\infty \rightarrow V^\infty$ is an $\begin{cases} \text{orientation-preserving} \\ \text{orientation-reversing} \end{cases}$

homotopy equivalence.

(iii) A *sum map* for V is any map

$$\nabla_V : V^\infty \rightarrow V^\infty \vee V^\infty$$

in the homotopy class of the compactification Umkehr map of an open embedding $e : V \times \{1, 2\} \hookrightarrow V$ such that the restrictions $e| : V \times \{i\} \hookrightarrow V$ ($i = 1, 2$) are orientation-preserving.

(iv) A *difference map* for V is a map

$$\bar{\nabla}_V : V^\infty \rightarrow V^\infty \vee V^\infty$$

in the homotopy class of the compactification Umkehr map of an open embedding $\bar{e} : V \times \{1, 2\} \hookrightarrow V$ such that the restriction $\bar{e}| : V \times \{i\} \hookrightarrow V$ is orientation-reversing for $i = 1$ and orientation-preserving for $i = 2$. \square

Remark 2.13. For $i = 1, 2$ let

$$\pi_i : V^\infty \vee V^\infty \rightarrow V^\infty ; v_j \mapsto \begin{cases} v_j & \text{if } i = j \\ \infty & \text{if } i \neq j \end{cases} \quad (j = 1, 2).$$

(i) If $\dim(V) \geq 2$ a map $\nabla_V : V^\infty \rightarrow V^\infty \vee V^\infty$ is a sum map if and only if the maps $\pi_1 \nabla_V, \pi_2 \nabla_V : V^\infty \rightarrow V^\infty$ are orientation-preserving homotopy equivalences.

(ii) If $\dim(V) \geq 2$ a map $\bar{\nabla}_V : V^\infty \rightarrow V^\infty \vee V^\infty$ is a difference map if and only if the maps $\pi_1 \bar{\nabla}_V, \pi_2 \bar{\nabla}_V : V^\infty \rightarrow V^\infty$ are homotopy equivalences with $\pi_1 \bar{\nabla}_V$ orientation-reversing and $\pi_2 \bar{\nabla}_V$ orientation-preserving. \square

Example 2.14. (i) For any $x \in V$ there is defined an orientation-preserving open embedding

$$e_x : V \rightarrow V ; v \mapsto x + \frac{v}{1 + \|v\|}$$

such that

$$e_x(0) = x \in V, \quad e_x(V) = \{v \in V \mid \|v - x\| < 1\}.$$

(ii) For any $y \in S(V)$ let

$$L_y = \{ty \mid t \in \mathbb{R}\}, \quad H_y = \{z \in V \mid \langle y, z \rangle = 0 \in \mathbb{R}\} \subset V,$$

so that H_y the hyperplane orthogonal to the line $L_y \subset V$ containing y . Reflection in H_y defines an orientation-reversing isomorphism

$$\begin{aligned} R_y &= -1_{L_y} \oplus 1_{H_y} : V = L_y \oplus H_y \rightarrow V = L_y \oplus H_y; \\ v &= (ty, z) \mapsto v - 2\langle v, y \rangle y = (-ty, z). \end{aligned}$$

For any $x \in V$ the composite $\bar{e}_{x,y} = e_x R_y : V \rightarrow V$ is an orientation-reversing open embedding such that

$$\bar{e}_{x,y}(0) = x \in V, \quad \bar{e}_{x,y}(V) = \{v \in V \mid \|v - x\| < 1\}.$$

(iii) For any $v \in V$ with $\|v\| = 1$ define an open embedding

$$f_v : V \times \{v, -v\} \hookrightarrow V; \quad (u, \pm v) \mapsto \pm v + \frac{u}{1 + \|u\|}$$

with compactification Umkehr map a sum map

$$\begin{aligned} F_v &= \nabla_v : V^\infty \rightarrow (V \times \{v, -v\})^\infty = V^\infty \vee V^\infty; \\ u &\mapsto \begin{cases} \left(\frac{u - v}{1 - \|u - v\|} \right)_1 & \text{if } \|u - v\| < 1 \\ \left(\frac{u + v}{1 - \|u + v\|} \right)_2 & \text{if } \|u + v\| < 1 \\ * & \text{otherwise} \end{cases} \end{aligned}$$

such that $F_{-v} = T F_v$ with $T : V^\infty \vee V^\infty \rightarrow V^\infty \vee V^\infty$ the transposition involution.

(iv) If $\nabla_V : V^\infty \rightarrow V^\infty \vee V^\infty$ is a sum map then

$$\bar{\nabla}_V = (R_y \vee 1) \nabla_V : V^\infty \rightarrow V^\infty \vee V^\infty$$

is a difference map, with R_y as in (ii).

(v) For a decomposition $V = \mathbb{R} \oplus W$ as in (ii) use the homeomorphism

$$V^\infty \cong \Sigma W^\infty; \quad (t, w) \mapsto \left(\frac{e^t}{e^t + 1}, w \right)$$

to identify $V^\infty = \Sigma W^\infty$ and define sum and difference maps

$$\begin{aligned}\nabla_V : V^\infty &\rightarrow V^\infty \vee V^\infty ; (t, w) \mapsto \begin{cases} (2t, w)_1 & \text{if } 0 \leq t \leq 1/2 \\ (2t-1, w)_2 & \text{if } 1/2 \leq t \leq 1 \end{cases} , \\ \bar{\nabla}_V : V^\infty &\rightarrow V^\infty \vee V^\infty ; (t, w) \mapsto \begin{cases} (1-2t, w)_1 & \text{if } 0 \leq t \leq 1/2 \\ (2t-1, w)_2 & \text{if } 1/2 \leq t \leq 1 \end{cases} .\end{aligned}$$

□

Proposition 2.15. *For any pointed spaces X, Y and an inner product space V the set $[V^\infty \wedge X, Y]$ of homotopy classes of maps $V^\infty \wedge X \rightarrow Y$ has a group structure for $\dim(V) \geq 1$ (abelian for $\dim(V) \geq 2$), with the sum and difference of $f, g : V^\infty \wedge X \rightarrow Y$ given by*

$$\begin{aligned}f + g &= (g \vee f)(\nabla_V \wedge 1_X) : V^\infty \wedge X \rightarrow Y , \\ f - g &= (g \vee f)(\bar{\nabla}_V \wedge 1_X) : V^\infty \wedge X \rightarrow Y .\end{aligned}$$

□

Example 2.16. (i) The map

$$\begin{aligned}\nabla_V : V^\infty &\rightarrow V^\infty \vee V^\infty ; \\ [t, u] = v &\mapsto \begin{cases} [2t, u]_1 = \left(\frac{2v}{1 - \|v\|} \right)_1 & \text{if } 0 \leq t \leq 1/2 \\ [2t-1, u]_2 = \left(\frac{(\|v\| - 1)v}{2\|v\|} \right)_2 & \text{if } 1/2 \leq t \leq 1 \end{cases}\end{aligned}$$

is a sum map such that $\nabla_V(0) = 0_1$, $\nabla_V(\infty) = \infty_2$.

(ii) The composite of the projection $V^\infty \rightarrow V^\infty/S(V)$ and the homeomorphism

$$\begin{aligned}V^\infty/S(V) &\rightarrow V^\infty \vee V^\infty ; \\ [t, u] = v &\mapsto \begin{cases} [1-2t, u]_1 = \left(\frac{(1 - \|v\|)v}{2\|v\|} \right)_1 & \text{if } 0 \leq t \leq 1/2 \\ [2t-1, u]_2 = \left(\frac{(\|v\| - 1)v}{2\|v\|} \right)_2 & \text{if } 1/2 \leq t \leq 1 \end{cases}\end{aligned}$$

is a difference map

$$\bar{\nabla}_V : V^\infty \rightarrow V^\infty/S(V) \cong V^\infty \vee V^\infty ; [t, u] \mapsto \begin{cases} [1-2t, u]_1 & \text{if } 0 \leq t \leq 1/2 \\ [2t-1, u]_2 & \text{if } 1/2 \leq t \leq 1 \end{cases}$$

such that $\bar{\nabla}_V(0) = \bar{\nabla}_V(\infty) = \infty$.

□

Proposition 2.17. *Let V be a non-zero finite-dimensional inner product space.*

(i) *There is defined a commutative diagram of maps and spaces*

$$\begin{array}{ccccccc} \mathbb{R} \times S(V) & \longrightarrow & S(\mathbb{R} \oplus V) & \longrightarrow & \Sigma S(V)^+ & \longrightarrow & \Sigma S(V)^+/S(V) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ V \setminus \{0\} & \longrightarrow & V^\infty & \longrightarrow & V^\infty/0^+ & \longrightarrow & V^\infty \vee_0 V^\infty \end{array}$$

with the vertical maps homeomorphisms, and

$$V^\infty \vee_0 V^\infty = (V^\infty \vee V^\infty)/\{0_1 \sim 0_2\} .$$

(ii) *There is defined a commutative square of homeomorphisms*

$$\begin{array}{ccc} (\mathbb{R} \times S(V))^\infty & \xrightarrow{\cong} & \Sigma S(V)^+ \\ \cong \downarrow & & \downarrow \cong \\ (V \setminus \{0\})^\infty & \xrightarrow{\cong} & V^\infty/0^+ \end{array}$$

(iii) *The maps $\alpha_V, \beta_V, \gamma_V, s_V, 0_V$ defined by*

$$\alpha_V : V^\infty \rightarrow \Sigma S(V)^+ ; [t, u] \mapsto (t, u) ,$$

$$\beta_V : V^\infty \rightarrow V^\infty \vee_0 V^\infty ;$$

$$[t, u] = v \mapsto \begin{cases} [1 - 2t, u]_1 = \left(\frac{1 - \|v\|}{2\|v\|} v \right)_1 & \text{if } 0 \leq t \leq 1/2 \\ [2t - 1, u]_2 = \left(\frac{\|v\| - 1}{2\|v\|} v \right)_2 & \text{if } 1/2 \leq t \leq 1 , \end{cases}$$

$$\gamma_V : \Sigma S(V)^+ \rightarrow \Sigma S(V)^+/S(V) \cong V^\infty \vee_0 V^\infty ;$$


$$(t, u) \mapsto \begin{cases} [1 - 2t, u]_1 & \text{if } 0 \leq t \leq 1/2 \\ [2t - 1, u]_2 & \text{if } 1/2 \leq t \leq 1 , \end{cases}$$

$$s_V = \text{projection} : S(V)^+ \rightarrow S(V)^+/S(V) = S^0 = \{0, \infty\} ;$$

$$u \mapsto 0 \ (u \in S(V)) ,$$

$$0_V : S^0 \rightarrow V^\infty ; \infty \mapsto \infty , 0 \mapsto 0$$

fit into a commutative braid of cofibrations of unpointed spaces



(which extends to a braid of cofibrations of pointed spaces on the right). For any $u \in S(V)$

$$\alpha_V(u) = \alpha_V[1/2, u] = (1/2, u) \in \Sigma S(V)^+,$$

$$\beta_V(u) = \beta_V[1/2, u] = \gamma_V(1/2, u) = 0_1 = 0_2 \in V^\infty \vee_0 V^\infty.$$

The map $\alpha_V : V^\infty \rightarrow (\mathbb{R} \times S(V))^\infty \cong \Sigma S(V)^+$ is the compactification Umkehr map of the open embedding

$$\mathbb{R} \times S(V) \hookrightarrow V; (x, u) \mapsto e^x u,$$


and γ_V pinches $S(V) \subset \Sigma S(V)^+$ to $0 \in V^\infty \vee_0 V^\infty$.

(iii) There exists a homotopy

$$\beta_V \simeq i\overline{\nabla}_V : V^\infty \rightarrow V^\infty \vee_0 V^\infty$$

with $\overline{\nabla}_V : V^\infty \rightarrow V^\infty \vee V^\infty$ a difference map for V , and $i : V^\infty \vee V^\infty \rightarrow V^\infty \vee_0 V^\infty$ the projection.

(iv) For finite-dimensional inner product spaces U, V there is defined a commutative braid of cofibrations of pointed spaces



Proof. (i) The projection

$$V^\infty \rightarrow (V \setminus \{0\})^\infty = V^\infty / 0^+$$

is the compactification Umkehr map of the open embedding $V \setminus \{0\} \hookrightarrow V$, and

$$\begin{aligned} \mathbb{R} \times S(V) &\cong V \setminus \{0\} ; (s, u) \mapsto e^s u , \\ \mathbb{R} \times S(V) &\longrightarrow S(\mathbb{R} \oplus V) ; (s, u) \mapsto \left(\frac{e^{-s} - e^s}{e^{-s} + e^s}, \frac{2u}{e^{-s} + e^s} \right) , \end{aligned}$$

and the other maps defined as follows. Stereographic projection defines inverse homeomorphisms

$$\begin{aligned} S(\mathbb{R} \oplus V) &\cong V^\infty ; (t, v) \mapsto \frac{v}{1-t} , (1, 0) \mapsto \infty , \\ V^\infty &\cong S(\mathbb{R} \oplus V) ; w \mapsto \left(\frac{\|w\|^2 - 1}{\|w\|^2 + 1}, \frac{2w}{\|w\|^2 + 1} \right) , \infty \mapsto (1, 0) . \end{aligned}$$

For any $t \in I$, $u \in S(V)^+$ define $[t, u] \in V^\infty$ by

$$[t, u] = \begin{cases} \frac{tu}{1-t} & \text{if } t < 1 \text{ and } u \neq \infty \\ \infty & \text{if } t = 1 \text{ or } u = \infty . \end{cases}$$

Every $v \neq 0 \in V$ has a unique expression as $v = [t, u]$ with

$$t = \frac{\|v\|}{1 + \|v\|} \in (0, 1) , u = \frac{v}{\|v\|} \in S(V) .$$

The projection

$$I \wedge S(V)^+ \rightarrow V^\infty ; (t, u) \mapsto [t, u]$$

pinches $\{0\} \times S(V)$ to $0 \subset V^\infty$, inducing the homeomorphism

$$\Sigma S(V)^+ \cong V^\infty / 0^+ ; (t, u) \mapsto [t, u]$$

with inverse

$$V^\infty / 0^+ \cong \Sigma S(V)^+ ; v \mapsto \begin{cases} \left(\frac{\|v\|}{1 + \|v\|}, \frac{v}{\|v\|} \right) & \text{if } v \neq 0 \\ \infty & \text{if } v = 0 . \end{cases}$$

The elements of V^∞ can thus be written as $[t, u]$ with $0 \leq t \leq 1$, $u \in S(V)$, identifying

$$0 = [0, u] , u = [1/2, u] , \infty = [1, u] \in V^\infty ,$$

and

$$S(V) = \{1/2\} \times S(V) \subset \Sigma S(V)^+ .$$

The homeomorphism $\Sigma S(V)^+/S(V) \cong V^\infty \vee_0 V^\infty$ is defined by

$$\Sigma S(V)^+/S(V) \cong V^\infty \vee_0 V^\infty ; (t, u) \mapsto \begin{cases} [1-2t, u]_1 & \text{if } 0 \leq t \leq 1/2 \\ [2t-1, u]_2 & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

(ii) Define

$$\begin{aligned} (\mathbb{R} \times S(V))^\infty &\cong (V \setminus \{0\})^\infty ; (x, u) \mapsto e^x u , \\ (\mathbb{R} \times S(V))^\infty &\cong \Sigma S(V)^+ ; (x, u) \mapsto \left(\frac{e^x}{1+e^x}, u \right) . \end{aligned}$$

(iii) Let $e : V \times \{1, 2\} \hookrightarrow V$ be an open embedding such that

1. $0 < \|e(0, 1)\| < 1 < \|e(0, 2)\|$,
2. $e(V \times \{1, 2\}) \cap (\{0\} \cup S(V)) = \emptyset$,
3. $e| : V \times \{1\} \hookrightarrow V$ is orientation-reversing,
4. $e| : V \times \{2\} \hookrightarrow V$ is orientation-preserving.

The compactification Umkehr of e is a difference map

$$F = \overline{\nabla}_V : V^\infty \rightarrow V^\infty \vee V^\infty$$

such that

1. $F(0) = F(\infty) = \infty$,
2. $F(e(0, i)) = 0_i$ ($i = 1, 2$),
3. the induced map

$$[F] : V^\infty/S(V) \rightarrow V^\infty \vee V^\infty$$
 is a homotopy equivalence,

4. the induced map

$$[F] : (V^\infty/S(V))/\{e(0, 1) \sim e(0, 2)\} \rightarrow V^\infty \vee_0 V^\infty$$

is a homotopy equivalence.

The composite

$$i\overline{\nabla}_V : V^\infty \rightarrow (V^\infty/S(V))/\{e(0,1) \sim e(0,2)\} \xrightarrow{[F]} V^\infty \vee_0 V^\infty$$

is homotopic to

$$V^\infty \rightarrow (V^\infty/S(V))/\{0 \sim \infty\} = (V^\infty/0^+)/S(V) \cong V^\infty \vee_0 V^\infty$$

which is just β_V .

(iv) By construction.

□

Definition 2.18. Let $p, q : V^\infty \wedge X \rightarrow Y$ be maps which agree on $X = 0^+ \wedge X \subset V^\infty \wedge X$, i.e. such that

$$p(0, x) = q(0, x) \in Y \quad (x \in X) .$$

(i) The *difference map* of p and q is the map

$$p - q = (q \vee p)(\overline{\nabla}_V \wedge 1_X) : V^\infty \wedge X \rightarrow Y ,$$

that is

$$p - q : V^\infty \wedge X \rightarrow Y ; ([t, u], x) \mapsto \begin{cases} q([1 - 2t, u], x) & \text{if } 0 \leq t \leq 1/2 \\ p([2t - 1, u], x) & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

(ii) The *relative difference of stable maps*, $\delta(p, q)$ of p and q is the rel X difference

$$\delta(p, q) : \Sigma S(V)^+ \wedge X \rightarrow Y ; (t, u, x) \mapsto \begin{cases} q([1 - 2t, u], x) & \text{if } 0 \leq t \leq 1/2 \\ p([2t - 1, u], x) & \text{if } 1/2 \leq t \leq 1 , \end{cases}$$

that is

$$\delta(p, q) = (q \vee_0 p)(\gamma_V \wedge 1_X) : \Sigma S(V)^+ \wedge X \rightarrow (V^\infty \vee_0 V^\infty) \wedge X \rightarrow Y .$$

□

Example 2.19. The relative difference of maps $p, q : V^\infty \wedge X \rightarrow Y$ such that

$$p(0, x) = q(0, x) = * \in Y \quad (x \in X)$$

is just the difference of the induced maps

$$[p], [q] : \Sigma S(V)^+ \wedge X = (V^\infty/0^+) \wedge X \rightarrow Y ,$$

that is

$$\delta(p, q) = [p] - [q] : \Sigma S(V)^+ \wedge X \rightarrow Y ,$$

with $[p] - [q] = ([q] \vee [p])\bar{\nabla}$ defined using the difference map

$$\bar{\nabla} : \Sigma S(V)^+ \rightarrow \Sigma S(V)^+ \vee \Sigma S(V)^+ ; (t, u) \mapsto \begin{cases} (1 - 2t, u) & \text{if } 0 \leq t \leq 1/2 \\ (2t - 1, u) & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

□

The difference map $p - q$ of Definition 2.18 (i) is just an explicit representative of the difference of the homotopy classes $p, q \in [V^\infty \wedge X, Y]$. Writing $[t, u] = v \in V^\infty$ the difference of $p, q : V^\infty \wedge X \rightarrow Y$ can be expressed as

$$p - q : V^\infty \wedge X \rightarrow Y ; (v, x) \mapsto \begin{cases} q(\frac{1 - \|v\|}{2\|v\|^2}v, x) & \text{if } 0 < \|v\| < 1 \\ p(0, x) = q(0, x) & \text{if } \|v\| = 1 \\ p(\frac{\|v\| - 1}{2\|v\|}v, x) & \text{if } \|v\| > 1 \\ \infty & \text{if } v = 0 \text{ or } * . \end{cases}$$

The difference is the composite

$$p - q : V^\infty \wedge X \rightarrow (V^\infty/0^+) \wedge X \cong \Sigma S(V)^+ \wedge X \xrightarrow{\delta(p, q)} Y ,$$

and the same formulae apply to the relative difference :

$$\delta(p, q) : (V^\infty/0^+) \wedge X \rightarrow Y ; (v, x) \mapsto \begin{cases} q(\frac{1 - \|v\|}{2\|v\|^2}v, x) & \text{if } 0 < \|v\| \leq 1 \\ p(\frac{\|v\| - 1}{2\|v\|}v, x) & \text{if } \|v\| \geq 1 \\ \infty & \text{if } v = 0 \text{ or } * . \end{cases}$$

The relative difference of maps $p, q : V^\infty \wedge X \rightarrow Y$ which agree on $0^\infty \wedge X$ can be interpreted as the difference of two null-homotopies:

Proposition 2.20. *Let $p, q : V^\infty \wedge X \rightarrow Y$ be pointed maps such that*

$$p(0, x) = q(0, x) \in Y \quad (x \in X) .$$

(i) *The pointed map defined by*

$$f : S(V)^+ \wedge X \rightarrow Y ; (u, x) \mapsto p(0, x) = q(0, x)$$

has two null-homotopies

$$\begin{aligned} \delta p & : CS(V)^+ \wedge X \rightarrow Y ; (t, v, x) \mapsto p([t, v], x) , \\ \delta q & : CS(V)^+ \wedge X \rightarrow Y ; (t, v, x) \mapsto q([t, v], x) \end{aligned}$$

and the relative difference of p, q is given by

$$\begin{aligned} \delta(p, q) & = -\delta q \cup \delta p : \\ \Sigma S(V)^+ \wedge X & = CS(V)^+ \wedge X \cup_{S(V)^+ \wedge X} CS(V)^+ \wedge X \rightarrow Y . \end{aligned}$$

(ii) Under the one-one correspondence of Proposition 2.11 (ii) between pointed maps $f : \Sigma S(V)^+ \wedge X \rightarrow Y$ and pairs (g, h) defined by a pointed map $g : S(U)^+ \wedge V^\infty \wedge X \rightarrow Y$ and a null-homotopy

$$h : g(j_{U,V} \wedge 1_X) \simeq \{*\} : S(U \oplus V)^+ \wedge X \rightarrow Y$$

the relative difference $f = \delta(p, q)$ corresponds to

$$\begin{aligned} g & = \delta(p, q)(k_{U,V} \wedge 1_X) = (p - q)(s_U \wedge 1_{V^\infty \wedge X}) : S(U)^+ \wedge V^\infty \wedge X \rightarrow Y , \\ h & = \delta(p, q)(\delta j_{U,V} \wedge 1_X) : C(S(U \oplus V))^+ \wedge X \rightarrow Y . \end{aligned}$$

Proof. (i) Immediate from Proposition 2.10 and Definition 1.3.

(ii) By construction. \square

A linear map $f : V \rightarrow W$ of inner product spaces is proper if and only if f is injective, in which case there is defined an injective pointed map $f^\infty : V^\infty \rightarrow W^\infty$ of the one-point compactifications.

Example 2.21. (i) Let $f, g : V \rightarrow W$ be injective linear maps of inner product spaces V, W , and let $X = S^0$, $Y = W^\infty$ in 2.20. The difference and relative difference maps of the pointed maps

$$f^\infty , g^\infty : V^\infty \wedge X = V^\infty \rightarrow Y = W^\infty$$

are

$$\begin{aligned} f^\infty - g^\infty & = (g^\infty \vee f^\infty) \overline{\nabla}_V : V^\infty \rightarrow W^\infty , \\ \delta(f^\infty, g^\infty) & = (g^\infty \vee f^\infty) \gamma_V : \Sigma S(V)^+ \rightarrow W^\infty . \end{aligned}$$

(ii) For $f = -1$, $g = 1 : V \rightarrow W = V$ (i) gives the difference and relative difference maps of the maps

$$p = 1^\infty = 1 , q = (-1)^\infty = -1 : V^\infty \wedge X = V^\infty \rightarrow Y = V^\infty$$

to be

$$\begin{aligned} p - q &= ((-1)^\infty \vee 1^\infty) \overline{\nabla}_V \simeq 2 : V^\infty \rightarrow V^\infty, \\ \delta(p, q) &= ((-1)^\infty \vee 1^\infty) \gamma_V : \Sigma S(V)^+ \rightarrow V^\infty. \end{aligned}$$

□

Example 2.22. Let $X = S^0$. The difference and relative difference maps of the maps

$$p, q : V^\infty \wedge X = V^\infty \rightarrow Y = V^\infty \vee_0 V^\infty$$

defined by

$$p(v) = v_2, \quad q(v) = v_1 \quad (v \in V).$$

are

$$\begin{aligned} p - q &= \beta_V : V^\infty \rightarrow V^\infty \vee_0 V^\infty, \\ \delta(p, q) &= \gamma_V : \Sigma S(V)^+ \rightarrow V^\infty \vee_0 V^\infty. \end{aligned}$$

□

Proposition 2.23. (i) *For any pointed spaces X, Y and inner product space V there is defined an exact sequence of pointed sets*

$$\begin{aligned} \dots \longrightarrow [\Sigma X, Y] &\xrightarrow{\Sigma s_V^*} [\Sigma S(V)^+ \wedge X, Y] \xrightarrow{\alpha_V^*} [V^\infty \wedge X, Y] \\ &\xrightarrow{0_V^*} [X, Y] \xrightarrow{s_V^*} [S(V)^+ \wedge X, Y] \end{aligned}$$

with

$$0_V^* : [V^\infty \wedge X, Y] \rightarrow [0^+ \wedge X, Y] = [X, Y] ; F \mapsto F|_{0^+ \wedge X}.$$

For any maps $p, q : V^\infty \wedge X \rightarrow Y$ which agree on $0^+ \wedge X$ the difference $p - q \in [V^\infty \wedge X, Y]$ has image $$ in $[X, Y]$, and $p - q$ is the image of $\delta(p, q) \in [\Sigma S(V)^+ \wedge X, Y]$, with a homotopy commutative diagram*

$$\begin{array}{ccc} V^\infty \wedge X & \xrightarrow{\alpha_V \wedge 1_X} & \Sigma S(V)^+ \wedge X \\ & \searrow p - q & \downarrow \delta(p, q) \\ & & Y \end{array}$$

that is

$$\begin{aligned} p - q &= \alpha_V^*(\delta(p, q)) \in \ker(0_V^* : [V^\infty \wedge X, Y] \rightarrow [X, Y]) \\ &= \text{im}(\alpha_V^* : [\Sigma S(V)^+ \wedge X, Y] \rightarrow [V^\infty \wedge X, Y]). \end{aligned}$$

(ii) *If the maps $p, q : V^\infty \wedge X \rightarrow Y$ in (i) are related by a homotopy*

$$r : p \simeq q : V^\infty \wedge X \rightarrow Y$$

then

$$p - q \simeq * : V^\infty \wedge X \rightarrow Y$$

and the relative difference $\delta(p, q) \in [\Sigma S(V)^+ \wedge X, Y]$ is the image of the map

$$r_0 : \Sigma X \rightarrow Y ; (t, x) \mapsto r(t, 0, x) ,$$

with a homotopy commutative diagram

$$\begin{array}{ccc} \Sigma S(V)^+ \wedge X & \xrightarrow{\Sigma s_V \wedge 1_X} & S^1 \wedge X = \Sigma X \\ & \searrow \delta(p, q) & \downarrow r_0 \\ & & Y \end{array}$$

i.e. if $p = q \in [V^\infty \wedge X, Y]$ then

$$\begin{aligned} \alpha_V^*(\delta(p, q)) &= p - q = * \in [V^\infty \wedge X, Y] , \\ \delta(p, q) &= \Sigma s_V^*(r_0) \in \ker(\alpha_V^* : [\Sigma S(V)^+ \wedge X, Y] \rightarrow [V^\infty \wedge X, Y]) \\ &= \text{im}(\Sigma s_V^* : [\Sigma X, Y] \rightarrow [\Sigma S(V)^+ \wedge X, Y]) . \end{aligned}$$

In particular, if

$$r(t, 0, x) = p(0, x) = q(0, x) \in Y \quad (t \in I, x \in X)$$

then $\delta(p, q) \simeq * : \Sigma S(V)^+ \wedge X \rightarrow Y$, i.e. if p, q are related by a homotopy which is constant on $\{0\} \times X$ then

$$\delta(p, q) = * \in [\Sigma S(V)^+ \wedge X, Y] .$$

(iii) If $p, p', q, q' : V^\infty \wedge X \rightarrow Y$ are maps which agree on $0^+ \wedge X$ and

$$f : p \simeq p' , g : q \simeq q' : V^\infty \wedge X \rightarrow Y$$

are homotopies which are constant on $0^+ \wedge X$ there is induced a homotopy of the relative differences

$$\delta(f, g) : \delta(p, q) \simeq \delta(p', q') : \Sigma S(V)^+ \wedge X \rightarrow Y .$$

(iv) If $p, q, r : V^\infty \wedge X \rightarrow Y$ are maps which agree on $0^+ \wedge X$ the relative differences are related by a homotopy

$$h : \delta(p, q) + \delta(q, r) \simeq \delta(p, r) : \Sigma S(V)^+ \wedge X \rightarrow Y .$$

(v) If $p, q : V^\infty \wedge X \rightarrow Y$ are maps such that

$$p(t, x) = \begin{cases} q(t, x) & \text{if } (t, x) \in U \\ * & \text{if } (t, x) \in \overline{(V^\infty \wedge X) \setminus U} \end{cases}$$

for some neighbourhood $U \subseteq V^\infty \wedge X$ of $0^+ \wedge X \subset V^\infty \wedge X$ there exists a homotopy

$$\gamma(p, q) : \delta(p, q) \simeq q' : \Sigma S(V)^+ \wedge X \rightarrow Y$$

with q' defined by

$$q' : \Sigma S(V)^+ \wedge X = (V^\infty / 0^+) \wedge X \rightarrow Y ;$$

$$(t, x) \mapsto \begin{cases} * & \text{if } (t, x) \in U \\ q(t, x) & \text{if } (t, x) \in \overline{V^\infty \wedge X \setminus U} . \end{cases}$$

(vi) The homotopy $\gamma(p, q)$ in (v) can be chosen to be natural, meaning that given commutative squares of maps

$$\begin{array}{ccc} V^\infty \wedge X_1 & \xrightarrow{p_1} & Y_1 \\ 1 \wedge h \downarrow & & \downarrow k \\ V^\infty \wedge X_2 & \xrightarrow{p_2} & Y_2 \end{array} , \quad \begin{array}{ccc} V^\infty \wedge X_1 & \xrightarrow{q_1} & Y_1 \\ 1 \wedge h \downarrow & & \downarrow k \\ V^\infty \wedge X_2 & \xrightarrow{q_2} & Y_2 \end{array}$$

and neighbourhoods $U_i \subseteq V^\infty \wedge X_i$ of $0^+ \wedge X_i \subset V^\infty \wedge X_i$ with $U_2 = (1 \wedge h)(U_1)$ such that

$$p_i(t, x) = \begin{cases} q_i(t, x) & \text{if } (t, x) \in U_i \\ * & \text{if } (t, x) \in \overline{V^\infty \wedge X_i \setminus U_i} \end{cases}$$

there is defined a commutative square

$$\begin{array}{ccc} I \times (\Sigma S(V)^+ \wedge X_1) & \xrightarrow{\gamma(p_1, q_1)} & Y_1 \\ 1 \wedge h \downarrow & & \downarrow k \\ I \times (\Sigma S(V)^+ \wedge X_2) & \xrightarrow{\gamma(p_2, q_2)} & Y_2 \end{array}$$

Proof. (i) Immediate from the cofibration sequence of pointed spaces

$$S(V)^+ \xrightarrow{s_V} S^0 = 0^+ \rightarrow V^\infty \xrightarrow{\alpha_V} \Sigma S(V)^+ \xrightarrow{\Sigma s_V} S^1 \rightarrow \dots$$

given by Proposition 2.17 (i).

(ii) Immediate from the homotopy $\beta_V \simeq i\bar{\nabla}_V$ given by Proposition 2.17 (iii).

(iii) The homotopies

$$f, g : I \times (V^\infty \wedge X) \rightarrow Y$$

which agree on $I \times 0^+ \wedge X$. The relative difference

$$\delta(f, g) : I \times (\Sigma S(V)^+ \wedge X) \rightarrow Y .$$

defines a homotopy $\delta(f, g) : \delta(p, q) \simeq \delta(p', q')$.

(iv) The sum of $\delta(p, q)$ and $\delta(q, r)$ is defined by

$$\begin{aligned} \delta(p, q) + \delta(q, r) &: \Sigma S(V)^+ \wedge X \rightarrow Y ; \\ (t, u, x) &\mapsto \begin{cases} \delta(q, r)(2t, u, x) & \text{if } 0 \leq t \leq 1/2 \\ \delta(p, q)(2t - 1, u, x) & \text{if } 1/2 \leq t \leq 1 . \end{cases} \end{aligned}$$

Define a homotopy $h : \delta(p, q) + \delta(q, r) \simeq \delta(p, r)$ by

$$h(s, t, u, x) = \begin{cases} r([\frac{1+s-4t}{1+s}, u], x) & \text{if } 0 \leq t \leq (1+s)/4 \\ q([4t-1-s, u], x) & \text{if } (1+s)/4 \leq t \leq 1/2 \\ q([3-s-4t, u], x) & \text{if } 1/2 \leq t \leq (3-s)/4 \\ p([\frac{s+4t-3}{1+s}, u], x) & \text{if } (3-s)/4 \leq t \leq 1 . \end{cases}$$

(v)+(vi) These are special cases of Proposition 1.7. □

Example 2.24. (i) For any inner product spaces V, W and injective linear maps $f, g : V \rightarrow W$ the maps $f^\infty, g^\infty : V^\infty \rightarrow W^\infty$ are such that

$$f^\infty(0_V) = g^\infty(0_V) = 0_W \in W^\infty ,$$

so that there is defined a relative difference map

$$\delta(f^\infty, g^\infty) : \Sigma S(V)^+ \rightarrow W^\infty ; [t, u] \mapsto \begin{cases} g([1-2t, u]) & \text{if } 0 \leq t \leq 1/2 \\ f([2t-1, u]) & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

By Proposition 2.23 (i) there exists a homotopy

$$\delta(f^\infty, g^\infty)\alpha_V \simeq f^\infty - g^\infty : V^\infty \rightarrow W^\infty .$$

(ii) Let $GL(V)$ be the space of linear automorphisms $a : V \rightarrow V$, with base point $1 : V \rightarrow V$. For any map $c : X \rightarrow GL(V)$ the maps defined by

$$\begin{aligned} p &= \text{adj}(1) : V^\infty \wedge X \rightarrow V^\infty ; (v, x) \mapsto v , \\ q &= \text{adj}(c) : V^\infty \wedge X \rightarrow V^\infty ; (v, x) \mapsto c(x)(v) \end{aligned}$$

are such that $p(0, x) = q(0, x) = 0 \in V^\infty$ ($x \in V$). Use the relative difference construction of (i) to define a function

$$\delta_V : [X, GL(V)] \rightarrow [\Sigma S(V)^+ \wedge X, V^\infty] ; c \mapsto \delta_V(c) = \delta(p, q)$$

with

$$\delta_V(c)([t, u], x) = \begin{cases} c(x)^\infty [1 - 2t, u] & \text{if } 0 \leq t \leq 1/2 \\ [2t - 1, u] & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

The composite

$$[X, GL(V)] \xrightarrow{\delta_V} [\Sigma S(V)^+ \wedge X, V^\infty] \xrightarrow{\alpha_V^*} [V^\infty \wedge X, V^\infty]$$

is given by

$$\alpha_V^* \delta_V : [X, GL(V)] \rightarrow [V^\infty \wedge X, V^\infty] ; c \mapsto p - q .$$

□

Example 2.25. Here are two special cases of Proposition 2.23 (v).

(i) For any map $p : V^\infty \wedge X \rightarrow Y$ take $q = p$, $U = V^\infty \wedge X$ to obtain a homotopy

$$\gamma(p, p) : \delta(p, p) \simeq * : \Sigma S(V)^+ \wedge X \rightarrow Y .$$

(ii) For any map $q : V^\infty \wedge X \rightarrow Y$ such that

$$q(0, x) = * \in Y \quad (x \in X)$$

take $q = *$, $U = 0^+ \wedge X$ to obtain a homotopy

$$\gamma(p, *) : \delta(p, *) \simeq [p] : \Sigma S(V)^+ \wedge X \rightarrow Y .$$

with $[p]$ induced from p using the homeomorphism $\Sigma S(V)^+ \cong V^\infty / 0^+$. □

Proposition 2.26. *The relative difference (2.18 (ii)) of maps $p, q : V^\infty \wedge X \rightarrow Y$ such that $p|_X = q|_X : X \rightarrow Y$ is the relative difference (1.3) of the null-homotopies $\delta p, \delta q$ of*

$$(p|_X)(s_V \wedge 1_X) = (q|_X)(s_V \wedge 1_X) : S(V)^+ \wedge X \rightarrow Y$$

given by 2.10, that is

$$\begin{aligned} \delta(p, q) : \Sigma S(V)^+ \wedge X &\xrightarrow{(-1 \cup 1) \wedge 1_X} \\ CS(V)^+ \wedge X \cup_{S(V)^+ \wedge X} CS(V)^+ \wedge X &\xrightarrow{\delta q \cup \delta p} Y \end{aligned}$$

with

$$\delta p(t, u, x) = p([t, u], x) , \quad \delta q(t, u, x) = q([t, u], x) .$$

Proof. For all $t \in I$, $u \in S(V)$, $x \in X$

$$\delta(p, q)(t, u, x) = \begin{cases} q(1 - 2t, u, x) & \text{if } 0 \leq t \leq 1/2 \\ p(2t - 1, u, x) & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

□

Chapter 3

Stable homotopy theory

This Chapter develops stable homotopy theory and bordism theory using inner product spaces.

3.1 Stable maps

A *stable map* $F : X \rightarrow Y$ between pointed spaces X, Y is a map of the type

$$F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$$

for some (finite-dimensional) inner product space V .

The *stable homotopy group* is the abelian group

$$\{X; Y\} = \varinjlim_V [V^\infty \wedge X, V^\infty \wedge Y]$$

with the direct limit is taken over all (finite-dimensional) inner product spaces V , and addition and subtraction maps induced by the sum and difference maps $\nabla_V, \overline{\nabla}_V : V^\infty \rightarrow V^\infty \vee V^\infty$. For reasonable spaces X, Y (e.g. finite *CW* complexes) an element $F \in \{X; Y\}$ is an equivalence class of stable maps $F : X \rightarrow Y$.

Example 3.1. For an open embedding $e : V \times M \hookrightarrow V \times N$ the adjunction Umkehr $F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge M^+$ defines a stable map $F : N^+ \rightarrow M^+$. \square

Given a map $F : A \rightarrow B$ let $G : B \rightarrow \mathcal{C}(F)$ be the inclusion in the mapping cone, and let $H : \mathcal{C}(F) \rightarrow \Sigma A$ be the projection, so that

$$A \xrightarrow{F} B \xrightarrow{G} \mathcal{C}(F) \xrightarrow{H} \Sigma A \xrightarrow{\Sigma F} \Sigma B \longrightarrow \dots$$

is a cofibration sequence, as before. By analogy with the cohomotopy Barratt-Puppe exact sequence (1.9) there is a stable homotopy exact sequence :

Proposition 3.2. *For any pointed space X there is induced a Barratt-Puppe exact sequence of stable homotopy groups*

$$\begin{aligned} \dots \longrightarrow \{\Sigma X; \mathcal{C}(F)\} &\xrightarrow{H} \{X; A\} \xrightarrow{F} \{X; B\} \\ &\xrightarrow{G} \{X; \mathcal{C}(F)\} \xrightarrow{H} \{X; \Sigma A\} \longrightarrow \dots \end{aligned}$$


□

The cofibration sequence

$$S(V)^+ \xrightarrow{s_V} S^0 \xrightarrow{0_V} V^\infty \xrightarrow{\alpha_V} \Sigma S(V)^+ \longrightarrow \dots$$

determines the following braid :

Proposition 3.3. *For any inner product spaces U, V and pointed spaces X, Y there is defined a commutative braid of exact sequences of stable homotopy groups*



with

$$\begin{aligned}
A_1 &= \{\Sigma X; V^\infty \wedge Y\}, \quad A_2 = \{\Sigma S(U \oplus V)^+ \wedge X; V^\infty \wedge Y\}, \\
A_3 &= \{S(U \oplus V)^+ \wedge X; V^\infty \wedge Y\}, \quad A_4 = \{\Sigma S(U)^+ \wedge X; Y\}.
\end{aligned}$$

Proof. These are the Barratt-Puppe exact sequences (3.2) determined by the commutative braid of cofibrations

$$\begin{array}{ccccc}
& & U^\infty \wedge S(V)^+ & & \Sigma S(U)^+ \\
& \nearrow & \searrow^{1 \wedge s_V} & \nearrow^{\alpha_V} & \searrow \\
S(U \oplus V)^+ & & U^\infty \wedge S^0 = U^\infty & & \Sigma S(U \oplus V)^+ \\
& \searrow^{s_{U \oplus V}} & \nearrow^{0_U} & \searrow^{1 \wedge 0_V} & \nearrow^{\alpha_{U \oplus V}} \\
& & S^0 & & U^\infty \wedge V^\infty = (U \oplus V)^\infty \\
& & \nearrow^{0_{U \oplus V}} & & \\
& & & &
\end{array}$$

given by Proposition 2.17 (iv).

□

We shall need the following version of exactness at $\{X; A\}$ in the Barratt-Puppe sequence (3.2)

$$\begin{aligned}
\ldots \longrightarrow \{\Sigma X; \mathcal{C}(F)\} &\xrightarrow{H} \{X; A\} \xrightarrow{F} \{X; B\} \\
&\xrightarrow{G} \{X; \mathcal{C}(F)\} \xrightarrow{H} \{X; \Sigma A\} \longrightarrow \ldots
\end{aligned}$$

Let X be a pointed space with a difference map $\overline{\nabla} : X \rightarrow X \vee X$, e.g. the co- H -space $X = V^\infty \wedge X_0$ for some non-zero inner product space V and a pointed space X_0 . The difference of maps $f_1, f_2 : X \rightarrow Y$ is defined by

$$f_1 - f_2 = (f_2 \vee f_1) \overline{\nabla} : X \rightarrow Y,$$

and for any map $f : X \rightarrow Y$ there is defined a standard null-homotopy

$$f - f \simeq * : X \rightarrow Y.$$

Definition 3.4. Let X be a pointed space with a difference map $\overline{\nabla} : X \rightarrow X \vee X$. Given maps $a_1, a_2 : X \rightarrow A$, $F : A \rightarrow B$ and a homotopy $b : Fa_1 \simeq Fa_2 : X \rightarrow B$ let

$$\bar{b} : Fa_2 - Fa_1 \simeq * : X \rightarrow B$$

be the null-homotopy of $Fa_2 - Fa_1 = F(a_2 - a_1) : X \rightarrow B$ defined by the concatenation of the homotopy

$$b - 1 : Fa_2 - Fa_1 \simeq Fa_1 - Fa_1 : X \rightarrow B$$

and the standard null-homotopy $Fa_1 - Fa_1 \simeq * : X \rightarrow B$. The *stable relative difference*

$$\delta(a_1, a_2, b) = c \in \{\Sigma X; \mathcal{C}(F)\}$$

is the stable homotopy class of the map $c : \Sigma X \rightarrow \mathcal{C}(F)$ in the map of cofibrations

$$\begin{array}{ccccccc} X & \longrightarrow & CX & \longrightarrow & \Sigma X & \xlongequal{\quad} & \Sigma X \\ a_1 - a_2 \downarrow & & \downarrow \bar{b} & & \downarrow c & & \downarrow \Sigma(a_1 - a_2) \\ A & \xrightarrow{F} & B & \xrightarrow{G} & \mathcal{C}(F) & \xrightarrow{H} & \Sigma A \end{array}$$

□

Proposition 3.5. *The stable relative difference $\delta(a_1, a_2, b) \in \{\Sigma X; \mathcal{C}(F)\}$ of 3.4 is such that*

$$\begin{aligned} a_1 - a_2 &= H(\delta(a_1, a_2, b)) \in \ker(F : \{X; A\} \rightarrow \{X; B\}) \\ &= \text{im}(H : \{\Sigma X; \mathcal{C}(F)\} \rightarrow \{X; A\}) \subseteq \{X; A\} = \{\Sigma X; \Sigma A\} . \end{aligned}$$

Proof. By construction

□

The cofibration sequence

$$S(V)^+ \wedge Y \xrightarrow{s_V} Y \xrightarrow{0_V} V^\infty \wedge Y \longrightarrow \mathcal{C}(0_V) \simeq \Sigma S(V)^+ \wedge Y \longrightarrow \dots$$

induces a long exact sequence of stable homotopy groups

$$\begin{aligned} \dots &\longrightarrow \{V^\infty \wedge X; S(V)^+ \wedge Y\} \xrightarrow{s_V} \{V^\infty \wedge X; Y\} \\ &\xrightarrow{0_V} \{V^\infty \wedge X; V^\infty \wedge Y\} = \{X; Y\} \longrightarrow \{V^\infty \wedge X; \Sigma S(V)^+ \wedge Y\} \longrightarrow \dots \end{aligned}$$

Definition 3.6. Given a map $p : V^\infty \wedge X \rightarrow Y$ define the map

$$f = 0_V p : V^\infty \wedge X \rightarrow V^\infty \wedge Y ; (v, x) \mapsto (0, p(v, x)) .$$

Given also a map $q : V^\infty \wedge X \rightarrow Y$ which agrees with p on $0^+ \wedge X \subset V^\infty \wedge X$ let $g = 0_V q : V^\infty \wedge X \rightarrow V^\infty \wedge Y$, and define a homotopy

$$h : f \simeq g : V^\infty \wedge X \rightarrow V^\infty \wedge Y$$

by

$$h : I \times V^\infty \wedge X \rightarrow V^\infty \wedge Y ;$$

$$(t, v, x) \mapsto \begin{cases} (2tv, q((1-2t)v, x)) & \text{if } 0 \leq t \leq 1/2 \\ ((2-2t)v, p((2t-1)v, x)) & \text{if } 1/2 \leq t \leq 1 . \end{cases}$$

The *stable relative difference* of p, q is the stable relative difference (3.4)

$$\delta'(p, q) = \delta(f, g, h) \in \{\Sigma(V^\infty \wedge X); \mathcal{C}(0_V)\} = \{V^\infty \wedge X; S(V)^+ \wedge Y\} .$$

□

Proposition 3.7. *The stable relative difference of 3.6 is such that*

$$\begin{aligned} p - q &= s_V \delta'(p, q) \in \text{im}(s_V : \{V^\infty \wedge X; S(V)^+ \wedge Y\} \rightarrow \{V^\infty \wedge X; Y\}) \\ &= \ker(0_V : \{V^\infty \wedge X; Y\} \rightarrow \{X; Y\}) . \end{aligned}$$

Proof. Immediate from Proposition 3.5.

□

The Umkehr maps of open embeddings $V \times X \hookrightarrow V \times Y$ are stable maps. Specifically, given a map $f : X \rightarrow Y$, an inner product space V and a map $g : V \times X \rightarrow Y$ such that

$$e = (g, f) : V \times X \rightarrow V \times Y ; (v, x) \mapsto (g(v, x), f(x))$$

is an open embedding then Definition 2.3 gives a compactification Umkehr stable map

$$F : (V \times Y)^\infty = V^\infty \wedge Y^\infty \rightarrow (V \times X)^\infty = V^\infty \wedge X^\infty .$$

For compact X, Y this is an adjunction Umkehr stable map $F : V^\infty \wedge Y^+ \rightarrow V^\infty \wedge X^+$.

Example 3.8. For any finite cover $f : \tilde{K} \rightarrow K$ of a CW complex K there is defined a transfer map

$$f^! : C(K) \rightarrow C(\tilde{K}) ; x \mapsto \sum_{y \in f^{-1}(x)} y .$$

For finite K there exist a finite-dimensional inner product space V and a map $g : V \times \tilde{K} \rightarrow V$ such that

$$e = (g, f) : V \times \tilde{K} \rightarrow V \times K ; (v, x) \mapsto (g(v, x), f(x))$$

is an open embedding, with compactification (= adjunction) Umkehr map

$$F : (V \times K)^\infty = V^\infty \wedge K^+ \rightarrow (V \times \tilde{K})^\infty = V^\infty \wedge \tilde{K}^+$$

inducing the transfer map $F = f^! : C(K) \rightarrow C(\tilde{K})$ (cf. Adams [2, §4.2], Proposition 4.32 and Example 6.29 below).

□

We shall need a version of the Umkehr construction for pointed maps.

Definition 3.9. Let $f : X \rightarrow Y$ be a pointed map, with $f(x_0) = y_0 \in Y$ for base points $x_0 \in X$, $y_0 \in Y$. Let V be an inner product space, and suppose given a map $g : V \times X \rightarrow V$ such that

$$e = (g, f) : V \times X \rightarrow V \times Y ; (v, x) \mapsto (g(v, x), f(x))$$

restricts to an open embedding

$$e| : V \times (X \setminus \{x_0\}) \rightarrow V \times Y$$

and also

$$g(v, x_0) = 0 \in V \quad (v \in V) .$$

The *Umkehr stable map* of e is

$$F : V^\infty \wedge Y \rightarrow V^\infty \wedge X ;$$

$$(w, y) \mapsto \begin{cases} (v, x) & \text{if } (w, y) = (g(v, x), f(x)) \in V \times Y \\ \infty & \text{otherwise .} \end{cases}$$

□

Example 3.10. (i) Let $f : X \rightarrow Y$ be a pointed map of finite CW complexes such that the restriction $f| : X \setminus \{x_0\} \rightarrow Y \setminus \{y_0\}$ is a finite cover, so that there is defined a transfer chain map

$$f^! : \dot{C}(Y) \rightarrow \dot{C}(X) ; y \mapsto \sum_{x \in f^{-1}(y)} x$$

of the reduced singular chain complexes. As in Adams [3, pp. 511–512] there exist a finite-dimensional inner product space V and a map $g : V \times X \rightarrow V$

such that

$$e = (g, f) : V \times X \rightarrow V \times Y ; (v, x) \mapsto (g(v, x), f(x))$$

satisfies the hypothesis of Definition 3.9, giving an Umkehr stable map

$$F : V^\infty \wedge Y \rightarrow V^\infty \wedge X$$

inducing $f^!$ on the chain level.

(ii) If $f : \tilde{K} \rightarrow K$ is a finite cover of a finite CW complex then for any finite-dimensional inner product space W

$$1 \wedge f^+ : X = W^\infty \wedge \tilde{K}^+ \rightarrow Y = W^\infty \wedge K^+$$

is a pointed map as in (i), and the Umkehr stable map

$$F : V^\infty \wedge Y = (V \oplus W)^\infty \wedge K^+ \rightarrow V^\infty \wedge X = (V \oplus W)^\infty \wedge \tilde{K}^+$$

is just the compactification Umkehr map of Example 3.8 above, inducing the transfer chain map

$$F = f^! : \dot{C}(Y) = C(K)_{*-\dim(W)} \rightarrow \dot{C}(X) = C(\tilde{K})_{*-\dim(W)} .$$

□

Definition 3.11. (i) The *stable homotopy* and *cohomotopy* groups of a space X are

$$\omega_n(X) = \{S^n; X^+\} , \omega^n(X) = \{X^+; S^n\} \quad (n \in \mathbb{Z}) .$$

(ii) The *reduced stable homotopy* and *cohomotopy* groups of a space X are

$$\tilde{\omega}_n(X) = \{S^n; X\} , \tilde{\omega}^n(X) = \{X; S^n\} .$$

In particular, for an unpointed space X

$$\omega_n(X) = \tilde{\omega}_n(X^+) , \omega^n(X) = \tilde{\omega}^n(X^+) .$$

□

Example 3.12. The stable homotopy and cohomotopy groups of $\{*\}$ are the stable homotopy groups of spheres

$$\omega_n(\{*\}) = \omega^{-n}(\{*\}) = \{S^n; S^0\} \quad (n \geq 0)$$

which we write as ω_n .

□

Definition 3.13. (i) A *spectrum* $\underline{X} = \{X(V)\}$ is a sequence of pointed spaces $X(V)$ indexed by finite-dimensional inner product spaces V , with structure maps

$$(V^\perp)^\infty \wedge X(V) \rightarrow X(W)$$

defined whenever $V \subseteq W$, where $V^\perp \subseteq W$ is the orthogonal complement of V in W . For $n \in \mathbb{Z}$ let

$$\pi_n(\underline{X}) = \begin{cases} \varinjlim_V [\Sigma^n V^\infty, X(V)] & \text{if } n \geq 0 \\ \varprojlim_V [V^\infty, \Sigma^{-n} X(V)] & \text{if } n \leq -1. \end{cases}$$

(ii) A spectrum \underline{X} is *connective* if

$$\pi_n(\underline{X}) = 0 \text{ for } n \leq -1.$$

In particular, this is the case if each $X(V)$ is $(\dim(V) - 1)$ -connected, i.e.

$$\pi_n(X(V)) = 0 \text{ for } n \leq (\dim(V) - 1).$$

(iii) The \underline{X} -coefficient homology groups of a space Y are

$$\underline{X}_n(Y) = \pi_n(\underline{X} \wedge Y^+) \quad (n \in \mathbb{Z}).$$

The *reduced \underline{X} -coefficient homology* groups of a pointed space Y are

$$\tilde{\underline{X}}_n(Y) = \pi_n(\underline{X} \wedge Y) \quad (n \in \mathbb{Z}).$$

If \underline{X} is connective then $\underline{X}_n(Y) = 0$ for $n \leq -1$, and similarly in the reduced case.

□

Example 3.14. (i) The suspension spectrum of a pointed space X is the connective spectrum

$$\underline{X} = \{V^\infty \wedge X \mid \dim(V) < \infty\},$$

such that for any pointed space Y

$$\underline{X}_n(Y) = \begin{cases} \tilde{\omega}_n(X \wedge Y) & \text{if } n \geq 0 \\ 0 & \text{if } n \leq -1. \end{cases}$$

In particular, for $Y = S^0$

$$\pi_n(\underline{X}) = \underline{X}_n(S^0) = \begin{cases} \tilde{\omega}_n(X) & \text{if } n \geq 0 \\ 0 & \text{if } n \leq -1. \end{cases}$$

(ii) The sphere spectrum $\underline{S} = \{V^\infty\}$ (= the suspension spectrum of S^0) is such that

$$\pi_n(\underline{S}) = \underline{S}_n(S^0) = \omega_n, \quad \underline{S}_n(X) = \tilde{\omega}_n(X).$$

□

3.2 Vector bundles

We shall only be considering vector bundles and spherical fibrations over CW complexes.

Definition 3.15. Let V be an inner product space.

(i) A V -*bundle* over a space X is a vector bundle

$$\xi : V \longrightarrow E(\xi) \xrightarrow{p_\xi} X$$

with each fibre $p^{-1}(x) = \xi_x$ ($x \in X$) an inner product space isomorphic to V , and the transition functions linear isometries.

(ii) The *Thom space* of ξ is the pointed space

$$T(\xi) = D(\xi)/S(\xi),$$

with $(D(\xi), S(\xi))$ the total space of the $(D(V), S(V))$ -bundle of ξ

$$(D(V), S(V)) \longrightarrow (D(\xi), S(\xi)) \xrightarrow{p_\xi|} X.$$

(iii) The *trivial V -bundle* ϵ_V over X has

$$\begin{aligned} p &= \text{projection} : E(\epsilon_V) = V \times X \rightarrow X, \\ D(\epsilon_V) &= D(V) \times X, \quad S(\epsilon_V) = S(V) \times X, \\ T(\epsilon_V) &= (D(V) \times X)/(S(V) \times X) = V^\infty \wedge X^+. \end{aligned}$$

In particular, for $V = 0$

$$D(\epsilon_V) = X, \quad S(\epsilon_V) = \emptyset, \quad T(\epsilon_V) = X^+.$$

(iv) The *Thom spectrum* of a V -bundle ξ is the suspension spectrum of $T(\xi)$

$$\underline{T}(\xi) = \{T(\xi \oplus \epsilon_V) \mid \dim(V) < \infty\}$$

with $T(\xi \oplus \epsilon_V) = V^\infty \wedge T(\xi)$.

□

Remark 3.16. The canonical map

$$E(\xi)^\infty \rightarrow T(\xi) ; (x, v) \mapsto (x, \frac{v}{1 + \|v\|}) \quad (x \in X, v \in V)$$

is a bijection which is a homeomorphism if and only if X is compact.

□

Definition 3.17. (i) The *pullback* of a V -bundle ξ over X along a map $f : Y \rightarrow X$ is the V -bundle over Y

$$f^*\xi : V \longrightarrow E(f^*\xi) \xrightarrow{pf^*\xi} Y$$

with $E(f^*\xi)$ fitting into a pullback square

$$\begin{array}{ccc} E(f^*\xi) & \longrightarrow & E(\xi) \\ pf^*\xi \downarrow & & \downarrow p_\xi \\ Y & \xrightarrow{f} & X \end{array}.$$

(ii) The *product* of a V -bundle over X and a W -bundle over Y

$$\xi : V \longrightarrow E(\xi) \xrightarrow{p_\xi} X \quad , \quad \eta : W \longrightarrow E(\eta) \xrightarrow{p_\eta} Y$$

is the $V \oplus W$ -bundle over $X \times Y$

$$\xi \times \eta : V \oplus W \longrightarrow E(\xi \times \eta) \xrightarrow{p_{\xi \times \eta}} X \times Y$$

with

$$p_{\xi \times \eta} = p_\xi \times p_\eta : E(\xi \times \eta) = E(\xi) \times E(\eta) \rightarrow X \times Y, \quad T(\xi \times \eta) = T(\xi) \wedge T(\eta).$$

(iii) The *Whitney sum* of a V -bundle over X and a W -bundle over X

$$\xi : V \longrightarrow E(\xi) \xrightarrow{p_\xi} X \quad , \quad \eta : W \longrightarrow E(\eta) \xrightarrow{p_\eta} X$$

is the pullback $V \oplus W$ -bundle over X

$$\xi \oplus \eta = \Delta^*(\xi \times \eta) : V \oplus W \longrightarrow E(\xi \oplus \eta) \xrightarrow{p_{\xi \oplus \eta}} X$$

with $\Delta : X \rightarrow X \times X; x \mapsto (x, x)$, such that

$$T((\xi \oplus \eta)^+) = T(\xi^+) \wedge_X T(\eta^+) .$$

The total space $E(\xi \oplus \eta)$ fits into a pullback square

$$\begin{array}{ccc} E(\xi \oplus \eta) & \longrightarrow & E(\xi) \\ \downarrow & \searrow p_{\xi \oplus \eta} & \downarrow p_\xi \\ E(\eta) & \xrightarrow{p_\eta} & X \end{array}$$

and the Thom space of $\xi \oplus \eta$ is

$$T(\xi \oplus \eta) = T(p_{D(\xi)}^* \eta) / T(p_{S(\xi)}^* \eta) .$$

□

Proposition 3.18. *Let*

$$\xi : V \rightarrow E(\xi) \rightarrow M, \quad \eta : W \rightarrow E(\eta) \rightarrow N$$

be vector bundles. Regard $E(\xi)$ as a subspace of the disk space $D(\xi)$ via the open embedding

$$E(\xi) \hookrightarrow D(\xi) ; (m, v) \mapsto (m, \frac{v}{1 + \|v\|}) \quad (m \in M, v \in V)$$

with $D(\xi) \setminus E(\xi) = S(\xi)$, and similarly for $E(\eta) \subset D(\eta)$. If

$$f : X = E(\xi) \hookrightarrow Y = E(\eta)$$

is an open embedding which extends to an embedding $\bar{f} : D(\xi) \hookrightarrow D(\eta)$ there is defined a homeomorphism

$$T(\xi) = D(\xi)/S(\xi) \rightarrow Y/(Y \setminus f(X)) ; x \mapsto \bar{f}(x)$$

and the adjunction Umkehr of \bar{f} with respect to $E(\eta) \subset D(\eta)$ is a map

$$F : D(\eta)/S(\eta) = T(\eta) \rightarrow Y/(Y \setminus f(X)) = T(\xi) .$$

Proof. By construction.

□

Example 3.19. Let V, W be finite dimensional inner product spaces. For an open embedding of the type

$$e : E(\epsilon_W) = W \times M \hookrightarrow E(\epsilon_V) = V \times N$$

which extends to an embedding

$$f : D(\epsilon_W) = D(W) \times M \hookrightarrow D(\epsilon_V) = D(V) \times N$$

the adjunction Umkehr of 3.18 is a map

$$F : T(\epsilon_V) = V^\infty \wedge N^+ \rightarrow T(\epsilon_W) = W^\infty \wedge M^+ .$$

□

Given a (topological) group G with a right action on a space X and a left action on a space Y let $X \times_G Y$ be the quotient of $X \times Y$ by the equivalence relation

$$(xg, y) \sim (x, gy) \text{ for all } g \in G, x \in X, y \in Y .$$

Similarly for pointed spaces, with $X \wedge_G Y$ the corresponding quotient of $X \wedge Y$.

Definition 3.20. Let V, W be inner product spaces.

(i) The *orthogonal group* $O(V)$ is the group of linear isometries $g : V \rightarrow V$, regarded as a pointed space with base point $1 : V \rightarrow V$. Define a left $O(V)$ -action on V by

$$O(V) \times V \rightarrow V ; (g, x) \mapsto g(x) .$$

(ii) Let $O(V, W)$ be the *Stiefel manifold* of linear isometries $i : V \hookrightarrow W$. Define a right $O(V)$ -action on $O(V, W)$ by

$$O(V, W) \times O(V) \rightarrow O(V, W) ; (i, g) \mapsto ig .$$

(iii) Let $G(V, W)$ be the *Grassmann manifold* of the subspaces $U \subseteq W$ which are isomorphic to V .

□

Here are some standard properties of the Stiefel and Grassmann manifolds:

Proposition 3.21. (i) For any finite-dimensional inner product space W and subspace $V \subseteq W$ there is defined a homeomorphism

$$G(V, W) \rightarrow G(V^\perp, W) ; U \mapsto U^\perp .$$

(ii) For any two inner product spaces U, V $O(U) \times O(V)$ has a right action on $O(U \oplus V)$

$$O(U \oplus V) \times (O(U) \times O(V)) \rightarrow O(U \oplus V) ; (f, g, h) \mapsto f(g \oplus h) .$$

Regard $O(V, U \oplus V)$ as a pointed space with base point the inclusion

$$j : V \hookrightarrow U \oplus V ; x \mapsto (0, x) .$$


The homeomorphisms

$$\begin{aligned} O(U \oplus V)/O(V) &\rightarrow O(U, U \oplus V) ; h \mapsto h|_U , \\ O(U \oplus V)/O(U) &\rightarrow O(V, U \oplus V) ; h \mapsto h|_V , \\ O(U \oplus V)/(O(U) \times O(V)) &\rightarrow G(V, U \oplus V) ; h \mapsto h(V) \end{aligned}$$

give identifications

$$\begin{aligned} O(U, U \oplus V) &= O(U \oplus V)/O(V) , \\ O(V, U \oplus V) &= O(U \oplus V)/O(U) , \\ G(V, U \oplus V) &= O(U \oplus V)/(O(U) \times O(V)) = G(U, U \oplus V) . \end{aligned}$$

(iii) For any finite-dimensional inner product spaces U, V, W there is defined a commutative braid of fibre bundles



where $O(U, V, W) = O(U \oplus V \oplus W)/\langle O(U \oplus V), O(U \oplus W) \rangle$.

□

Definition 3.22. The canonical U - and V -bundles over the Grassmann manifold $G = G(U, U \oplus V) = G(V, U \oplus V)$ are

$$\begin{aligned} \xi(U) : U &\rightarrow E(\xi(U)) = O(U, U \oplus V) \times_{O(U)} U \rightarrow O(U, U \oplus V)/O(U) = G , \\ \xi(V) : V &\rightarrow E(\xi(V)) = O(V, U \oplus V) \times_{O(V)} V \rightarrow O(V, U \oplus V)/O(V) = G \end{aligned}$$

with Thom spaces

$$\begin{aligned} T(\xi(U)) &= O(U, U \oplus V)^+ \wedge_{O(U)} U^\infty , \\ T(\xi(V)) &= O(V, U \oplus V)^+ \wedge_{O(V)} V^\infty . \end{aligned}$$

An element $(W, x) \in E(\xi(U))$ (resp. $(W, y) \in E(\xi(V))$) can be regarded as a subspace $W \subseteq U \oplus V$ isomorphic to U , together with an element $x \in W$

(resp. $y \in W^\perp$).

□

Let $p_U : U \oplus V \rightarrow U$, $p_V : U \oplus V \rightarrow V$ be the projections. For any $W \in G(U, U \oplus V)$ there exists $h \in O(U \oplus V)$ such that

$$h(U) = W, \quad h(V) = W^\perp \subseteq U \oplus V,$$

The maps

$$\begin{aligned} G \times (U \oplus V) &\rightarrow E(\xi(U)) = O(U \oplus V)/O(V) \times_{O(U)} U; \\ &\quad (W, u, v) \mapsto (h, p_U h^{-1}(u, v)) \\ G \times (U \oplus V) &\rightarrow E(\xi(V)) = O(U \oplus V)/O(U) \times_{O(V)} V; \\ &\quad (W, u, v) \mapsto (h, p_V h^{-1}(u, v)) \end{aligned}$$

fit into a pullback square

$$\begin{array}{ccc} G \times (U \oplus V) & \longrightarrow & E(\xi(U)) \\ \downarrow & & \downarrow \\ E(\xi(V)) & \longrightarrow & G \end{array}$$

defining the canonical isomorphism

$$\xi(U) \oplus \xi(V) \cong \epsilon_{U \oplus V}.$$

Definition 3.23. Let V be a finite-dimensional inner product space.

(i) The *classifying space* for V -bundles is

$$BO(V) = \varinjlim_U G(V, U \oplus V) = \varinjlim_U O(U \oplus V)/(O(U) \times O(V))$$

with U running over finite-dimensional inner product spaces.

(ii) The *universal V -bundle* is

$$\xi(V) = \varinjlim_U \xi(V) : V \rightarrow EO(V) = \varinjlim_U O(V, U \oplus V) \times_{O(V)} V \rightarrow BO(V).$$

(iii) The *universal Thom spectrum* is

$$\underline{MO} = \{MO(V) \mid \dim(V) < \infty\}$$

with $EO(V)$ a contractible space with free $O(V)$ -action

$$MO(V) = T(\xi(V)) = \varinjlim_U O(V, U \oplus V)^+ \wedge_{O(V)} V^\infty.$$

□

Terminology 3.24 The Stiefel manifold

$$V_{j+k,k} = O(\mathbb{R}^k, \mathbb{R}^{j+k}) = O(\mathbb{R}^{j+k})/O(\mathbb{R}^j)$$

of linear isometries $\mathbb{R}^k \hookrightarrow \mathbb{R}^{j+k}$ is (homeomorphic to) the space of orthogonal k -frames in \mathbb{R}^{j+k} . The Grassmann manifold

$$G(\mathbb{R}^k, \mathbb{R}^{j+k}) = V_{j+k,k}/O(\mathbb{R}^k) = O(\mathbb{R}^{j+k})/(O(\mathbb{R}^j) \times O(\mathbb{R}^k))$$

is (homeomorphic to) the space of k -dimensional subspaces

$$W = \text{im}(\mathbb{R}^k \hookrightarrow \mathbb{R}^{j+k}) \subseteq \mathbb{R}^{j+k}.$$

The canonical \mathbb{R}^j -bundle $\xi(\mathbb{R}^j)$ and the canonical \mathbb{R}^k -bundle $\xi(\mathbb{R}^k)$ over the Grassmann manifold

$$\begin{aligned} \mathbb{R}^j &\rightarrow E(\xi(\mathbb{R}^j)) \rightarrow G(\mathbb{R}^k, \mathbb{R}^{j+k}), \\ \mathbb{R}^k &\rightarrow E(\xi(\mathbb{R}^k)) \rightarrow G(\mathbb{R}^k, \mathbb{R}^{j+k}) \end{aligned}$$

are such that

$$\begin{aligned} E(\xi(\mathbb{R}^k)) &= V_{j+k,k} \times_{O(\mathbb{R}^k)} \mathbb{R}^k = \{(W \subseteq \mathbb{R}^{j+k}, x \in W) \mid \dim(W) = k\}, \\ E(\xi(\mathbb{R}^j)) &= \{(W \subseteq \mathbb{R}^{j+k}, y \in W^\perp) \mid \dim(W) = k\}, \\ \xi(\mathbb{R}^j) \oplus \xi(\mathbb{R}^k) &= \epsilon_{\mathbb{R}^{j+k}}. \end{aligned}$$

Passing to the limit as $j \rightarrow \infty$ gives the universal \mathbb{R}^k -bundle

$$\begin{aligned} \mathbb{R}^k \rightarrow EO(\mathbb{R}^k) &= \varinjlim_j V_{j+k,k} \times_{O(\mathbb{R}^k)} \mathbb{R}^k \\ &\rightarrow BO(\mathbb{R}^k) = \varinjlim_j G(\mathbb{R}^k, \mathbb{R}^{j+k}). \end{aligned}$$

with Thom space $MO(\mathbb{R}^k)$ and Thom spectrum $\underline{MO} = \{MO(\mathbb{R}^k)\}$.

□

Remark 3.25. We recall some more standard facts about the Stiefel spaces.

(i) $V_{j+k,k} = O(\mathbb{R}^{j+k})/O(\mathbb{R}^j)$ is $(j-1)$ -connected, with

$$\pi_i(V_{j+k,k}) = \begin{cases} 0 & \text{if } i < j \\ \mathbb{Z} & \text{if } i = j \text{ is even, or if } i = j, k = 1 \\ \mathbb{Z}_2 & \text{if } i = j \text{ is odd and } k \geq 2 \end{cases}$$

(ii) $V_{j+k,k}$ fits into a fibre bundle

$$V_{j+k,k} \rightarrow BO(\mathbb{R}^j) \rightarrow BO(\mathbb{R}^{j+k}) .$$

The homotopy classes of maps $c : X \rightarrow V_{j+k,k}$ are in one-one correspondence with the equivalence classes of pairs $(\xi, \delta\xi)$ with $\xi : X \rightarrow BO(\mathbb{R}^j)$ an \mathbb{R}^j -bundle over X and $\delta\xi : \xi \oplus \epsilon_{\mathbb{R}^k} \cong \epsilon_{\mathbb{R}^{j+k}}$ a k -stable isomorphism, with

$$E(\xi) = \{(x \in X, y \in c(x)^\perp \subset \mathbb{R}^{j+k})\} \subset E(\epsilon_{\mathbb{R}^{j+k}}) = X \times \mathbb{R}^{j+k} .$$

(iii) $V_{j+1,1} = S^j$, with $1 : S^j \rightarrow V_{j+1,1}$ classifying the tangent \mathbb{R}^j -bundle $\tau_{S^j} : S^j \rightarrow BO(\mathbb{R}^j)$ with the stable isomorphism $\tau_{S^j} \oplus \epsilon_{\mathbb{R}} \cong \epsilon_{\mathbb{R}^{j+1}}$ determined by the standard framed embedding $S^j \subset \mathbb{R}^{j+1}$.

(iv) Let $j \geq 0, k \geq 1$. For any \mathbb{R}^{j+k} -bundle $\xi : X \rightarrow BO(\mathbb{R}^{j+k})$ over a *CW* complex X it is possible to split $\xi|_{X^{(j)}} \cong \xi' \oplus \epsilon_{\mathbb{R}^k}$ for some \mathbb{R}^j -bundle ξ' over $X^{(j)}$. The $(j+1)$ -th *Stiefel-Whitney class* of ξ

$$w_{j+1}(\xi) \in H^{j+1}(X; \{\pi_j(V_{j+k,k})\})$$

is the primary obstruction to splitting $\xi \cong \xi'' \oplus \epsilon_{\mathbb{R}^k}$ for some \mathbb{R}^j -bundle ξ'' over X .

(v) The *Euler class* of an \mathbb{R}^{j+1} -bundle $\xi : X \rightarrow BO(\mathbb{R}^{j+1})$ is the $(j+1)$ th Stiefel-Whitney class

$$\gamma(\xi) = w_{j+1}(\xi) \in H^{j+1}(X; \{\mathbb{Z}\}) ,$$

the primary obstruction to splitting $\xi = \xi' \oplus \epsilon_{\mathbb{R}}$.

(vi) The mod 2 Stiefel-Whitney class $w_{j+1}(\xi) \in H^{j+1}(X; \mathbb{Z}_2)$ of $\xi : X \rightarrow BO(\mathbb{R}^{j+k})$ is given by the evaluation of the Steenrod square

$$Sq^{j+1} : \dot{H}^{j+k}(T(\xi); \mathbb{Z}_2) \rightarrow \dot{H}^{2j+k+1}(T(\xi); \mathbb{Z}_2)$$

on the mod 2 Thom class $U_\xi \in \dot{H}^{j+k}(T(\xi); \mathbb{Z}_2)$, with

$$Sq^{j+1}(U_\xi) = U_\xi \cup w_{j+1}(\xi) \in \dot{H}^{2j+k+1}(T(\xi); \mathbb{Z}_2) .$$

□

The projection

$$p : O(V, U \oplus V) \rightarrow G(U, U \oplus V) = O(V, U \oplus V)/O(V)$$

is such that there is defined a pullback square

$$\begin{array}{ccc} O(V, U \oplus V) \times V & \longrightarrow & E(\xi(V)) = O(V, U \oplus V) \times_{O(V)} V \\ \downarrow p^* \xi(V) & & \downarrow \xi(V) \\ O(V, U \oplus V) & \xrightarrow{p} & O(V, U \oplus V)/O(V) \end{array}$$

defining an isomorphism of V -bundles over $O(V, U \oplus V)$

$$p^*\xi(V) \cong \epsilon_V .$$

The pullback

$$p^*\xi(U) : U \rightarrow O(U \oplus V) \times_{O(U)} U \rightarrow O(U \oplus V)/O(U) = O(V, U \oplus V)$$

is such that

$$\begin{aligned} O(U \oplus V) \times_{O(U)} (U \oplus V) &\rightarrow O(V, U \oplus V) \times (U \oplus V) ; \\ (h, u, v) &\mapsto (h|_V, p_U h^{-1}(u), p_V h^{-1}(u) + v) \end{aligned}$$

defines an isomorphism of $U \oplus V$ -bundles over $O(V, U \oplus V)$

$$p^*\xi(U) \oplus \epsilon_V \cong \epsilon_{U \oplus V} ,$$

corresponding to the fibration sequence

$$O(U) \longrightarrow O(U \oplus V) \longrightarrow O(V, U \oplus V) \xrightarrow{p} BO(U) \longrightarrow BO(U \oplus V) .$$

Proposition 3.26. *Let X be a reasonable space, such as a finite CW complex.*

(i) *The isomorphism classes of V -bundles over X*

$$\xi : V \longrightarrow E(\xi) \longrightarrow X$$

are in bijective correspondence with the set $[X, BO(V)]$ of homotopy classes of maps

$$f : X \rightarrow G(V, U \oplus V) \subset BO(V) \quad (\dim(U) \text{ large})$$

with

$$\begin{aligned} \xi &= f^*\xi(V) , \\ E(\xi) &= f^*E(\xi(V)) = \{(u, v, x) \in U \oplus V \times X \mid (u, v) \in f(x) \subseteq U \oplus V\} . \end{aligned}$$

The V -bundles over X

$$\xi : |V| \longrightarrow E(\xi) \longrightarrow X$$

are classified by the homotopy classes of maps

$$f : X \rightarrow G(V, V \oplus \mathbb{R}^p) \subset BO(V) \quad (p \text{ large}) ,$$

with

$$\xi = f^*\xi(V) , \quad E(\xi) = f^*E(\xi(V)) = \{(x, y) \in X \times V \oplus \mathbb{R}^p \mid y \in f(x)\} .$$

The function

$$\begin{aligned} [X, BO(V)] &\rightarrow \{\text{isomorphism classes of } V\text{-bundles over } X\} ; \\ (f : X \rightarrow BO(V)) &\mapsto f^*\xi \end{aligned}$$

is a bijection.

(ii) The isomorphism classes of pairs $(\xi, \delta\xi)$ with ξ a V -bundle over X and

$$\delta\xi : \xi \oplus \epsilon_U \cong \epsilon_{U \oplus V}$$

a $U \oplus V$ -bundle isomorphism are in bijective correspondence with the homotopy classes of maps $g : X \rightarrow O(U, U \oplus V)$, with

$$\begin{aligned} \xi &= (pg)^*\xi(U) : X \xrightarrow{g} O(U, U \oplus V) \xrightarrow{p} G(V, U \oplus V) \subset BO(V) , \\ E(\xi) &= (pg)^*E(\xi(V)) = \{(u, v, x) \in U \oplus V \times X \mid (u, v) \in pg(x) \subseteq U \oplus V\} , \\ \delta\xi : E(\xi \oplus \epsilon_U) &= U \times E(\xi) \rightarrow E(\epsilon_{U \oplus V}) = U \oplus V \times X ; \\ &(t, u, v, x) \mapsto (t + u, v, x) \quad ((u, v) \in g(x)(U)^\perp \subseteq U \oplus V) . \end{aligned}$$

□

Example 3.27. Let M^m an m -dimensional manifold with an embedding $M \subset \mathbb{R}^m \oplus V$, for some inner product space V . The tangent \mathbb{R}^m -bundle τ_M and the normal V -bundle $\nu_M = \nu_{M \subset \mathbb{R}^m \oplus V}$ are classified by

$$\begin{aligned} \tau_M &: M \rightarrow G(\mathbb{R}^m, \mathbb{R}^m \oplus V) \subset BO(\mathbb{R}^m) , \\ \nu_M &= \tau_M^\perp : M \rightarrow G(V, \mathbb{R}^m \oplus V) \subset BO(V) \end{aligned}$$

with

$$\tau_M \oplus \nu_M = \tau_{\mathbb{R}^m \oplus V}|_M = \epsilon_{\mathbb{R}^m \oplus V} : M \rightarrow G(\mathbb{R}^m \oplus V, \mathbb{R}^m \oplus V) \subset BO(\mathbb{R}^m \oplus V) .$$

□

Example 3.28. Let $V = U \oplus \mathbb{R}$, for some finite-dimensional inner product space U . The homeomorphism

$$S(V) \rightarrow O(\mathbb{R}, V) ; \quad x \mapsto (t \mapsto tx)$$

is the clutching function of the tangent U -bundle $\tau_{S(V)} : S(V) \rightarrow BO(U)$

$$U \rightarrow E(\tau_{S(V)}) \rightarrow S(V)$$

with sphere bundle

$$S(\tau_{S(V)}) = \{(x, y) \in V \oplus V \mid \|x\| = 1, \langle x, y \rangle = 0\}.$$

The open embedding

$$S(V) \times \mathbb{R} \hookrightarrow V ; (x, t) \mapsto e^t x$$

corresponds to the V -bundle isomorphism

$$\delta\tau_{S(V)} : \tau_{S(V)} \oplus \nu_{S(V) \subset V} = \tau_{S(V)} \oplus \epsilon_{\mathbb{R}} \cong \tau_V| = \epsilon_V$$

classified by $S(V) \rightarrow O(\mathbb{R}, V) = O(V)/O(U)$.

□

Definition 3.29. Let $\xi : X \rightarrow BO(V)$, $\xi' : X \rightarrow BO(V')$ be vector bundles over a space X , for some finite-dimensional inner product spaces V, V' .

(i) The *dimension* of the virtual bundle $\xi - \xi'$ is

$$\dim(\xi - \xi') = \dim(V) - \dim(V') \in \mathbb{Z}.$$

(ii) The *virtual Thom space* of $\xi - \xi'$ is the spectrum

$$\underline{T}(\xi - \xi') = \{T(\xi - \xi')_U \mid U\}$$

defined by

$$T(\xi - \xi')_U = T(\xi \oplus \eta)$$

for finite-dimensional inner product spaces U with an isometry $V' \rightarrow U$ and a vector bundle η over X such that $\xi' \oplus \eta = \epsilon_U$. Note that

$$\dim(\xi - \xi') = \dim(\xi \oplus \eta) - \dim(U) \in \mathbb{Z}.$$

(iii) Write

$$\omega_*(X; \xi - \xi') = \tilde{\omega}_*(\underline{T}(\xi - \xi')) , \quad \omega^*(X; \xi - \xi') = \tilde{\omega}^*(\underline{T}(\xi - \xi'))$$

for the reduced stable homotopy and cohomotopy of the virtual Thom space of $\xi - \xi'$, so that

$$\begin{aligned} \omega_*(X; \xi - \xi') &= \tilde{\omega}_{*+\dim(U)}(T(\xi \oplus \eta)) , \\ \omega^*(X; \xi - \xi') &= \tilde{\omega}^{*+\dim(U)}(T(\xi \oplus \eta)) . \end{aligned}$$

□

Example 3.30. (i) In the special case $\xi' = \epsilon_{\mathbb{R}^\ell}$ ($\ell \geq 0$) the virtual Thom space $\underline{T}(\xi - \xi')$ is just the ℓ -fold desuspension of the suspension spectrum $\underline{T}(\xi)$ of the actual Thom space $T(\xi)$

$$\begin{aligned}\underline{T}(\xi - \xi') &= \Sigma^{-\ell} \{T(\xi \oplus \epsilon_U) | U\} , \\ &= \{T(\xi \oplus \epsilon_{U/\mathbb{R}^\ell}) | U, \mathbb{R}^\ell \subseteq U\} ,\end{aligned}$$

and

$$\begin{aligned}\omega_*(X; \xi - \epsilon_{\mathbb{R}^\ell}) &= \tilde{\omega}_{*+\ell}(T(\xi)) , \\ \omega^*(X; \xi - \epsilon_{\mathbb{R}^\ell}) &= \tilde{\omega}^{*+\ell}(T(\xi)) .\end{aligned}$$

(iii) For a virtual trivial bundle $\xi - \xi' = \epsilon_{\mathbb{R}^k} - \epsilon_{\mathbb{R}^\ell}$ over a compact space X

$$\begin{aligned}\omega_*(X; \epsilon_{\mathbb{R}^k} - \epsilon_{\mathbb{R}^\ell}) &= \omega_{*-k+\ell}(X) , \\ \omega^*(X; \epsilon_{\mathbb{R}^k} - \epsilon_{\mathbb{R}^\ell}) &= \omega^{*-k+\ell}(X) .\end{aligned}$$

□

Proposition 3.31. (The tubular neighbourhood theorem).

(i) Every embedding of manifolds $M^m \hookrightarrow N^n$ has a normal \mathbb{R}^{n-m} -bundle

$$\nu_{M \subset N} : M \rightarrow BO(n-m)$$

such that

$$\tau_M \oplus \nu_{M \subset N} = \tau_N|_M : M \rightarrow BO(\mathbb{R}^n) ,$$

with a codimension 0 embedding $E(\nu_{M \subset N}) \subset N$ and hence an Umkehr map

$$F : N^\infty \rightarrow N/(N \setminus E(\nu_{M \subset N})) = T(\nu_{M \subset N}) .$$

(ii) Every immersion of manifolds $f : M^m \looparrowright N^n$ has a normal \mathbb{R}^{n-m} -normal W -bundle

$$\nu_{M \looparrowright N} : \mathbb{R}^n \rightarrow E(\nu_{M \looparrowright N}) \rightarrow M ,$$

with a codimension 0 immersion $E(\nu_{M \looparrowright N}) \looparrowright N$. For any inner product space V with $\dim(V) = j$ the immersion

$$(0, f) : M \looparrowright V \times N ; x \mapsto (0, f(x))$$

has codimension $j + n - m$, and normal bundle $\nu_{(0, f)} = \nu_{M \looparrowright N} \oplus \epsilon_V$. For $j \geq 2m - n + 1$ $(0, f)$ is regular homotopic to an embedding of the form

$$(e, f) : M \hookrightarrow V \times N ; x \mapsto (e(x), f(x))$$

for some $e : M \rightarrow V$, with $\nu_{(e, f)} = \nu_{M \looparrowright N} \oplus \epsilon_V$. There is an extension of (e, f) to an open embedding

$$E(\nu_{M \looparrowright N} \oplus \epsilon_V) \hookrightarrow E(\epsilon_V) = V \times N$$

with adjunction Umkehr map

$$F : T(\epsilon_V) = V^\infty \wedge N^+ \rightarrow T(\nu_{M \looparrowright N} \oplus \epsilon_V) = V^\infty \wedge T(\nu_{M \looparrowright N}) .$$

□

Example 3.32. For any m -dimensional manifold M there exists an embedding $M \subset \mathbb{R}^m \oplus V$, for some finite-dimensional inner product space V . By the tubular neighbourhood theorem the embedding has a normal V -bundle ν_M with an embedding $E(\nu_M) \subset \mathbb{R}^m \oplus V$. The composite of the Umkehr map

$$\begin{aligned} \alpha_{\mathbb{R}^m \oplus V} : (\mathbb{R}^m \oplus V)^\infty \\ \rightarrow (\mathbb{R}^m \oplus V)/(\mathbb{R}^m \oplus V \setminus E(\nu_M)) = D(\nu_M)/S(\nu_M) = T(\nu_M) \end{aligned}$$

and the diagonal map $\Delta : T(\nu_M) \rightarrow M^+ \wedge T(\nu_M)$

$$\Delta \alpha_{\mathbb{R}^m \oplus V} : (\mathbb{R}^m \oplus V)^\infty \rightarrow M^+ \wedge T(\nu_M)$$

represents an element

$$(M, 1) = \Delta \alpha_{\mathbb{R}^m \oplus V} \in \omega_m(M, \nu_M - \epsilon_V) = \tilde{\omega}_{m+\dim(V)}(T(\nu_M)) .$$

□

3.3 Bordism

Every m -dimensional manifold M admits an embedding $M \subset \mathbb{R}^{m+n}$ (n large). The tangent \mathbb{R}^m -bundle $\tau_M : M \rightarrow BO(\mathbb{R}^m)$ and the normal \mathbb{R}^n -bundle $\nu_{M \subset \mathbb{R}^{m+n}} : M \rightarrow BO(\mathbb{R}^n)$ are such that

$$\tau_M \oplus \nu_M = \tau_{\mathbb{R}^{m+n}}|_M = \epsilon_{\mathbb{R}^{m+n}} : M \rightarrow BO(\mathbb{R}^{m+n}) .$$

The Pontryagin-Thom Umkehr map $\alpha_M : S^{m+n} \rightarrow T(\nu_M)$ is transverse regular at the zero section $M \subset T(\nu_M)$ with

$$(\alpha_M)^{-1}(M) = M \subset \mathbb{R}^{m+n} \subset (\mathbb{R}^{m+n})^\infty = S^{m+n} .$$

Definition 3.33. Let $\xi - \xi'$ be a virtual bundle over a space X , represented by bundles ξ, ξ' with $\dim(\xi) = i$, $\dim(\xi') = i'$.

(i) A m -dimensional normal map

$$(f, b) : (M, \nu_M - \epsilon_{\mathbb{R}^n}) \rightarrow (X, \xi - \xi')$$

consists of an m -dimensional manifold M with an embedding $M \subset \mathbb{R}^{m+n+i-i'}$ (n large), a map $f : M \rightarrow X$ and a bundle map over f

$$b : \nu_M \oplus f^* \xi' \rightarrow \xi \oplus \epsilon_{\mathbb{R}^n}$$

with

$$\nu_M = \nu_{M \subset \mathbb{R}^{m+n+i-i'}} : M \rightarrow BO(n+i-i') .$$

Equivalently, b can be regarded as a bundle map

$$b : \tau_M \oplus f^* \xi \oplus \epsilon_{\mathbb{R}^n} \rightarrow \xi' \oplus \epsilon_{\mathbb{R}^{m+n+i-i'}}$$

and as a virtual bundle map $b : \nu_M - \epsilon_{\mathbb{R}^n} \rightarrow \xi - \xi'$. Let $\eta : M \rightarrow BO(j)$ be a bundle such that $\xi' \oplus \eta = \epsilon_{\mathbb{R}^{i'+j}}$, so that $M \subset \mathbb{R}^{m+n+i+j}$ has normal bundle

$$\begin{aligned} \nu_{M \subset \mathbb{R}^{m+n+i+j}} &= \nu_M \oplus \epsilon_{\mathbb{R}^{i'+j}} \\ &= \nu_M \oplus f^* \xi' \oplus f^* \eta \\ &= f^*(\xi \oplus \eta \oplus \epsilon_{\mathbb{R}^n}) : M \rightarrow BO(n+i+j) \end{aligned}$$

allowing the *Pontryagin-Thom map* to be defined by

$$\begin{aligned} \alpha(f, b) : S^{m+n+i+j} &\xrightarrow{\alpha_M} \Sigma^{i'+j} T(\nu_M) = T(\nu_{M \subset \mathbb{R}^{m+n+i+j}}) \\ &\xrightarrow{T(b)} T(\xi \oplus \eta \oplus \epsilon_{\mathbb{R}^n}) . \end{aligned}$$

(ii) Let $\Omega_m(X, \xi - \xi')$ be the bordism group of m -dimensional normal maps $(f, b) : (M, \nu_M - \epsilon_{\mathbb{R}^n}) \rightarrow (X, \xi - \xi')$, with *Pontryagin-Thom map*

$$\begin{aligned} PT : \Omega_m(X, \xi - \xi') &\rightarrow \omega_{m+i-i'}(X; \xi - \xi') = \{S^{m+n+i+j}; T(\xi \oplus \eta \oplus \epsilon_{\mathbb{R}^n})\} ; \\ (f, b) &\mapsto \alpha(f, b) . \end{aligned}$$

(iii) Let $\mathfrak{N}_m(X)$ be the bordism group of m -dimensional manifolds M with a map $f : M \rightarrow X$.

□

Proposition 3.34. (i) *The Pontryagin-Thom maps*

$$PT : \Omega_m(X, \xi - \xi') \rightarrow \omega_{m+i-i'}(X; \xi - \xi') \quad (m \geq 0)$$

are isomorphisms, with inverses given by the transversality construction. Every element

$$F \in \omega_{m+i-i'}(X; \xi - \xi') = \{S^{m+n+i+j}; T(\xi \oplus \eta \oplus \epsilon_{\mathbb{R}^n})\}$$

is represented by a map $F : S^{m+n+i+j} \rightarrow T(\xi \oplus \eta \oplus \epsilon_{\mathbb{R}^n})$ which is transverse regular at the zero section $X \subset T(\xi \oplus \eta \oplus \epsilon_{\mathbb{R}^n})$ with the restriction

$$f = F| : M^m = F^{-1}(X) \rightarrow X$$

such that

$$\begin{aligned}\nu_{M \subset \mathbb{R}^{m+n+i+j}} \oplus f^* \xi' &= f^*(\xi \oplus \xi' \oplus \eta \oplus \epsilon_{\mathbb{R}^n}) \\ &= f^* \xi \oplus \epsilon_{\mathbb{R}^{n+i'+j}} : M \rightarrow BO(\mathbb{R}^{n+i+j}) ,\end{aligned}$$

and

$$T(b)\alpha_M = F \in \omega_{m+i-i'}(X; \xi - \xi') .$$

(ii) The maps

$$\begin{aligned}\mathfrak{N}_m(X) &\rightarrow \varinjlim_V \Omega_m(X \times BO(V), 0 \times \xi(V) - \epsilon_V) ; \\ (f : M \rightarrow X) &\mapsto (f \times \nu_M : (M, \nu_M - \epsilon_V) \rightarrow (X \times BO(V), \xi(V) - \epsilon_V))\end{aligned}$$

are isomorphisms. The Pontryagin-Thom maps

$$\begin{aligned}PT : \mathfrak{N}_m(X) &= \varinjlim_V \Omega_m(X \times BO(V), 0 \times \xi(V)) \rightarrow \\ &MO_m(X) = \varinjlim_V [\Sigma^m V^\infty; X^+ \wedge MO(V)]\end{aligned}$$

are isomorphisms. □

Definition 3.35. (i) An m -dimensional manifold M is *framed* if the normal bundle of an embedding $M \subset \mathbb{R}^{m+n}$ is trivialized,

$$\nu_M \cong \epsilon_{\mathbb{R}^n} ,$$

so that the Pontryagin-Thom Umkehr map

$$\alpha_M : (\mathbb{R}^{m+n})^\infty = S^{m+n} \rightarrow T(\nu_M) = \Sigma^n M^+$$

represents an element

$$\alpha_M \in \{S^{m+n}, \Sigma^n M^+\} = \omega_m(M) .$$

(ii) The m -dimensional framed bordism group of a space X

$$\Omega_m^{fr}(X) = \Omega_m(X, 0) \quad (\xi = \epsilon_V)$$

is the bordism group of m -dimensional framed manifolds M with a map $f : M \rightarrow X$, and the *framed Pontryagin-Thom map* is

$$PT^{fr} : \Omega_m^{fr}(X) \rightarrow \omega_m(X) ; ((f, b) : M \rightarrow X) \mapsto f_* \alpha_M .$$

□

Proposition 3.36. *The framed Pontryagin-Thom maps are isomorphisms*

$$PT^{fr} : \Omega_m^{fr}(X) \xrightarrow{\cong} \omega_m(X) .$$

Proof. This is just the special case $\xi - \xi' = 0$ of Proposition 3.34. \square

Example 3.37. In particular, the Pontryagin-Thom isomorphism identifies Ω_m^{fr} with $\omega_m = \{S^m; S^0\}$. \square

As usual, there are also relative bordism groups $\mathfrak{N}_m(X, Y)$ for a pair of spaces $(X, Y \subseteq X)$, with elements the bordism classes of maps $f : (M, \partial M) \rightarrow (X, Y)$ from m -dimensional manifolds with boundary. The relative bordism groups fits into an exact sequence

$$\cdots \rightarrow \mathfrak{N}_m(Y) \rightarrow \mathfrak{N}_m(X) \rightarrow \mathfrak{N}_m(X, Y) \rightarrow \mathfrak{N}_{m-1}(Y) \rightarrow \cdots$$

and the natural maps

$$\begin{aligned} \mathfrak{N}_*(X, Y) &= MO_m(X, Y) \rightarrow \\ \widetilde{\mathfrak{N}}_*(X/Y) &= \widetilde{MO}_m(X/Y) = \varinjlim_V [\Sigma^m V^\infty; X/Y \wedge MO(V)] \end{aligned}$$

are isomorphisms. For any \mathbb{R}^i -bundle $\xi : X \rightarrow BO(i)$ there are defined Thom isomorphisms

$$\begin{aligned} \mathfrak{N}_m(X) \rightarrow \mathfrak{N}_{m+i}(D(\xi), S(\xi)) &= \widetilde{\Omega}_{m+i}(T(\xi)) ; \\ (M, f : M \rightarrow X) &\mapsto (D(f^*\xi), S(f^*\xi)) . \end{aligned}$$

Example 3.38. (i) For pointed spaces X, Y a stable map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ induces morphisms

$$F_* : \widetilde{\mathfrak{N}}_m(X) = \widetilde{MO}_m(X) \rightarrow \widetilde{\mathfrak{N}}_m(Y) = \widetilde{MO}_m(Y)$$

(ii) For a map $F : X \rightarrow Y$ of unpointed spaces the corresponding pointed map $F^+ : X^+ \rightarrow Y^+$ is such that

$$F_*^+ : \mathfrak{N}_m(X) \rightarrow \mathfrak{N}_m(Y) ; (M, f : M \rightarrow X) \mapsto (M, Ff : M \rightarrow Y) .$$

(iii) The Umkehr map $F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f)$ of an immersion $f : M^m \looparrowright N^n$ is such that

$$F_* : \mathfrak{N}_n(N) \rightarrow \widetilde{\mathfrak{N}}_n(T(\nu_f)) = \mathfrak{N}_n(D(\nu_f), S(\nu_f)) = \mathfrak{N}_m(M) ;$$

$$(N, 1) \mapsto (N, \emptyset, z) = (M, 1)$$

with $z : N \rightarrow D(f)$ the zero section.

□

3.4 S -duality

Definition 3.39. Let X, Y be pointed spaces.

(i) Define the *cap* and *slant products* in stable (co)homotopy

$$\begin{aligned} \tilde{\omega}_N(X \wedge Y) \otimes \tilde{\omega}^i(X) &\rightarrow \tilde{\omega}_{N-i}(Y) ; \\ (\sigma : S^N \rightarrow X \wedge Y) \otimes (f : X \rightarrow S^i) &\mapsto ((1 \wedge f)\sigma : S^N \rightarrow \Sigma^i Y) , \\ \tilde{\omega}^N(X \wedge Y) \otimes \tilde{\omega}_i(X) &\rightarrow \tilde{\omega}^{N-i}(Y) ; \\ (\sigma^* : X \wedge Y \rightarrow S^N) \otimes (f : S^i \rightarrow X) &\mapsto (\sigma^*(f \wedge 1) : S^i \wedge Y \rightarrow S^N) \end{aligned}$$

for $i, N \in \mathbb{Z}$.

(ii) An element $\sigma \in \tilde{\omega}_N(X \wedge Y)$ is an S -duality if the products

$$\sigma \otimes - : \tilde{\omega}^i(X) \rightarrow \tilde{\omega}_{N-i}(Y) \quad (i \in \mathbb{Z})$$

are isomorphisms.

(iii) An element $\sigma^* \in \tilde{\omega}^N(X \wedge Y)$ is a *reverse S -duality* if the products

$$\sigma^* \otimes - : \tilde{\omega}_i(X) \rightarrow \tilde{\omega}^{N-i}(Y) \quad (i \in \mathbb{Z})$$

are isomorphisms.

□

Proposition 3.40. (i) If $\sigma \in \tilde{\omega}_N(X \wedge Y)$ is an S -duality there are induced isomorphisms

$$\begin{aligned} \sigma : \{X \wedge A; B\} &\xrightarrow{\cong} \{\Sigma^N A; B \wedge Y\} ; F \mapsto (F \wedge 1_Y)(\sigma \wedge 1_A) , \\ \sigma : \{A \wedge Y; B\} &\xrightarrow{\cong} \{\Sigma^N A; X \wedge B\} ; G \mapsto (1_X \wedge G)(1_A \wedge \sigma) \end{aligned}$$

for any pointed CW complexes A, B .

(ii) If $\sigma^* \in \tilde{\omega}^N(X \wedge Y)$ is a reverse S -duality there are induced isomorphisms

$$\begin{aligned}\sigma^* & : \{Z; X \wedge A\} \xrightarrow{\cong} \{Z \wedge Y; \Sigma^N A\} ; F \mapsto (\sigma^* \wedge 1_A)(F \wedge 1_Y) , \\ \sigma^* & : \{Z; A \wedge Y\} \xrightarrow{\cong} \{X \wedge Z; \Sigma^N A\} ; G \mapsto (\sigma^* \wedge 1_A)(1_X \wedge G)\end{aligned}$$

for any pointed CW complexes A, Z .

(iii) If X, Y are finite pointed CW complexes an element $\sigma \in \tilde{\omega}_N(X \wedge Y)$ is an S -duality if and only if the products

$$[\sigma] \otimes - : \tilde{H}^*(X) \rightarrow \tilde{H}_{N-*}(Y)$$

are isomorphisms, with $[\sigma] \in \tilde{H}_N(X \wedge Y)$ the Hurewicz image. Similarly for reverse S -duality.

(iv) If X, Y are finite pointed CW complexes there exists an S -duality $\sigma \in \tilde{\omega}_N(X \wedge Y)$ if and only if there exists a reverse S -duality $\sigma^* \in \tilde{\omega}^N(X \wedge Y)$, with $\sigma^* \sigma = 1 \in \{X; X\}$.

(v) For any finite pointed CW complex X there exists a finite pointed CW complex Y with an S -duality $\sigma \in \tilde{\omega}_N(X \wedge Y)$ and a reverse S -duality $\sigma^* \in \tilde{\omega}^N(X \wedge Y)$, with $\sigma^* \sigma = 1 \in \{X; X\}$.

(vi) Let M be an m -dimensional manifold with an embedding $M \subset \mathbb{R}^m \oplus V$ for $V = \mathbb{R}^i$, and let ν_M be the normal V -bundle. The composite of the Umkehr map of $E(\nu_M) \subset \mathbb{R}^m \oplus V$

$$\begin{aligned}\alpha_{\mathbb{R}^m \oplus V} & : (\mathbb{R}^m \oplus V)^\infty \\ & \rightarrow (\mathbb{R}^m \oplus V)/(\mathbb{R}^m \oplus V \setminus E(\nu_M)) = D(\nu_M)/S(\nu_M) = T(\nu_M)\end{aligned}$$

and the diagonal map $\Delta : T(\nu_M) \rightarrow M^+ \wedge T(\nu_M)$ represents an S -duality map

$$\Delta \alpha_{\mathbb{R}^m \oplus V} \in \omega_m(M \times M; 0 \times \nu_M - \epsilon_{\mathbb{R}^i}) = \tilde{\omega}_{m+i}(M^+ \wedge T(\nu_M)) .$$

Proof. Standard. □

Proposition 3.41. *Let M be an m -dimensional manifold with an embedding $M \subset \mathbb{R}^{m+n}$ with normal \mathbb{R}^n -bundle ν_M . For any bundles $\xi : M \rightarrow BO(\mathbb{R}^i)$, $\eta : M \rightarrow BO(\mathbb{R}^{n-i})$ such that*

$$\xi \oplus \eta = \nu_M : M \rightarrow BO(\mathbb{R}^n)$$

the composite

$$\sigma = \Delta \alpha_M : S^{m+n} \xrightarrow{\alpha_M} T(\nu_M) \xrightarrow{\Delta} T(\xi) \wedge T(\eta)$$

is an S -duality map such that products with the S -duality

$$\sigma \in \omega_m(M \times M; \xi \times \eta - \epsilon_{\mathbb{R}^n}) = \tilde{\omega}_{m+n}(T(\xi) \wedge T(\eta))$$

give Poincaré duality isomorphisms in stable homotopy

$$\sigma : \omega^*(M; \eta - \epsilon_{\mathbb{R}^{n-i}}) \cong \omega_{m-*}(M; \xi - \epsilon_{\mathbb{R}^i}) = \Omega_{m-*}(M, \xi) .$$

□

Example 3.42. Take $\xi = \nu_M$, $\eta = 0$ in Proposition 3.41, so that $i = n$ and the S -duality map is

$$\sigma = \Delta \alpha_M : S^{m+n} \xrightarrow{\alpha_M} T(\nu_M) \xrightarrow{\Delta} M^+ \wedge T(\nu_M)$$

Products with the S -duality $\sigma \in \tilde{\omega}_{m+n}(M^+ \wedge T(\nu_M))$ define Poincaré duality isomorphisms

$$\sigma : \omega^*(M) \cong \omega_{m-*}(M; \nu_M - \epsilon_{\mathbb{R}^n}) = \Omega_{m-*}(M, \nu_M) .$$

In particular, $1 \in \omega^0(M)$ is the Poincaré dual of

$$\sigma(1) = \alpha_M = (M, 1) \in \omega_m(M; \nu_M - \epsilon_{\mathbb{R}^n}) = \Omega_m(M, \nu_M) .$$

□

Example 3.43. (i) For any finite CW complex L there exists an embedding $L \subset S^N$ ($N \geq 2 \dim(L) + 1$). Let $W \subset S^N$ be a closed regular neighbourhood of $L \subset S^N$, so that the projection $p : W \rightarrow L$ is a homotopy equivalence with contractible point inverses. For any subcomplex $K \subseteq L$ the inverse image

$$V = p^{-1}(K) \subset W$$

is a closed regular neighbourhood of $K \subset S^N$, with $p| : V \rightarrow K$ a homotopy equivalence with contractible point inverses. The codimension 0 submanifolds

$$U = \overline{W \setminus V} , \quad V \subset W$$

have boundaries

$$\partial U = \partial_V U \cup \partial_W U , \quad \partial V = \partial_V U \cup \partial_W V$$

with

$$\begin{aligned} \partial_W U &= U \cap \partial W \subseteq U , \quad \partial_W V = V \cap \partial W \subseteq V , \\ \partial_V U &= U \cap V , \quad U/\partial_V U = W/V \simeq L/K . \end{aligned}$$

The composite

$$\sigma : S^N \longrightarrow S^N/(S^N \setminus U) = U/\partial U \xrightarrow{\Delta} U/\partial_V U \wedge U/\partial_W U \simeq L/K \wedge U/\partial_W U$$

represents an S -duality $\sigma \in \tilde{\omega}_N((L/K) \wedge (U/\partial_W U))$.

(ii) The special case $K = \{*\} \subset L$ in (i) gives an S -duality $\sigma \in \tilde{\omega}_N(L \wedge (U/\partial_W U))$ for any finite pointed CW complex L .

(iii) For a finite unpointed CW complex L the special case $K = \emptyset \subset L$ (with $V = \emptyset$, $(U, \partial_W U) = (W, \partial W)$) in (i) gives an S -duality map

$$\sigma : S^N \rightarrow L^+ \wedge W/\partial W .$$

If L is an n -dimensional geometric Poincaré complex (e.g. an n -dimensional manifold) then

$$S^{N-n-1} \rightarrow \partial W \rightarrow W \simeq L$$

is the Spivak normal fibration ν_L , and

$$\sigma \in \tilde{\omega}_N(L^+ \wedge W/\partial W) = \tilde{\omega}_N(L^+ \wedge T(\nu_L))$$

gives the Atiyah-Wall S -duality between L^+ and the Thom space $T(\nu_L) = W/\partial W$.

□

Proposition 3.44. *Let V be a finite-dimensional inner product space.*

(i) *The composite of the projection*

$$\alpha_V : V^\infty \rightarrow V^\infty/0^+ = \Sigma S(V)^+ ;$$

$$v = [t, u] = \frac{tu}{1-t} \mapsto \begin{cases} (t, u) = (\frac{\|v\|}{1+\|v\|}, \frac{v}{\|v\|}) & \text{if } v \neq 0, \infty \\ \infty & \text{if } v = 0 \text{ or } \infty \end{cases}$$

and the diagonal map

$$\Delta : \Sigma S(V)^+ \rightarrow S(V)^+ \wedge \Sigma S(V)^+ ; (t, u) \mapsto (u, (t, u))$$

is an S -duality map

$$\sigma_V = \Delta \alpha_V : V^\infty \xrightarrow{\alpha_V} \Sigma S(V)^+ \xrightarrow{\Delta} S(V)^+ \wedge \Sigma S(V)^+ ;$$

$$v = [t, u] \mapsto \begin{cases} (u, (t, u)) & \text{if } v \neq 0, \infty \\ \infty & \text{if } v = 0 \text{ or } \infty . \end{cases}$$

with S -duality isomorphisms

$$\begin{aligned}\sigma_V &: \{\Sigma S(V)^+ \wedge X; Y\} \xrightarrow{\cong} \{V^\infty \wedge X; S(V)^+ \wedge Y\} ; F \mapsto F\sigma_V , \\ \sigma_V &: \{S(V)^+ \wedge X; Y\} \xrightarrow{\cong} \{V^\infty \wedge X; \Sigma S(V)^+ \wedge Y\} ; G \mapsto G\sigma_V\end{aligned}$$

for any pointed spaces X, Y . The corresponding reverse S -duality map $\sigma_V^* : S(V)^+ \wedge \Sigma S(V)^+ \rightarrow V^\infty$ is given by the composite

$$\sigma_V^* : S(V)^+ \wedge \Sigma S(V)^+ \xrightarrow{\Sigma F} \Sigma T(\tau_{S(V)}) = V^\infty \wedge S(V)^+ \longrightarrow V^\infty$$

with $F : S(V)^+ \wedge S(V)^+ \rightarrow T(\tau_{S(V)})$ the Pontryagin-Thom map of the diagonal embedding $\Delta : S(V) \hookrightarrow S(V) \times S(V)$ with normal bundle $\nu_\Delta = \tau_{S(V)}$ such that $\nu_\Delta \oplus \epsilon_{\mathbb{R}} = \epsilon_V$.

(ii) The S -duality isomorphism

$$\{X; S(V)^+ \wedge Y\} \xrightarrow{\cong} \{\Sigma S(V)^+ \wedge X; V^\infty \wedge Y\} ; f \mapsto \Delta\alpha_V \wedge f$$

sends the stable relative difference $\delta'(p, q) \in \{X; S(V)^+ \wedge Y\}$ (3.6) of maps $p, q : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ which agree on $0^+ \wedge X \subset V^\infty \wedge X$ to the stable homotopy class of the relative difference $\delta(p, q) \in \{\Sigma S(V)^+ \wedge X; V^\infty \wedge Y\}$ (2.18), with

$$\begin{aligned}s_V \delta'(p, q) &= s_V \delta(p, q) = q - p \\ &\in \ker(0_V : \{X; Y\} \rightarrow \{X; V^\infty \wedge Y\}) = \text{im}(s_V : \{X; S(V)^+ \wedge Y\} \rightarrow \{X; Y\}) .\end{aligned}$$

(iii) The inclusion $S(V) \subset V$ is an embedding with trivial normal \mathbb{R} -bundle $\nu_{S(V) \subset V} = \epsilon_{S(V)}$. By (i) the composite of

$$\alpha_V : V^\infty \rightarrow T(\epsilon_{S(V)}) = \Sigma S(V)^+ ; [t, u] \mapsto (t, u)$$

and the diagonal map $\Delta : \Sigma S(V)^+ \rightarrow S(V)^+ \wedge \Sigma S(V)^+$ is an S -duality map

$$\sigma_V = \Delta\alpha_V : V^\infty \rightarrow S(V)^+ \wedge \Sigma S(V)^+ .$$

Thus for any pointed CW complexes A, B there is defined an S -duality isomorphism

$$\{\Sigma S(V)^+ \wedge A; V^\infty \wedge B\} \rightarrow \{A; S(V)^+ \wedge B\} ; F \mapsto (1 \wedge F)(\Delta\alpha_V \wedge 1) .$$

□

Example 3.45. Let M be an m -dimensional manifold, with tangent bundle $\tau_M : M \rightarrow BO(\mathbb{R}^m)$. Let $\nu_M : M \rightarrow BO(k)$ be the normal bundle of an embedding $M \subset \mathbb{R}^{m+k}$, so that

$$\tau_M \oplus \nu_M = \tau_{\mathbb{R}^{m+k}}|_M = \epsilon^{m+k} ,$$

and by the tubular neighbourhood theorem there is defined a codimension 0 embedding $E(\nu_M) \subset S^{m+k}$. By Proposition 3.40 (vi) the composite

$$\begin{aligned} \sigma_M : S^{m+k} &\rightarrow S^{m+k}/(S^{m+k} \setminus E(\nu_M)) = E(\nu_M)/S(\nu_M) = T(\nu_M) \\ &\xrightarrow{\Delta} E(\nu_M) \times E(\nu_M)/(E(\nu_M) \times S(\nu_M)) \simeq M^+ \wedge T(\nu_M) \end{aligned}$$

is an S -duality map. Let $z : M \hookrightarrow E(\nu_M)$ be the zero section. The embedding

$$(1 \times z)\Delta : M \hookrightarrow M \times E(\nu_M) ; x \mapsto (x, z(x))$$

has normal bundle

$$\nu_{(1 \times z)\Delta} = \tau_M \oplus \nu_M = \epsilon^{m+k} : M \rightarrow BO(\mathbb{R}^{m+k})$$

with an adjunction Umkehr map (2.2)

$$M^+ \wedge T(\nu_M) \rightarrow T(\nu_{(1 \times z)\Delta}) = \Sigma^{m+k} M^+ .$$

The composite

$$\sigma_M^* : M^+ \wedge T(\nu_M) \longrightarrow \Sigma^{m+k} M^+ \longrightarrow S^{m+k}$$

is a reverse S -duality map such that $\sigma_M^* \sigma_M = 1 \in \{M^+; M^+\}$, with S -duality isomorphisms

$$\begin{aligned} \sigma_M^* : \{X; T(\nu_M)\} &\xrightarrow{\cong} \{M^+ \wedge X; S^{m+k}\} ; F \mapsto \sigma_M^* F , \\ \sigma_M^* : \{X; M^+\} &\xrightarrow{\cong} \{X \wedge T(\nu_M); S^{m+k}\} ; G \mapsto \sigma_M^* G . \end{aligned}$$

□

3.5 The stable cohomotopy Thom and Euler classes

Proposition 3.46. *Let*

$$\xi : U \longrightarrow E(\xi) \xrightarrow{p_\xi} X , \quad \eta : V \longrightarrow E(\eta) \xrightarrow{p_\eta} X$$

be bundles over a space X .

(i) *There is defined a cofibration sequence*

$$T(\eta|_{S(\xi)}) \longrightarrow T(\eta) \xrightarrow{z} T(\xi \oplus \eta)$$

with

$$z : T(\eta) \rightarrow T(\xi \oplus \eta) ; x \mapsto (0, x) .$$

(ii) There is defined an isomorphism of exact sequences

$$\begin{array}{ccccccc} \dots \rightarrow \{T(\xi \oplus \eta); (U \oplus V)^\infty\} & \xrightarrow{z^*} & \{T(\eta); (U \oplus V)^\infty\} & \rightarrow & \{T(\eta|_{S(\xi)}); (U \oplus V)^\infty\} & \rightarrow & \dots \\ & \downarrow \cong & \downarrow \cong & & \downarrow \cong & & \\ \dots \rightarrow \omega^0(D(\xi), S(\xi); \eta) & \longrightarrow & \omega^0(X; \eta) & \xrightarrow{p_{S(\xi)}^*} & \omega^0(S(\xi); \eta) & \longrightarrow & \dots \end{array}$$

(iii) If $\xi \oplus \eta = \epsilon_{U \oplus V}$ and $\dim(U \oplus V) = m$

$$\begin{aligned} E(\xi \oplus \eta) &= (U \oplus V) \times X = \mathbb{R}^m \times X , \\ T(\xi \oplus \eta) &= T(\epsilon_{U \oplus V}) = (U \oplus V)^\infty \wedge X^+ = \Sigma^m X^+ , \\ \omega^n(D(\xi); -\xi) &= \tilde{\omega}^{m+n}(T(\eta)) = \{T(\eta); \Sigma^n(U \oplus V)^\infty\} = \tilde{\omega}^{m+n}(T(\eta)) , \\ \omega^n(D(\xi), S(\xi); -\xi) &= \omega^{m+n}(T(\eta), T(\eta|_{S(\xi)})) \\ &= \{T(\xi \oplus \eta); \Sigma^n(U \oplus V)^\infty\} = \{X^+; S^n\} = \omega^n(X) . \end{aligned}$$

□

Definition 3.47. (Crabb [12, §§2,5])

Let ξ be a U -bundle over a space X .

(i) The *stable cohomotopy Thom class* of ξ is

$$u(\xi) = 1 \in \omega^0(D(\xi), S(\xi); -\xi) = \omega^0(X)$$

with

$$1 : X^+ \rightarrow S^0 ; x \mapsto -1 , \infty \mapsto 1 .$$

(ii) The *stable cohomotopy Euler class* of ξ is

$$\gamma(\xi) = z^* u(\xi) \in \omega^0(X; -\xi) = \omega^0(D(\xi); -\xi) = \{T(\eta); (U \oplus V)^\infty\}$$

with η any V -bundle over X such that $\xi \oplus \eta = \epsilon_{U \oplus V}$, and is represented by

$$T(\eta) \xrightarrow{z} T(\xi \oplus \eta) = (U \oplus V)^\infty \wedge X^+ \rightarrow (U \oplus V)^\infty .$$

□

Example 3.48. For the trivial U -bundle $\xi = \epsilon_U$ over X can take $V = \{0\}$, $\eta = 0$, so that $\gamma(\epsilon_U)$ is represented by the constant map

$$T(\eta) = X^+ \rightarrow U^\infty ; x \mapsto 0$$

and

$$\gamma(\epsilon_U) = 0 \in \omega^0(X; -\epsilon_U) = \{X^+; U^\infty\} .$$

□

Example 3.49. Let M be an m -dimensional manifold, with tangent bundle $\tau_M : M \rightarrow BO(m)$, and let $M \subset \mathbb{R}^{m+k}$ be an embedding with normal bundle $\nu_M : M \rightarrow BO(k)$, so that

$$\tau_M \oplus \nu_M = \epsilon_{\mathbb{R}^{m+k}} : M \rightarrow BO(m+k) .$$

The stable cohomotopy Euler class of τ_M

$$\gamma(\tau_M) \in \omega^0(M; -\tau_M) = \{T(\nu_M); S^{m+k}\}$$

is represented by the composite

$$T(\nu_M) \xrightarrow{z} T(\tau_M \oplus \nu_M) = \Sigma^{m+k} M^+ \rightarrow S^{m+k} .$$

□

Proposition 3.50. (Crabb [12, §§2,5])

(i) *The stable cohomotopy Euler class of the Whitney sum $V \oplus W$ -bundle $\xi \oplus \eta$ of a V -bundle ξ over M and a W -bundle η over X is the product of the stable cohomotopy Euler classes of ξ, η*

$$\gamma(\xi \oplus \eta) = \gamma(\xi)\gamma(\eta) \in \omega^0(X; -(\xi \oplus \eta)) .$$

(ii) *For any V -bundle ξ over X there is a bijective correspondence between the splittings $\xi \cong \xi_1 \oplus \epsilon_{\mathbb{R}}$ and sections $s : X \rightarrow S(\xi)$ of $p_{S(\xi)} : S(\xi) \rightarrow X$. If there exists such a splitting (or section) then*

$$\gamma(\xi) = \gamma(\xi_1)\gamma(\epsilon_{\mathbb{R}}) = 0 \in \omega^0(X; -\xi)$$

(since $\gamma(\epsilon_{\mathbb{R}}) = 0$).

(iii) *If ξ is a V -bundle over X and $s : Y \rightarrow S(\xi|_Y)$ is a section of $p_{S(\xi|_Y)} : S(\xi|_Y) \rightarrow Y$ (for some $Y \subseteq X$) then using any extension $\tilde{s} : (X, Y) \rightarrow (D(\xi), S(\xi))$ of s there is defined an Euler class rel Y*

$$\gamma(\xi, s) = \tilde{s}^*(u) \in \omega^0(X, Y; -\xi) ,$$

with image

$$[\gamma(\xi, s)] = \gamma(\xi) \in \omega^0(X; -\xi) .$$

(iv) *If ξ is a V -bundle over X and $s_0, s_1 : Y \rightarrow S(\xi|_Y)$ are sections of $p_{S(\xi|_Y)} : S(\xi|_Y) \rightarrow Y$ which agree on $Z \subseteq Y$ there is defined a difference class*

$$\delta(s_0, s_1) \in \omega^{-1}(Y, Z; -\xi)$$

with image $\gamma(\xi, s_0) - \gamma(\xi, s_1) \in \omega^0(X, Y; -\xi)$.

(v) Let X be an n -dimensional manifold, and let $X \subset \mathbb{R}^n \oplus U$ be an embedding, with normal U -bundle ν_X . Given a V -bundle ξ over X let $g : X \rightarrow E(\xi)$ be a generic section transverse at the zero section $X \subset E(\xi)$, so that the restriction

$$(f, b) = g| : M^m = g^{-1}(X) \rightarrow X$$

is the normal map defined by the inclusion of an m -dimensional submanifold $M^m \subset X$ with

$$\begin{aligned} m &= n - \dim(V) , \quad \nu_{M \subset X} = \xi|_M , \\ b : \nu_M &= \nu_{M \subset X \subset \mathbb{R}^n \oplus U} = (\xi \oplus \nu_X)|_M \rightarrow \xi \oplus \nu_X . \end{aligned}$$

For any W -bundle η over X such that $\xi \oplus \eta = \epsilon_{V \oplus W}$ the restriction $\eta|_M$ is trivial, and the Euler class

$$\gamma(\xi) \in \omega^0(X; -\xi) = \{T(\eta); (V \oplus W)^\infty\}$$

is the stable homotopy class of the composite

$$T(\eta) \xrightarrow{F} T(\eta|_M) = M^+ \wedge (V \oplus W)^\infty \longrightarrow (V \oplus W)^\infty$$

with F the adjunction Umkehr map (2.2) of the inclusion

$$E(\eta|_M) = M \times (V \oplus W) \subset E(\eta) .$$

In terms of the S -dual formulation

$$\begin{aligned} \gamma(\xi) &= T(b)_*(M, 1) \\ &\in \omega^0(X; -\xi) = \omega_{m+\dim(U)+\dim(V)}(X; \xi \oplus \nu_X) = \Omega_m(X; \xi \oplus \nu_X) \end{aligned}$$

with $(M, 1) \in \omega_{m+\dim(U)+\dim(V)}(M; \nu_M)$.

(vi) Let $n, U, V, W, X, \nu_X, \xi, \eta$ be as in (v). The fibre product bundles

$$\begin{aligned} S(U) \times S(U) &\rightarrow S(\xi) \times_X S(\xi) \rightarrow X , \\ S(U \oplus U) &\rightarrow S(\xi \oplus \xi) = S(\xi) \times_X D(\xi) \cup D(\xi) \times_X S(\xi) \rightarrow X \end{aligned}$$

are such that there are defined bundle isomorphisms

$$\begin{aligned} \nu_{S(\xi) \times_X S(\xi) \hookrightarrow S(\xi \oplus \xi)} &\cong \epsilon_{\mathbb{R}} , \\ \nu_{\Delta: S(\xi) \hookrightarrow S(\xi) \times_X S(\xi)} \oplus \epsilon_{\mathbb{R}} &\cong \nu_{\Delta: S(\xi) \hookrightarrow S(\xi \oplus \xi)} \cong \epsilon_U . \end{aligned}$$

Given two sections $s_0, s_1 : X \rightarrow S(\xi)$ of ξ which agree on a submanifold $Y \subseteq X$ the section $(s_0, -s_1) : X \rightarrow S(\xi) \times_X S(\xi)$ is homotopic to a section

$(t_0, -t_1) : X \rightarrow S(\xi) \times_X S(\xi)$ transverse at the diagonal $\Delta_{S(\xi)} \subset S(\xi) \times_X S(\xi)$.
The inverse image

$$C = (t_0, -t_1)^{-1}(\Delta_{S(\xi)}) \subseteq X \setminus Y$$

is an $(m+1)$ -dimensional submanifold with

$$m = n - \dim(V) , \quad (\nu_{C \subset X}) \oplus \epsilon_{\mathbb{R}} \cong \xi|_C .$$

Inclusion defines a normal map $(f, b) : C \rightarrow X \setminus Y$ with

$$b : \nu_C = \nu_{C \subset X \subset \mathbb{R}^n \oplus U} \oplus \epsilon_{\mathbb{R}} = \nu_{C \subset X} \oplus \epsilon_{\mathbb{R}} \oplus \nu_X|_C \rightarrow (\xi \oplus \nu_X)|_{X \setminus Y} .$$

The rel Y difference class

$$\delta(s_0, s_1) \in \omega^{-1}(X, Y; -\xi) = \{\Sigma(T(\eta)/T(\eta|_Y)); (U \oplus V)^\infty\}$$

is the stable homotopy class of the composite

$$\begin{aligned} \Sigma(T(\eta)/T(\eta|_Y)) &= T(\eta \oplus \epsilon_{\mathbb{R}})/T(\eta \oplus \epsilon_{\mathbb{R}}|_Y) \xrightarrow{F} \\ T(\xi \oplus \eta|_C) &= C^+ \wedge (U \oplus V)^\infty \longrightarrow (U \oplus V)^\infty \end{aligned}$$

with F the adjunction Umkehr map of the inclusion

$$E((\eta \oplus \epsilon_{\mathbb{R}})|_C) = C \times (U \oplus V) \subset E(\eta \oplus \epsilon_{\mathbb{R}}) .$$

In terms of the S -dual formulation

$$\begin{aligned} \delta(s_0, s_1) &= T(b)_*(C, 1) \in \omega^{-1}(X, Y; -\xi) \\ &= \omega_{m+\dim(U)+\dim(V)+1}(X \setminus Y; (\xi \oplus \nu_X)|_{X \setminus Y}) = \Omega_{m+1}(X \setminus Y; (\xi \oplus \nu_X)|_{X \setminus Y}) \end{aligned}$$

with $(C, 1) \in \omega_{m+\dim(U)+\dim(V)+1}(C; \nu_C)$.

□

Example 3.51. (i) A non-zero section of the trivial V -bundle ϵ_V over a space X is essentially the same as a map of the type

$$X \rightarrow S(\epsilon_V) = X \times S(V) ; \quad x \mapsto (x, s(x)) ,$$

as determined by a map $s : X \rightarrow S(V)$. Any such map s determines a map

$$p_s : CX^+ \rightarrow V^\infty ; \quad (t, x) \mapsto \frac{ts(x)}{1-t}$$

sending $X = \{0\} \times X \subset CX^+$ to $0 \in V^\infty$. The rel $X \cup CY$ difference class of the non-zero sections (3.50 (iv)) given by two such maps $s_0, s_1 : X \rightarrow S(V)$

which agree on $Y \subseteq X$ is just the rel $X \cup CY$ difference (1.6 (iii)) of $p_{s_0}, p_{s_1} : CX^+ \rightarrow V^\infty$

$$\delta(s_0, s_1) = \delta(p_{s_0}, p_{s_1}) \in \omega^{-1}(X, Y; -\epsilon_V) = \{\Sigma(X/Y); V^\infty\}.$$

(ii) If X is an n -dimensional manifold and $s_0, s_1 : X \rightarrow S(V)$ are maps such that

$$d : X \rightarrow S(V) \times S(V) ; x \mapsto (s_0(x), -s_1(x))$$

is transverse regular at $\Delta_{S(V)} \subset S(V) \times S(V)$ then

$$C = d^{-1}(\Delta_{S(V)}) = \{x \in X \mid s_0(x) = -s_1(x) \in S(V)\} \subset X$$

is an $(m+1)$ -dimensional submanifold with

$$m = n - \dim(V), \quad \nu_{C \subset X} = (d|_C)^* \nu_{\Delta : S(V) \hookrightarrow S(V) \times S(V)} = (d|_C)^* \tau_{S(V)}$$

such that $\nu_{C \subset X} \oplus \epsilon_{\mathbb{R}} \cong \epsilon_V$, and

$$\delta(s_0, s_1) : \Sigma X^+ \xrightarrow{\Sigma F} \Sigma T(\nu_{C \subset X}) = C^+ \wedge V^\infty \longrightarrow V^\infty$$

with $F : X^+ \rightarrow T(\nu_{C \subset X})$ the adjunction Umkehr map of $E(\nu_{C \subset X}) \subset X$.

(iii) For any maps $s_0, s_1 : X = S(V) \rightarrow S(V)$ with

$$C = \{x \in X \mid s_0(x) = -s_1(x) \in S(V)\}$$

a 0-dimensional submanifold (= finite subset) of X , the difference class of the corresponding sections

$$s_i : X \rightarrow S(\epsilon_V) = X \times S(V) ; x \mapsto (x, s_i(x)) \quad (i = 0, 1)$$

of $p_{S(\epsilon_V)}$ is

$$\delta(s_0, s_1) = |C| \in \{\Sigma X^+; V^\infty\} = \mathbb{Z},$$

counting the points of C algebraically. In particular, if

$$s_0(x) = x, \quad s_1(x) = x_0 \in S(V) \quad (x \in S(V))$$

for some point $x_0 \in S(V)$ then $C = \{-x_0\} \subset X$ and

$$\delta(s_0, s_1) = |C| = 1 \in \{\Sigma X^+; V^\infty\} = \mathbb{Z}.$$

□

Definition 3.52. An element $c \in O(V, U \oplus V)$ is a linear isometry $c : V \rightarrow U \oplus V$. The *adjoint* of a map $c : X \rightarrow O(V, U \oplus V)$ is the pointed map

$$F_c : V^\infty \wedge X^+ \rightarrow (U \oplus V)^\infty ; (v, x) \mapsto c(x)(v).$$

□

Proposition 3.53. (i) *The stable cohomotopy Euler class of the canonical V -bundle $\xi(V) : G = G(V, U \oplus V) \rightarrow BO(V)$*

$$\gamma(\xi(V)) \in \omega^0(G; -\xi(V)) = \{T(\xi(V)); (U \oplus V)^\infty\}$$

is the stable homotopy class of the adjoint map

$$\gamma(\xi(V)) : T(\xi(U)) = O(U, U \oplus V) \wedge_{O(U)} U^\infty \rightarrow (U \oplus V)^\infty ; (i, x) \mapsto i(x) .$$

(ii) *Any V -bundle $\xi : X \rightarrow BO(V)$ over a finite CW complex X is isomorphic to the pullback $f^*\xi(V)$ of $\xi(V)$ along a map $f : X \rightarrow G = G(V, U \oplus V)$, for sufficiently large $\dim(U)$. The pullback $\eta = f^*\xi(U) : X \rightarrow BO(U)$ is a U -bundle over X such that $\xi \oplus \eta \cong \epsilon_{U \oplus V}$, and the stable cohomotopy Euler class*

$$\gamma(\xi) = f^*\gamma(\xi(V)) \in \omega^0(X; -\xi) = \{T(\eta); (U \oplus V)^\infty\}$$

is the stable homotopy class of the composite

$$\gamma(\xi) : T(\eta) \xrightarrow{f} T(\xi(U)) = O(U, U \oplus V)^+ \wedge_{O(U)} U^\infty \xrightarrow{\gamma(\xi(U))} (U \oplus V)^\infty .$$

A $U \oplus V$ -bundle isomorphism $\delta\xi : \xi \oplus \epsilon_U \cong \epsilon_{U \oplus V}$ corresponds to a lift of $\xi : X \rightarrow BO(V)$ to a map $\delta\xi : X \rightarrow O(U, U \oplus V)$, in which case $\eta \cong \epsilon_U$ and $\gamma(\xi) \in \omega^0(X; -\xi) = \{X^+; V^\infty\}$ is the stable homotopy class of the adjoint of $\delta\xi$

$$\gamma(\xi) : U^\infty \wedge X^+ \rightarrow (U \oplus V)^\infty ; (u, x) \mapsto \delta\xi(x)(u) .$$

(iii) *A section $s : X \rightarrow S(\xi)$ of $\xi : X \rightarrow BO(V)$ determines a null-homotopy*

$$\gamma(s) : \gamma(\xi) \simeq * : T(\eta) \rightarrow (U \oplus V)^\infty ,$$

as given by an extension of $\gamma(\xi)$ to a map $\gamma(s) : CT(\eta) \rightarrow (U \oplus V)^\infty$. The rel Y difference class (3.50 (iv)) of sections $s_0, s_1 : X \rightarrow S(\xi)$ which agree on $Y \subseteq X$ is just the rel $T(\eta) \cup CT(\eta|_Y)$ difference (1.6 (iii))

$$\delta(s_0, s_1) = \delta(\gamma(s_0), \gamma(s_1)) \in \omega^{-1}(X, Y; -\xi) = \{\Sigma(T(\eta)/T(\eta|_Y)); (U \oplus V)^\infty\} .$$

Proof. (i) The composite

$$\begin{aligned} T(\xi(U)) &= O(U, U \oplus V)^+ \wedge_{O(U)} U^\infty \xrightarrow{z=1 \wedge j} \\ T(\xi(U) \oplus \xi(V)) &= O(U, U \oplus V)^+ \wedge_{O(U)} (U \oplus V)^\infty \\ &\cong G^+ \wedge (U \oplus V)^\infty \longrightarrow (U \oplus V)^\infty \end{aligned}$$

is given by $(h, x) \mapsto h(x)$.

(ii) Since X is a finite CW complex there exists a sufficiently high-dimensional U such that the classifying map $\xi : X \rightarrow BO(V)$ factors up to homotopy as

$$\xi : X \xrightarrow{f} G = G(V, U \oplus V) \longrightarrow BO(V)$$

with $\xi \cong f^*\xi(V)$, $\xi \oplus f^*\xi(U) \cong \epsilon_{U \oplus V}$.

(iii) The stable map representing $\gamma(\xi)$ is the composite

$$\gamma(\xi) : T(\eta) \xrightarrow{z} T(\xi \oplus \eta) \cong (U \oplus V)^\infty \wedge X^+ \longrightarrow (U \oplus V)^\infty .$$

The Thom space of $\xi \oplus \eta$ fits into a cofibration sequence

$$T(p^*\eta) \xrightarrow{T(p)} T(\eta) \xrightarrow{z} T(\xi \oplus \eta) \longrightarrow \dots$$

with $p = p_{S(\xi)} : S(\xi) \rightarrow X$, and $T(p) : T(p^*\eta) \rightarrow T(\eta)$ the map of Thom spaces induced by the map of V -bundles $p : p^*\eta \rightarrow \eta$ over $p : S(\xi) \rightarrow X$. For any section $s : X \rightarrow S(\xi)$ of $p : S(\xi) \rightarrow X$ it follows from $p \circ s = 1 : X \rightarrow X$ that $s^*(p^*\eta) = \eta$, so that s lifts to a map of U -bundles $s : \eta \rightarrow p^*\eta$ inducing a section $T(s) : T(\eta) \rightarrow T(p^*\eta)$ of $T(p)$, giving the null-homotopies $z \simeq *$, $\gamma(\xi) \simeq *$.

□

Chapter 4

\mathbb{Z}_2 -equivariant homotopy and bordism theory

Intersections and self-intersections of maps of manifolds are expressed in terms of algebraic topology by means of the \mathbb{Z}_2 -equivariant homotopy properties of the diagonal maps

$$\begin{aligned}\Delta &: X \rightarrow X \times X ; x \mapsto (x, x) \text{ (} X \text{ unpointed) ,} \\ \Delta &: X \rightarrow X \wedge X ; x \mapsto (x, x) \text{ (} X \text{ pointed) .}\end{aligned}$$

4.1 π -equivariant homotopy theory

Let π be a group. We shall be concerned with pointed spaces X with a π -action

$$\pi \times X \rightarrow X ; (g, x) \mapsto gx ,$$

particularly for $\pi = \mathbb{Z}_2$.

Definition 4.1. (i) Given a (pointed) π -space X let $|X|$ be the (pointed) space defined by X with the π -action forgotten.

(ii) Given pointed π -spaces X, Y let $[X, Y]_\pi$ be the set of (pointed) π -equivariant homotopy classes of π -equivariant maps. For any pointed π -spaces X, Y there is defined a forgetful function

$$[X, Y]_\pi \rightarrow [|X|, |Y|] .$$

(iii) The *fixed point set* of a π -space X is

$$X^\pi = \{x \in X \mid gx = x \text{ for all } g \in \pi\} .$$

A π -equivariant map $f : X \rightarrow Y$ restricts to a map of the fixed point sets

$$\rho(f) : X^\pi \rightarrow Y^\pi .$$

Similarly for π -equivariant homotopies, with a fixed point function

$$\rho : [X, Y]_\pi \rightarrow [X^\pi, Y^\pi]$$

for any pointed π -spaces X, Y .

(iv) A π -space X has the *trivial* π -action if

$$gx = x \quad (x \in X, g \in \pi) ,$$

or equivalently $X^\pi = X$. Given a (pointed) space X let X also denote the (pointed) π -space defined by the trivial π -action on X .

(v) A π -space X is *free* if

$$X^\pi = \emptyset ,$$

or equivalently if the quotient map $X \rightarrow X/\pi$ is a regular covering projection.

(vi) A pointed π -space X is *semifree* if

$$X^\pi = \{*\} ,$$

or equivalently if $X \setminus \{*\}$ is a free π -space.

□

Example 4.2. (i) If X is a free π -space then X^+ is a semifree π -space.

(ii) For any π -spaces X, Y

$$(X \times Y)^\pi = X^\pi \times Y^\pi$$

so that if X is free then so is $X \times Y$.

(iii) For any pointed \mathbb{Z}_2 -spaces X, Y

$$(X \wedge Y)^\pi = X^\pi \wedge Y^\pi$$

so that if X is semifree then so is $X \wedge Y$.

□

Definition 4.3. (i) For any pointed spaces A, B let $\text{map}_*(A, B)$ be the space of pointed maps $A \rightarrow B$, with path component set

$$\pi_0(\text{map}_*(A, B)) = [A, B] .$$

(ii) For any pointed π -spaces A, B let $\text{map}_*^\pi(A, B)$ be the space of π -equivariant pointed maps $A \rightarrow B$, with path component set

$$\pi_0(\text{map}_*^\pi(A, B)) = [A, B]_\pi .$$

□

Proposition 4.4. (i) For any space X and π -space Y a π -equivariant map $f : X \rightarrow Y$ is the composite

$$f : X \longrightarrow Y^\pi \longrightarrow Y$$

of a map $X \rightarrow Y^\pi$ and the inclusion $Y^\pi \rightarrow Y$.

(ii) For a pointed space X and pointed π -space Y

$$\text{map}_*^\pi(X, Y) = \text{map}_*(X, Y^\pi) , [X, Y]_\pi = [X, Y^\pi] .$$

Proof. By construction.

□

We now specialize to the case $\pi = \mathbb{Z}_2$. In Chapter 7 we shall deal with arbitrary π .

Example 4.5. (i) A (pointed) \mathbb{Z}_2 -space X is a (pointed) space with an involution

$$T : X \rightarrow X .$$

(ii) For any pointed space Y regard $Y \wedge Y$ as a pointed \mathbb{Z}_2 -space by the transposition

$$T : Y \wedge Y \rightarrow Y \wedge Y ; (y_1, y_2) \mapsto (y_2, y_1) .$$

The inclusion of the fixed point set is just the diagonal map

$$\Delta : (Y \wedge Y)^{\mathbb{Z}_2} = Y \rightarrow Y \wedge Y ; y \mapsto (y, y) ,$$

so that for any pointed space X

$$[X, Y \wedge Y]_{\mathbb{Z}_2} = [X, Y] .$$

□

Definition 4.6. (i) An *inner product \mathbb{Z}_2 -space* V is an inner product space with a \mathbb{Z}_2 -action $T : V \rightarrow V$ which is an isometry. We shall write $|V|$ for the underlying inner product space. Let $S(V)$ and $P(V)$ denote the \mathbb{Z}_2 -spaces defined by the unit sphere $S(|V|)$ and the projective space $P(|V|)$ with the \mathbb{Z}_2 -action induced by T .

- (ii) An inner product space V can be regarded as an inner product \mathbb{Z}_2 -space with the trivial \mathbb{Z}_2 -action $T = 1 : V \rightarrow V$, so that $|V| = V$. The \mathbb{Z}_2 -actions on $S(V)$ and $P(V)$ are trivial.
- (iii) Let L be the $\mathbb{R}[\mathbb{Z}_2]$ -module \mathbb{R} with the involution -1 . For any inner product \mathbb{Z}_2 -space V there is then defined an inner product \mathbb{Z}_2 -space

$$LV = L \otimes_{\mathbb{R}} V, \quad T_{LV} : LV \rightarrow LV ; v \mapsto -T_V(v),$$

with $|LV| = |V|$. The \mathbb{Z}_2 -action on $S(LV)$ is non-trivial, and the \mathbb{Z}_2 -action on $P(LV)$ is trivial. For $V = \mathbb{R}^n$ write

$$LV^\infty = LS^n.$$

□

Proposition 4.7. *Let V be an inner product \mathbb{Z}_2 -space.*

- (i) *The inner product spaces*

$$\begin{aligned} V_+ &= V^{\mathbb{Z}_2} = \{x \in V \mid Tx = x\}, \\ V_- &= (LV)^{\mathbb{Z}_2} = \{x \in V \mid Tx = -x\} \subseteq V \end{aligned}$$

are such that as an inner product \mathbb{Z}_2 -space V has a decomposition

$$V = V_+ \oplus LV_- , \quad T(x_+, x_-) = (x_+, -x_-) .$$

- (ii) *The one-point compactification V^∞ of $V = V_+ \oplus LV_-$ is a pointed \mathbb{Z}_2 -space*

$$V^\infty = V_+^\infty \wedge LV_-^\infty$$

with

$$\begin{aligned} T : V^\infty &\rightarrow V^\infty ; (x_+, x_-) \mapsto (x_+, -x_-) , \\ (V^\infty)^{\mathbb{Z}_2} &= V_+^\infty \wedge LV_-^\infty = V_+^\infty \wedge \{0\}^\infty = V_+^\infty , \\ V^\infty/\mathbb{Z}_2 &= V_+^\infty/\mathbb{Z}_2 \wedge LV_-^\infty/\mathbb{Z}_2 = V_+^\infty \wedge sP(V_-) . \end{aligned}$$

- (iii) *The unit sphere $S(V)$ of $V = V_+ \oplus LV_-$ is a \mathbb{Z}_2 -space such that the homeomorphism of Proposition 2.11*

$$\begin{aligned} \lambda_{V_+, LV_-} : S(V_+) * S(LV_-) &\rightarrow S(V_+ \oplus LV_-) ; \\ (t, v_+, v_-) &\mapsto (v_+ \cos(\pi t/2), v_- \sin(\pi t/2)) \end{aligned}$$

is \mathbb{Z}_2 -equivariant, with

$$\begin{aligned} T : S(V_+) * S(LV_-) &\rightarrow S(V_+) * S(LV_-) ; (t, v_+, v_-) \mapsto (t, v_+, -v_-) , \\ S(V)^{\mathbb{Z}_2} &= S(V_+) * \emptyset = S(V_+) , \quad S(V)/\mathbb{Z}_2 = S(V_+) * P(V_-) . \end{aligned}$$

The induced \mathbb{Z}_2 -action on the projective space $P(V)$

$$T : P(V) \rightarrow P(V) ; [v_+, v_-] \mapsto [v_+, -v_-]$$

is such that

$$P(V)^{\mathbb{Z}_2} = P(V_+) \sqcup P(V_-) .$$

(iv) A subspace $W \subseteq V$ is \mathbb{Z}_2 -invariant (i.e. $TW = W$) if and only if $W = W_+ \oplus W_-$ for subspaces $W_+ \subseteq V_+$, $W_- \subseteq V_-$.

Proof. (i) It is clear that $V_+ \cap V_- = \{0\}$, and every $x \in V$ can be written as

$$x = \frac{x + Tx}{2} + \frac{x - Tx}{2} \in V_+ + V_- .$$

(ii)+(iii)+(iv) By construction. □

Proposition 4.8. Let X, Y be pointed \mathbb{Z}_2 -spaces.

(i) The pointed \mathbb{Z}_2 -space

$$S(L\mathbb{R})^+ \wedge X = X \vee X , \quad T(\pm x) = \mp Tx \quad (x \in X)$$

is the one-point union of two copies of X which are transposed by the \mathbb{Z}_2 -action. The map

$$S(L\mathbb{R})^+ \wedge X \rightarrow S(L\mathbb{R})^+ \wedge |X| ; \quad \begin{cases} (+, x) \mapsto (+, x) \\ (-, x) \mapsto (-, Tx) \end{cases}$$

is a \mathbb{Z}_2 -equivariant homeomorphism.

(ii) A \mathbb{Z}_2 -equivariant map $S(L\mathbb{R})^+ \wedge X \rightarrow Y$ is essentially the same as a map $|X| \rightarrow |Y|$, so that

$$[S(L\mathbb{R})^+ \wedge X, Y]_{\mathbb{Z}_2} = [|X|, |Y|] .$$

(iii) The \mathbb{Z}_2 -equivariant cofibration sequence

$$S(L\mathbb{R})^+ \rightarrow S^0 \rightarrow L\mathbb{R}^\infty \rightarrow \Sigma S(L\mathbb{R})^+ \rightarrow \dots$$

induces the exact sequence

$$\begin{aligned} \dots \rightarrow [\Sigma S(L\mathbb{R})^+ \wedge X, Y]_{\mathbb{Z}_2} &= [\Sigma X, Y] \rightarrow [L\mathbb{R}^\infty \wedge X, Y]_{\mathbb{Z}_2} \\ &\rightarrow [X, Y]_{\mathbb{Z}_2} \rightarrow [S(L\mathbb{R})^+ \wedge X, Y]_{\mathbb{Z}_2} = [|X|, |Y|] \end{aligned}$$

with $[X, Y]_{\mathbb{Z}_2} \rightarrow [|X|, |Y|]$ the function which forgets \mathbb{Z}_2 -equivariance, and

$$[L\mathbb{R}^\infty \wedge X, Y]_{\mathbb{Z}_2} \rightarrow [X, Y]_{\mathbb{Z}_2} ; \quad F \mapsto (x \mapsto F(0, x)) .$$

□

Example 4.9. The zero map $0_{L\mathbb{R}} : S^0 \rightarrow L\mathbb{R}^\infty$ induces a bijection

$$0_{L\mathbb{R}}^* : [L\mathbb{R}^\infty, L\mathbb{R}^\infty]_{\mathbb{Z}_2} \rightarrow [S^0, L\mathbb{R}^\infty]_{\mathbb{Z}_2}$$

which sends $1 : L\mathbb{R}^\infty \rightarrow L\mathbb{R}^\infty$ to $0_{L\mathbb{R}}$.

□

Proposition 4.10. (i) *For any inner product \mathbb{Z}_2 -spaces U, V*

$$P(U \oplus V)/P(V) = S(U)^+ \wedge_{\mathbb{Z}_2} V^\infty .$$

(ii) *The continuous bijection $V^\infty \rightarrow sS(V)$ of Proposition 2.8 (which is a homeomorphism for finite-dimensional V) is \mathbb{Z}_2 -equivariant, allowing the identifications*

$$V^\infty = sS(V) , (V^\infty)^{\mathbb{Z}_2} = sS(V_+) , V^\infty/\mathbb{Z}_2 = S(V_+) * sP(V_-) .$$

(iii) *For an inner product space V*

$$S(LV)^{\mathbb{Z}_2} = \emptyset , S(LV)/\mathbb{Z}_2 = P(V) .$$

For any \mathbb{Z}_2 -space X the \mathbb{Z}_2 -space $S(LV) \times X$ is free; for any pointed \mathbb{Z}_2 -space X the pointed \mathbb{Z}_2 -space $S(LV)^+ \wedge X$ is semifree.

Proof. (i) Immediate from the \mathbb{Z}_2 -equivariant cofibration sequence

$$S(V) \rightarrow S(U \oplus V) \rightarrow S(U)^+ \wedge V^\infty .$$

(ii)+(iii) By construction.

□

Proposition 4.11. *For any pointed \mathbb{Z}_2 -spaces X, Y and inner product space V there is defined a long exact sequence of abelian groups/pointed sets*

$$\begin{aligned} \dots \longrightarrow [\Sigma X, Y]_{\mathbb{Z}_2} &\xrightarrow{s_{LV}^*} [\Sigma S(LV)^+ \wedge X, Y]_{\mathbb{Z}_2} \xrightarrow{\alpha_{LV}^*} \\ &[LV^\infty \wedge X, Y]_{\mathbb{Z}_2} \xrightarrow{0_{LV}^*} [X, Y]_{\mathbb{Z}_2} . \end{aligned}$$

Proof. Immediate from the \mathbb{Z}_2 -equivariant cofibration sequence

$$S(LV)^+ \xrightarrow{s_{LV}} S^0 \xrightarrow{0_{LV}} LV^\infty \xrightarrow{\alpha_{LV}} \Sigma S(LV)^+ \longrightarrow \dots$$

of Proposition 2.17 (iii).

□

Terminology 4.12 The infinite-dimensional inner product space

$$\mathbb{R}(\infty) = \varinjlim_k \mathbb{R}^k$$

is denoted by $\mathbb{R}(\infty)$ to avoid confusion with the one-point compactification \mathbb{R}^∞ of \mathbb{R} . The unit sphere

$$S(\infty) = S(L\mathbb{R}(\infty)) = \varinjlim_k S(L\mathbb{R}^k)$$

is a contractible space with a free \mathbb{Z}_2 -action, with quotient the infinite-dimensional real projective space

$$P(\infty) = S(\infty)/\mathbb{Z}_2.$$

The unreduced suspension

$$sS(\infty) = \varinjlim_k sS(L\mathbb{R}^k) = \varinjlim_k (L\mathbb{R}^k)^\infty$$

is a contractible space with a non-free \mathbb{Z}_2 -action, such that there are two fixed points

$$sS(\infty)^{\mathbb{Z}_2} = S^0.$$

In dealing with inner product spaces V we interpret LV^∞ in the infinite-dimensional case $V = \mathbb{R}(\infty)$ to be

$$sS(\infty) = \varinjlim_{U \subset \mathbb{R}(\infty) \text{ finite-dimensional}} LU^\infty$$

(which is not compact, and in particular not the actual one-point compactification $L\mathbb{R}(\infty)^\infty$ of $L\mathbb{R}(\infty) = \varinjlim_U LU$).

□

Proposition 4.13. *For any pointed \mathbb{Z}_2 -spaces X, Y and inner product space V let*

$$\rho_V : \text{map}_*^{\mathbb{Z}_2}(X, LV^\infty \wedge Y) \rightarrow \text{map}_*(X^{\mathbb{Z}_2}, Y^{\mathbb{Z}_2})$$

be the fixed point map, and let

$$\begin{aligned} \rho_\infty &= \varinjlim_V \rho_V : \varinjlim_V \text{map}_*^{\mathbb{Z}_2}(X, LV^\infty \wedge Y) = \text{map}_*^{\mathbb{Z}_2}(X, L\mathbb{R}(\infty)^\infty \wedge Y) \\ &\rightarrow \text{map}_*(X^{\mathbb{Z}_2}, Y^{\mathbb{Z}_2}) \end{aligned}$$

be the map obtained by passing to the limit over finite-dimensional V .

(i) If X is a CW \mathbb{Z}_2 -complex then ρ_∞ is a fibration with contractible point inverses, and so induces isomorphisms in the homotopy groups. In particular, there is induced a bijection

$$\begin{aligned} \rho_\infty : \pi_0(\text{map}_*^{\mathbb{Z}_2}(X, L\mathbb{R}(\infty)^\infty \wedge Y)) &= [X, L\mathbb{R}(\infty)^\infty \wedge Y]_{\mathbb{Z}_2} \\ &\xrightarrow{\cong} \pi_0(\text{map}_*(X^{\mathbb{Z}_2}, Y^{\mathbb{Z}_2})) = [X^{\mathbb{Z}_2}, Y^{\mathbb{Z}_2}] . \end{aligned}$$

(ii) If the \mathbb{Z}_2 -action on X is trivial then ρ_V is a homeomorphism, with the inclusion

$$\sigma_V : Y^{\mathbb{Z}_2} = (LV^\infty \wedge Y)^{\mathbb{Z}_2} \rightarrow LV^\infty \wedge Y ; y \mapsto (0, y)$$

such that

$$\rho_V^{-1} = \sigma_V : \text{map}_*(X, Y^{\mathbb{Z}_2}) \rightarrow \text{map}_*^{\mathbb{Z}_2}(X, LV^\infty \wedge Y) ,$$

and

$$[X, LV^\infty \wedge Y]_{\mathbb{Z}_2} = [X, Y^{\mathbb{Z}_2}] .$$

The \mathbb{Z}_2 -equivariant homotopy group $[LV^\infty \wedge X, LV^\infty \wedge Y]_{\mathbb{Z}_2}$ fits into a direct sum system

$$[\Sigma S(LV)^+ \wedge X, LV^\infty \wedge Y]_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} [LV^\infty \wedge X, LV^\infty \wedge Y]_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} [X, Y^{\mathbb{Z}_2}]$$

with

$$\delta\gamma = 1 , \rho\sigma = 1 , \gamma\delta + \sigma\rho = 1 .$$

Here $\gamma = \alpha_{LV}^*$, and σ is defined by

$$\sigma : [X, Y^{\mathbb{Z}_2}] \rightarrow [LV^\infty \wedge X, LV^\infty \wedge Y]_{\mathbb{Z}_2} ; G \mapsto 1_{LV^\infty} \wedge \sigma_V G .$$

A \mathbb{Z}_2 -equivariant map $F : LV^\infty \wedge X \rightarrow LV^\infty \wedge Y$ with fixed point map $G = \rho(F) : X \rightarrow Y^{\mathbb{Z}_2}$ is such that F and $\sigma(G)$ agree on $0^+ \wedge X$, and $\delta(F)$ is defined to be the relative difference \mathbb{Z}_2 -equivariant map

$$\delta(F) = \delta(F, \sigma(G)) : \Sigma S(LV)^+ \wedge X \rightarrow LV^\infty \wedge Y ,$$

such that

$$F - \sigma(G) = \alpha_{LV}^* \delta(F, \sigma(G)) \in \text{im}(\alpha_{LV}^*) = \ker(\rho) .$$

Proof. (i) We repeat the argument of Sinha [59, p.277] (a space level version of Crabb [12, Lemma (A.1), p.60]). The fibre over a component of $\text{map}_*(X^{\mathbb{Z}_2}, Y^{\mathbb{Z}_2})$ is the space of \mathbb{Z}_2 -maps $f : X \rightarrow L\mathbb{R}(\infty)^\infty \wedge Y$ which are specified on $X^{\mathbb{Z}_2}$ by $\rho(f) : X^{\mathbb{Z}_2} \rightarrow Y^{\mathbb{Z}_2}$. We consider the effect on this mapping space of attaching \mathbb{Z}_2 -cells to X . There are two types: single cells with trivial \mathbb{Z}_2 -action D^n and pairs of cells with free \mathbb{Z}_2 -action $\mathbb{Z}_2 \times D^n$. For the single cells the extension of f is itself specified by the extension of $\rho(f)$. For the pairs of cells note that for any space W the \mathbb{Z}_2 -maps $g : \mathbb{Z}_2 \times W \rightarrow L\mathbb{R}(\infty)^\infty \wedge Y$ are just the maps $g : W \rightarrow L\mathbb{R}(\infty)^\infty \wedge Y$, and that $L\mathbb{R}(\infty)^\infty \wedge Y$ is contractible, so that the space of \mathbb{Z}_2 -maps $g : \mathbb{Z}_2 \times D^n \rightarrow L\mathbb{R}(\infty)^\infty \wedge Y$ extending a given \mathbb{Z}_2 -map $\partial g : \mathbb{Z}_2 \times S^{n-1} \rightarrow L\mathbb{R}(\infty)^\infty \wedge Y$ is contractible.

(ii) By Proposition 4.11 there is defined an exact sequence

$$\begin{aligned} \cdots \rightarrow [\Sigma S(LV)^+ \wedge X, LV^\infty \wedge Y]_{\mathbb{Z}_2} &\xrightarrow{\alpha_{LV}^*} [LV^\infty \wedge X, LV^\infty \wedge Y]_{\mathbb{Z}_2} \\ &\xrightarrow{0_{LV}^*} [X, LV^\infty \wedge Y]_{\mathbb{Z}_2} = [X, Y^{\mathbb{Z}_2}] \end{aligned}$$

□

4.2 The bi-degree

The degree is a \mathbb{Z} -valued homotopy invariant of a pointed map $V^\infty \rightarrow V^\infty$ for a finite-dimensional inner product space V , which defines an isomorphism

$$[V^\infty, V^\infty] \cong \mathbb{Z}.$$

Similarly, the bi-degree is a $\mathbb{Z} \oplus \mathbb{Z}$ -valued \mathbb{Z}_2 -homotopy invariant of a pointed \mathbb{Z}_2 -equivariant map $V^\infty \rightarrow V^\infty$ for a finite-dimensional inner product \mathbb{Z}_2 -space V , which defines an isomorphism

$$[V^\infty, V^\infty]_{\mathbb{Z}_2} \cong \mathbb{Z} \oplus \mathbb{Z}.$$

We start by recollecting the main properties of the degree.

Definition 4.14. Let V be a finite-dimensional inner product space. The *degree* of a pointed map $F : V^\infty \rightarrow V^\infty$ is the homotopy invariant defined by

$$\text{degree}(F) = F_*(1) \in \mathbb{Z},$$

with

$$F_* : \dot{H}_{\dim(V)}(V^\infty) = \mathbb{Z} \rightarrow \dot{H}_{\dim(V)}(V^\infty) = \mathbb{Z} .$$

□

Example 4.15. (i) The identity $1 : V^\infty \rightarrow V^\infty; v \mapsto v$ has $\text{degree}(1) = 1$.
(ii) The constant map $\infty : V^\infty \rightarrow V^\infty; v \mapsto \infty$ has $\text{degree}(\infty) = 0$.

□

Proposition 4.16. *Let V be a finite-dimensional inner product space.*

(i) *The degree function*

$$[V^\infty, V^\infty] \rightarrow \begin{cases} \mathbb{Z} & \text{if } \dim(V) > 0 \\ \{0, 1\} & \text{if } \dim(V) = 0 \end{cases} ; F \mapsto \text{degree}(F)$$

is a bijection of pointed sets, which is an isomorphism of abelian groups for $\dim(V) > 0$.

(ii) *For any pointed map $F : V^\infty \rightarrow V^\infty$ and finite-dimensional inner product space W the pointed map*

$$F' = F \wedge 1 : (V \oplus W)^\infty = V^\infty \wedge W^\infty \rightarrow (V \oplus W)^\infty = V^\infty \wedge W^\infty$$

has

$$\text{degree}(F') = \text{degree}(F) \in \mathbb{Z} .$$

□

We now move on to the bi-degree itself.

Proposition 4.17. *Let V be a non-zero inner product space.*

(i) *The cellular $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes of $S(LV)$ and LV^∞ are given by*

$$\begin{aligned} C^{\text{cell}}(S(LV)) : C(S(LV))_{\dim(V)-1} &= \mathbb{Z}[\mathbb{Z}_2] \xrightarrow{1+(-)^{\dim(V)-1}T} \\ C(S(LV))_{\dim(V)-2} &= \mathbb{Z}[\mathbb{Z}_2] \xrightarrow{1+(-)^{\dim(V)}T} \dots \xrightarrow{1-T} C(S(LV))_0 = \mathbb{Z}[\mathbb{Z}_2] , \\ \dot{C}^{\text{cell}}(LV^\infty) &= S^{\dim(V)}(\mathbb{Z}, (-1)^{\dim(V)}) \end{aligned}$$

where $(\mathbb{Z}, \pm 1)$ denotes the $\mathbb{Z}[\mathbb{Z}_2]$ -module \mathbb{Z} with $T \in \mathbb{Z}_2$ acting by ± 1 .

(ii) *A $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map*

$$e : \dot{C}^{\text{cell}}(\Sigma S(LV)^+) = SC^{\text{cell}}(S(LV)) \rightarrow \dot{C}^{\text{cell}}(LV^\infty)$$

induces a $\mathbb{Z}[\mathbb{Z}_2]$ -module morphism

$$e_* : H_{\dim(V)-1}(S(LV)) = \begin{cases} (\mathbb{Z}, (-)^{\dim(V)}) & \text{if } \dim(V) > 1 \\ \mathbb{Z}[\mathbb{Z}_2] & \text{if } \dim(V) = 1 \end{cases}$$

$$\rightarrow \dot{H}_{\dim(V)}(LV^\infty) = (\mathbb{Z}, (-)^{\dim(V)})$$

$$\text{with } \begin{cases} e_*(1) \in 2\mathbb{Z} \\ e_*(1) = -e_*(T) \in \mathbb{Z}. \end{cases}$$

Proof. (i) By construction.

(ii) The generator $1 \in H_{\dim(V)-1}(S(LV)) = \mathbb{Z}$ is represented by

$$1 + (-)^{\dim(V)}T \in C(S(LV))_{\dim(V)-1} = \mathbb{Z}[\mathbb{Z}_2].$$

A $\mathbb{Z}[\mathbb{Z}_2]$ -module morphism

$$e : C(S(LV))_{\dim(V)-1} = \mathbb{Z}[\mathbb{Z}_2] \rightarrow (\mathbb{Z}, (-)^{\dim(V)})$$

is of the form

$$e(a + bT) = na + (-)^{\dim(V)}nb \quad (a, b \in \mathbb{Z})$$

for some $n \in \mathbb{Z}$, and

$$\begin{cases} e(1 + (-)^{\dim(V)}T) = 2n & \text{if } \dim(V) > 1 \\ e(1) = -e(T) = n & \text{if } \dim(V) = 1. \end{cases}$$

□

Definition 4.18. (i) The *semidegree* of a $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$e : \dot{C}^{cell}(\Sigma S(LV)^+) \rightarrow \dot{C}^{cell}(LV^\infty)$$

is

$$\text{semidegree}(e) = \begin{cases} e_*(1)/2 \in \mathbb{Z} & \text{if } \dim(V) > 1 \\ e_*(1) = -e_*(T) \in \mathbb{Z} & \text{if } \dim(V) = 1. \end{cases}$$

(ii) Let V, W be finite-dimensional inner product spaces. The *semidegree* of a \mathbb{Z}_2 -equivariant pointed map

$$E : \Sigma S(LV)^+ \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty$$

is the semidegree of the induced $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map, that is

$$\text{semidegree}(E) = \begin{cases} E_*(1)/2 \in \mathbb{Z} & \text{if } \dim(V) > 1 \\ E_*(1) = -E_*(T) \in \mathbb{Z} & \text{if } \dim(V) = 1 \end{cases}$$

with

$$\begin{aligned} E_* : \dot{H}_{\dim(V)+\dim(W)}(\Sigma S(LV)^+ \wedge W^\infty) &= \begin{cases} \mathbb{Z} & \text{if } \dim(V) > 1 \\ \mathbb{Z}[\mathbb{Z}_2] & \text{if } \dim(V) = 1 \end{cases} \\ \rightarrow \dot{H}_{\dim(V)+\dim(W)}(LV^\infty \wedge W^\infty) &= \mathbb{Z} . \end{aligned}$$

□

Proposition 4.19. (i) *The function*

$$H_0(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(\dot{C}^{cell}(\Sigma S(LV)^+), \dot{C}^{cell}(LV^\infty))) \rightarrow \mathbb{Z} ; e \mapsto \text{semidegree}(e)$$

is an isomorphism of abelian groups.

(ii) *The semidegree of a \mathbb{Z}_2 -equivariant pointed map*

$$E : \Sigma S(LV)^+ \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty$$

is a \mathbb{Z}_2 -equivariant homotopy invariant, such that for any $j, k \geq 0$ the semidegrees of E and the \mathbb{Z}_2 -equivariant pointed map defined by

$$\begin{aligned} \Sigma^{j,k} E : \Sigma S(LV \oplus L\mathbb{R}^k)^+ \wedge (W \oplus \mathbb{R}^j)^\infty \\ \xrightarrow{j_{LV, L\mathbb{R}} \wedge 1} \Sigma S(LV)^+ \wedge (L\mathbb{R}^k)^\infty \wedge W^\infty \wedge (\mathbb{R}^j)^\infty \\ \xrightarrow{E \wedge 1} LV^\infty \wedge (L\mathbb{R}^k)^\infty \wedge W^\infty \wedge (\mathbb{R}^j)^\infty = (LV \oplus L\mathbb{R}^k)^\infty \wedge (W \oplus \mathbb{R}^j)^\infty \end{aligned}$$

are related by

$$\text{semidegree}(\Sigma^{j,k} E) = \text{semidegree}(E) \in \mathbb{Z} .$$

Proof. (i) For each $n \in \mathbb{Z}$ construct a chain map e with semidegree n as in the proof of Proposition 4.17.

(ii) The only non-trivial case is for $V = \mathbb{R}$, $W = \{0\}$, $j = 0$, $k = 1$. Consider the commutative diagram

$$\begin{array}{ccc} \dot{H}_2(\Sigma S(L\mathbb{R}^2)^+) = \mathbb{Z} & \xrightarrow{\Sigma^{0,1} E_*} & \dot{H}_2((L\mathbb{R}^2)^\infty) = \mathbb{Z} \\ \downarrow 1 - T & & \downarrow 1 \\ \dot{H}_1(\Sigma S(L\mathbb{R})^+) = \mathbb{Z}[\mathbb{Z}_2] & \xrightarrow{E_*} & \dot{H}_1(L\mathbb{R}^\infty) = \mathbb{Z} \end{array}$$

The evaluations of the two composites on the generator $1 \in \dot{H}_2(\Sigma S(L\mathbb{R}^2)^+) = \mathbb{Z}$ are

$$\begin{aligned} E_*(1 - T)(1) &= 2 \text{semidegree}(E) , \\ \Sigma^{1,0} E_*(1) &= 2 \text{semidegree}(\Sigma^{1,0} E) \in \mathbb{Z} , \end{aligned}$$

so

$$\text{semidegree}(\Sigma^{0,1}E) = \text{semidegree}(E) \in \mathbb{Z} .$$

□

Definition 4.20. Given a \mathbb{Z}_2 -equivariant pointed map

$$F : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty$$

define the pointed map of the \mathbb{Z}_2 -fixed point sets

$$G = \rho(F) : (LV^\infty \wedge W^\infty)^{\mathbb{Z}_2} = W^\infty \rightarrow W^\infty ;$$

$$w \mapsto \begin{cases} w' & \text{if } F(0, w) = (0, w') \\ \infty & \text{if } F(0, w) = \infty . \end{cases}$$

The \mathbb{Z}_2 -equivariant pointed map

$$\sigma(G) = 1 \wedge G : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty ; (v, w) \mapsto (v, G(w))$$

is such that

$$F(0, w) = (0, G(w)) = \sigma(G)(0, w) \in LV^\infty \wedge W^\infty \quad (w \in W) ,$$

so that the relative difference is defined, a \mathbb{Z}_2 -equivariant pointed map

$$\delta(F, \sigma(G)) : \Sigma S(LV)^+ \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty .$$

The *bi-degree* of F is defined by

$$\begin{aligned} \text{bi-degree}(F) &= (\text{semidegree}(\delta(F, \sigma(G))), \text{degree}(G)) \\ &\in \begin{cases} \mathbb{Z} \oplus \mathbb{Z} & \text{if } \dim(W) > 0 \\ \mathbb{Z} \times \{0, 1\} & \text{if } \dim(W) = 0 . \end{cases} \end{aligned}$$

□

Proposition 4.21. (i) *The bi-degree of a \mathbb{Z}_2 -equivariant pointed map $F : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty$ is*

$$\begin{aligned} \text{bi-degree}(F) &= \left(\frac{\text{degree}(F) - \text{degree}(G)}{2}, \text{degree}(G) \right) \\ &\in \begin{cases} \mathbb{Z} \oplus \mathbb{Z} & \text{if } \dim(W) > 0 \\ \mathbb{Z} \times \{0, 1\} & \text{if } \dim(W) = 0 \end{cases} \end{aligned}$$

with $G = \rho(F) : W^\infty \rightarrow W^\infty$. In particular

$$\deg(F) \equiv \deg(G) \pmod{2} .$$

(ii) *The degree of a \mathbb{Z}_2 -equivariant pointed map $F : LV^\infty \rightarrow LV^\infty$ is odd*

$$\deg(F) \equiv 1 \pmod{2} ,$$

and for any pointed map $G : W^\infty \rightarrow W^\infty$ the bi-degree of the \mathbb{Z}_2 -equivariant pointed map

$$F \wedge G : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty$$

is

$$\begin{aligned} \text{bi-degree}(F \wedge G) &= \left(\frac{(\deg(F) - 1)}{2} \deg(G), \deg(G) \right) \\ &\in \begin{cases} \mathbb{Z} \oplus \mathbb{Z} & \text{if } \dim(W) > 0 \\ \mathbb{Z} \times \{0, 1\} & \text{if } \dim(W) = 0 . \end{cases} \end{aligned}$$

(iii) *If $F_1, F_2 : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty$ are \mathbb{Z}_2 -equivariant pointed maps with*

$$\rho(F_1) = \rho(F_2) = G : W^\infty \rightarrow W^\infty$$

then

$$\text{bi-degree}(F_1) - \text{bi-degree}(F_2) = \left(\frac{\deg(F_1) - \deg(F_2)}{2}, 0 \right) \in \mathbb{Z} \oplus \mathbb{Z} .$$

In particular

$$\deg(F_1) \equiv \deg(F_2) \pmod{2} ,$$

with

$$\deg(F_1) - \deg(F_2) = 2 \text{ semidegree}(\delta(F_1, F_2)) \in \mathbb{Z} .$$

(iv) *The degree, semidegree and bi-degree functions define an isomorphism of direct sum systems of abelian groups*

$$\begin{array}{ccc} [\Sigma S(LV)^+ \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} & \xrightarrow[\cong]{\text{semidegree}} & \mathbb{Z} \\ \delta \updownarrow \gamma = \alpha_{LV}^* & & (1 \ 0) \updownarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ [LV^\infty \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} & \xrightarrow[\cong]{\text{bi-degree}} & \mathbb{Z} \oplus \mathbb{Z} \\ \sigma \updownarrow \rho = s_{LV}^* & & \begin{pmatrix} 0 \\ 1 \end{pmatrix} \updownarrow (0 \ 1) \\ [W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} = [W^\infty, W^\infty] & \xrightarrow[\cong]{\text{degree}} & \mathbb{Z} \end{array}$$

for non-zero W . The forgetful map

$$[LV^\infty \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} = \mathbb{Z} \oplus \mathbb{Z} \rightarrow [LV^\infty \wedge W^\infty, LV^\infty \wedge W^\infty] = \mathbb{Z}$$

sends a \mathbb{Z}_2 -equivariant map $F : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty$ with bi-degree $(a, b) \in \mathbb{Z} \oplus \mathbb{Z}$ to a map F with

$$\text{degree}(F) = 2a + b \in \mathbb{Z} .$$

For $W = \{0\}$ the functions define a bijection of products of pointed sets

$$\begin{array}{ccc} [\Sigma S(LV)^+, LV^\infty]_{\mathbb{Z}_2} & \xrightarrow[\cong]{\text{semidegree}} & \mathbb{Z} \\ \delta \uparrow \downarrow \alpha_{LV}^* & & \uparrow \downarrow \\ [LV^\infty, LV^\infty]_{\mathbb{Z}_2} & \xrightarrow[\cong]{\text{bi-degree}} & \mathbb{Z} \times \{0, 1\} \\ \sigma \uparrow \downarrow s_{LV}^* & & \uparrow \downarrow \\ [S^0, LV^\infty]_{\mathbb{Z}_2} = [S^0, S^0] & \xrightarrow[\cong]{\text{degree}} & \{0, 1\} \end{array}$$

Proof. (i) The map $\alpha_{LV} : LV^\infty \rightarrow \Sigma S(LV)^+$ induces

$$\begin{aligned} (\alpha_{LV})_* &= \begin{cases} 1 \\ 1 - T \end{cases} : \dot{H}_{\dim(V)}(LV^\infty) = \mathbb{Z} \\ \rightarrow \dot{H}_{\dim(V)}(\Sigma S(LV)^+) &= H_{\dim(V)-1}(S(LV)) = \begin{cases} \mathbb{Z} \\ \mathbb{Z}[\mathbb{Z}_2] \end{cases} \quad \text{if } \begin{cases} \dim(V) > 1 \\ \dim(V) = 1 \end{cases} . \end{aligned}$$

The composite

$$F - \sigma(G) : LV^\infty \wedge W^\infty \xrightarrow{\alpha_{LV} \wedge 1} \Sigma S(LV)^+ \wedge W^\infty \xrightarrow{\delta(F, \sigma(G))} LV^\infty \wedge W^\infty$$

has degree

$$\text{degree}(\delta(F, \sigma(G))(\alpha_{LV} \wedge 1)) = \text{degree}(F) - \text{degree}(\sigma(G))$$

so

$$\begin{aligned} \text{semidegree}(\delta(F, \sigma(G))) &= \frac{\text{degree}(F) - \text{degree}(\sigma(G))}{2} \\ &= \frac{\text{degree}(F) - \text{degree}(G)}{2} \in \mathbb{Z} . \end{aligned}$$

(ii) Immediate from (i).

(iii) Taking $X = Y = S^0$ in Proposition 4.11 (ii), we have a direct sum system

$$\begin{array}{ccc}
[\Sigma S(LV)^+ \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} & \xrightleftharpoons[\delta]{\gamma} & \\
[LV^\infty \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} & \xrightleftharpoons[\sigma]{\rho} & [W^\infty, W^\infty] .
\end{array}$$

The function

$$\begin{aligned}
(\alpha_{LV}^*, \sigma) &: [\Sigma S(LV)^+ \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} \times [W^\infty, W^\infty] \\
&\longrightarrow [LV^\infty \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2}
\end{aligned}$$

is an isomorphism such that

$$\begin{aligned}
F &= (\alpha_{LV}^*, \sigma)(\delta(F, \sigma(G)), G) \\
&\in \text{im}((\alpha_{LV}^*, \sigma) : [\Sigma S(LV)^+ \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2} \times [W^\infty, W^\infty] \\
&\xrightarrow{\cong} [LV^\infty \wedge W^\infty, LV^\infty \wedge W^\infty]_{\mathbb{Z}_2}) .
\end{aligned}$$

□

Example 4.22. Given a \mathbb{Z}_2 -equivariant pointed map $f : LV^\infty \rightarrow LV^\infty$ and a pointed map $g : W^\infty \rightarrow W^\infty$ define the \mathbb{Z}_2 -equivariant pointed map

$$F = f \wedge g : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty ; (v, w) \mapsto (f(v), g(w)) .$$

The \mathbb{Z}_2 -fixed point map is

$$G = \rho(F) : W^\infty \rightarrow W^\infty ; w \mapsto \begin{cases} g(w) & \text{if } f(0) = 0 \\ \infty & \text{if } f(0) = \infty \end{cases}$$

and

$$\sigma(G) : LV^\infty \wedge W^\infty \rightarrow LV^\infty \wedge W^\infty ; (v, w) \mapsto \begin{cases} (v, g(w)) & \text{if } f(0) = 0 \\ \infty & \text{if } f(0) = \infty , \end{cases}$$

so that

$$\begin{aligned}
\text{bi-degree}(F) &= (\text{semidegree}(\delta(F, \sigma(G))), \text{degree}(G)) \\
&= \left(\frac{\text{degree}(F) - \text{degree}(G)}{2}, \text{degree}(G) \right) \\
&= \begin{cases} \left(\frac{(\text{degree}(f) - 1)\text{degree}(g)}{2}, \text{degree}(g) \right) & \text{if } f(0) = 0 \\ \left(\frac{\text{degree}(f)\text{degree}(g)}{2}, 0 \right) & \text{if } f(0) = \infty . \end{cases}
\end{aligned}$$

□

Example 4.23. The \mathbb{Z}_2 -equivariant pointed map defined for any $\lambda \in \mathbb{R}$ by

$$F = \lambda : L\mathbb{R}^\infty \rightarrow L\mathbb{R}^\infty ; x \mapsto x/\lambda$$

has

$$G = \rho(F) : (L\mathbb{R}^\infty)^{\mathbb{Z}_2} = \{0, \infty\} \rightarrow \{0, \infty\} ; \infty \mapsto \infty, 0 \mapsto \begin{cases} 0 & \text{if } \lambda \neq 0 \\ \infty & \text{if } \lambda = 0, \end{cases}$$

$$\text{bi-degree}(F) = (\text{semidegree } \delta(F, \sigma(G)), \text{degree } G) = \begin{cases} (0, 1) & \text{if } \lambda > 0 \\ (0, 0) & \text{if } \lambda = 0 \\ (-1, 1) & \text{if } \lambda < 0 \end{cases}$$

$$\in [L\mathbb{R}^\infty, L\mathbb{R}^\infty]_{\mathbb{Z}_2} = [\Sigma S(L\mathbb{R})^+, L\mathbb{R}^\infty]_{\mathbb{Z}_2} \times [S^0, S^0] = \mathbb{Z} \times \{0, 1\}.$$

□

4.3 Stable \mathbb{Z}_2 -equivariant homotopy theory

Definition 4.24. The *stable \mathbb{Z}_2 -equivariant homotopy group* of pointed \mathbb{Z}_2 -spaces X, Y is defined by

$$\{X; Y\}_{\mathbb{Z}_2} = \varinjlim_V [V^\infty \wedge X, V^\infty \wedge Y]_{\mathbb{Z}_2}$$

with the direct limit running over all the finite-dimensional inner product \mathbb{Z}_2 -spaces V .

□

Remark 4.25. In dealing with

$$\{X; Y\}_{\mathbb{Z}_2} = \varinjlim_j \varinjlim_k [(L\mathbb{R}^j \oplus \mathbb{R}^k)^\infty \wedge X, (L\mathbb{R}^j \oplus \mathbb{R}^k)^\infty \wedge Y]_{\mathbb{Z}_2}$$

there is no loss of generality in only considering $j = k$. Noting that $(L\mathbb{R}^k \oplus \mathbb{R}^k)^\infty \wedge X$ is \mathbb{Z}_2 -equivariantly homeomorphic to

$$\Sigma^{k,k} X = S^k \wedge S^k \wedge X, \quad T : \Sigma^{k,k} X \rightarrow \Sigma^{k,k} X ; (s, t, x) \mapsto (t, s, Tx)$$

(see Definition 4.40 below for details) we can identify

$$\{X; Y\}_{\mathbb{Z}_2} = \varinjlim_k [\Sigma^{k,k} X, \Sigma^{k,k} Y]_{\mathbb{Z}_2} .$$

□

Example 4.26. For any non-zero inner product space V Proposition 4.21 (ii) gives an isomorphism of direct sum systems of abelian groups

$$\begin{array}{ccc} \{\Sigma S(LV)^+; LV^\infty\}_{\mathbb{Z}_2} & \xrightarrow[\cong]{\text{semidegree}} & \mathbb{Z} \\ \delta \updownarrow \gamma = \alpha_{LV}^* & & (1 \ 0) \updownarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ \{LV^\infty; LV^\infty\}_{\mathbb{Z}_2} = \{S^0; S^0\}_{\mathbb{Z}_2} & \xrightarrow[\cong]{\text{bi-degree}} & \mathbb{Z} \oplus \mathbb{Z} \\ \sigma \updownarrow \rho = s_{LV}^* & & \begin{pmatrix} 0 \\ 1 \end{pmatrix} \updownarrow \begin{pmatrix} 0 & 1 \end{pmatrix} \\ \{S^0; LV^\infty\}_{\mathbb{Z}_2} = \{S^0; S^0\} & \xrightarrow[\cong]{\text{degree}} & \mathbb{Z} \end{array}$$

□

Proposition 4.27. *Let X, Y be pointed \mathbb{Z}_2 -spaces.*

(i) *Given a pointed map $F : |X| \rightarrow |Y|$ define the \mathbb{Z}_2 -equivariant map*

$$F \vee TFT : S(L\mathbb{R})^+ \wedge X \rightarrow Y ; \quad \begin{cases} (+, x) \mapsto F(x) \\ (-, x) \mapsto TFT(x) . \end{cases}$$

The functions

$$\begin{aligned} \{|X|; |Y|\} &\rightarrow \{S(L\mathbb{R})^+ \wedge X; Y\}_{\mathbb{Z}_2} ; \quad F \mapsto F \vee TFT , \\ \{S(L\mathbb{R})^+ \wedge X; Y\}_{\mathbb{Z}_2} &\rightarrow \{|X|; |Y|\} ; \quad G \mapsto G|_{(+, X)} \end{aligned}$$

are inverse isomorphisms of abelian groups. The \mathbb{Z}_2 -equivariant cofibration sequence

$$S(L\mathbb{R})^+ \rightarrow S^0 \rightarrow L\mathbb{R}^\infty \rightarrow \Sigma S(L\mathbb{R})^+ \rightarrow \dots$$

induces the exact sequence

$$\begin{aligned} \dots &\rightarrow \{\Sigma S(L\mathbb{R})^+ \wedge X, Y\}_{\mathbb{Z}_2} = \{\Sigma X, Y\} \rightarrow \{L\mathbb{R}^\infty \wedge X, Y\}_{\mathbb{Z}_2} \\ &\rightarrow \{X, Y\}_{\mathbb{Z}_2} \rightarrow \{S(L\mathbb{R})^+ \wedge X, Y\}_{\mathbb{Z}_2} = \{|X|, |Y|\} \rightarrow \dots . \end{aligned}$$

(ii) *Given a pointed map $F : |X| \rightarrow |Y|$ define the \mathbb{Z}_2 -equivariant map*

$$\begin{aligned} F + TFT : L\mathbb{R}^\infty \wedge X &\rightarrow L\mathbb{R}^\infty \wedge (S(L\mathbb{R})^+ \wedge Y) ; \\ (t, x) &\mapsto \begin{cases} (2t, (+, F(x))) & \text{if } 0 \leq t \leq 1/2 \\ (2t - 1, (-, TFT(x))) & \text{if } 1/2 \leq t \leq 1 . \end{cases} \end{aligned}$$

The function

$$q : \{|X|; |Y|\} \rightarrow \{X; S(L\mathbb{R})^+ \wedge Y\}_{\mathbb{Z}_2} ; F \mapsto F + TFT$$

is an isomorphism of abelian groups.

Proof. (i) By Proposition 4.8 (i)

$$[S(L\mathbb{R})^+ \wedge V^\infty \wedge X, V^\infty \wedge Y]_{\mathbb{Z}_2} = [|V^\infty| \wedge |X|, |V^\infty| \wedge |Y|] .$$

The exact sequence is the stable version of 4.8 (iii).

(ii) Use the projection $p : S(L\mathbb{R})^+ \wedge Y \rightarrow S(L\mathbb{R})^+ \wedge_{\mathbb{Z}_2} Y$ and the homeomorphism

$$|Y| \rightarrow S(L\mathbb{R})^+ \wedge_{\mathbb{Z}_2} Y ; y \mapsto (+, y)$$

to define a morphism

$$p_* : \{X; S(L\mathbb{R})^+ \wedge Y\}_{\mathbb{Z}_2} \rightarrow \{|X|; S(L\mathbb{R})^+ \wedge_{\mathbb{Z}_2} Y\} = \{|X|; |Y|\}$$

such that $p_*q = 1$, $qp_* = 1$.

□

Given pointed \mathbb{Z}_2 -spaces A, B and a \mathbb{Z}_2 -equivariant pointed map $F : A \rightarrow B$ let $G : B \rightarrow \mathcal{C}(F)$ be the inclusion in the mapping cone, and let $H : \mathcal{C}(F) \rightarrow \Sigma A$ be the projection, so that

$$A \xrightarrow{F} B \xrightarrow{G} \mathcal{C}(F) \xrightarrow{H} \Sigma A \xrightarrow{\Sigma F} \Sigma B \longrightarrow \dots$$

is a \mathbb{Z}_2 -equivariant cofibration sequence. By analogy with the nonequivariant case (3.2):

Proposition 4.28. *For any pointed \mathbb{Z}_2 -space X there is induced a Barratt-Puppe exact sequence of stable \mathbb{Z}_2 -equivariant homotopy groups*


$$\begin{aligned} \dots \longrightarrow \{\Sigma X; \mathcal{C}(F)\}_{\mathbb{Z}_2} &\xrightarrow{H} \{X; A\}_{\mathbb{Z}_2} \xrightarrow{F} \{X; B\}_{\mathbb{Z}_2} \\ &\xrightarrow{G} \{X; \mathcal{C}(F)\}_{\mathbb{Z}_2} \xrightarrow{H} \{X; \Sigma A\}_{\mathbb{Z}_2} \longrightarrow \dots \end{aligned}$$

□

By analogy with the braid of stable homotopy groups of Proposition 3.3 :

Proposition 4.29. *For any inner product \mathbb{Z}_2 -spaces U, V and pointed \mathbb{Z}_2 -spaces X, Y there is defined a commutative braid of exact sequences of stable*


homotopy groups



with

$$\begin{aligned} A_1 &= \{\Sigma X; LV^\infty \wedge Y\}_{\mathbb{Z}_2}, \quad A_2 = \{\Sigma S(U \oplus V)^+ \wedge X; V^\infty \wedge Y\}_{\mathbb{Z}_2}, \\ A_3 &= \{S(U \oplus V)^+ \wedge X; V^\infty \wedge Y\}_{\mathbb{Z}_2}, \quad A_4 = \{\Sigma S(U)^+ \wedge X; Y\}_{\mathbb{Z}_2}. \end{aligned}$$

Proof. These are the Barratt-Puppe exact sequences (4.28) determined by the commutative braid of \mathbb{Z}_2 -equivariant cofibrations



□

Proposition 4.30. *Let Y be a pointed \mathbb{Z}_2 -space.*

(i) *For any pointed \mathbb{Z}_2 -space X the forgetful map $\{X; Y\}_{\mathbb{Z}_2} \rightarrow \{|X|; |Y|\}$ fits into a long exact sequence*

$$\cdots \rightarrow \{L\mathbb{R}^\infty \wedge X; Y\}_{\mathbb{Z}_2} \rightarrow \{X; Y\}_{\mathbb{Z}_2} \rightarrow \{|X|; |Y|\} \rightarrow \{L\mathbb{R}^\infty \wedge X; \Sigma Y\}_{\mathbb{Z}_2} \rightarrow \cdots$$

(ii) *For any CW \mathbb{Z}_2 -spectrum X the fixed point map*

$$\rho : \{X; L\mathbb{R}(\infty)^\infty \wedge Y\}_{\mathbb{Z}_2} \rightarrow \{X^{\mathbb{Z}_2}; Y^{\mathbb{Z}_2}\}$$

is an isomorphism, and the fixed point map

$$\begin{aligned} \rho : \{X; Y\}_{\mathbb{Z}_2} &\rightarrow \{X^{\mathbb{Z}_2}; Y^{\mathbb{Z}_2}\}; \\ (F : V^\infty \wedge X &\rightarrow V^\infty \wedge Y) \mapsto (\rho(F) : V_+^\infty \wedge X^{\mathbb{Z}_2} \rightarrow V_+^\infty \wedge Y^{\mathbb{Z}_2}) \end{aligned}$$

fits into a long exact sequence

$$\begin{aligned} \cdots \rightarrow \{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} &\rightarrow \{X; Y\}_{\mathbb{Z}_2} \xrightarrow{\rho} \{X^{\mathbb{Z}_2}; Y^{\mathbb{Z}_2}\} \\ &\rightarrow \{X; \Sigma S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \rightarrow \cdots \end{aligned}$$

(iii) For any semifree CW \mathbb{Z}_2 -spectrum X

$$\{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} = \{X; Y\}_{\mathbb{Z}_2}, \quad \{X^{\mathbb{Z}_2}; Y^{\mathbb{Z}_2}\} = 0.$$

(iv) For CW \mathbb{Z}_2 -spectrum X with the trivial \mathbb{Z}_2 -action the fixed point map $\rho : \{X; Y\}_{\mathbb{Z}_2} \rightarrow \{X; Y^{\mathbb{Z}_2}\}$ is a surjection which is split by the map $\sigma : \{X; Y^{\mathbb{Z}_2}\} \rightarrow \{X; Y\}_{\mathbb{Z}_2}$ induced by

$$\sigma : Y^{\mathbb{Z}_2} = (L\mathbb{R}(\infty)^\infty \wedge Y)^{\mathbb{Z}_2} \rightarrow L\mathbb{R}(\infty)^\infty \wedge Y; \quad y \mapsto (0, y).$$

Thus $\{X; Y\}_{\mathbb{Z}_2}$ fits into a direct sum system

$$\{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \{X; Y\}_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \{X; Y^{\mathbb{Z}_2}\}$$

(v) The direct sum system in (iv) is natural with respect to \mathbb{Z}_2 -equivariant maps. A stable \mathbb{Z}_2 -equivariant map $F : V^\infty \wedge Y \rightarrow V^\infty \wedge Y'$ induces

$$\begin{aligned} F &= \begin{pmatrix} 1 \wedge F & \delta(F i_Y, i_{Y'} \rho(F)) \\ 0 & \rho(F) \end{pmatrix} : \\ \{X; Y\}_{\mathbb{Z}_2} &= \{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \oplus \{X; Y^{\mathbb{Z}_2}\} \\ &\rightarrow \{X; Y'\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge Y'\}_{\mathbb{Z}_2} \oplus \{X; Y'^{\mathbb{Z}_2}\} \end{aligned}$$

with $i_Y : Y^{\mathbb{Z}_2} \rightarrow Y$, $i_{Y'} : Y'^{\mathbb{Z}_2} \rightarrow Y'$ the inclusions.

Proof. (i) Immediate from the \mathbb{Z}_2 -equivariant cofibration sequence

$$S(L\mathbb{R})^+ \xrightarrow{s_{L\mathbb{R}}} S^0 \xrightarrow{0_{L\mathbb{R}}} L\mathbb{R}^\infty \xrightarrow{\alpha_{L\mathbb{R}}} \Sigma S(L\mathbb{R})^+ \longrightarrow \cdots$$

of Proposition 2.17 (iii) and the isomorphism

$$\{X; Y\} \rightarrow \{S(L\mathbb{R})^+ \wedge X; Y\}_{\mathbb{Z}_2} ; F \mapsto F \vee TF$$

given by Proposition 4.8.

(ii) Combine Proposition 4.13 (i) with the long exact sequence

$$\begin{aligned} \cdots \rightarrow \{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} &\rightarrow \{X; Y\}_{\mathbb{Z}_2} \rightarrow \{X; L\mathbb{R}(\infty)^\infty \wedge Y\}_{\mathbb{Z}_2} \\ &\rightarrow \{X; \Sigma S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \rightarrow \cdots \end{aligned}$$

induced by the \mathbb{Z}_2 -equivariant cofibration sequence

$$S(\infty)^+ \xrightarrow{s_{S(\infty)}} S^0 \xrightarrow{0_{L\mathbb{R}(\infty)}} L\mathbb{R}(\infty)^\infty \xrightarrow{\alpha_{L\mathbb{R}(\infty)}} \Sigma S(\infty)^+ \longrightarrow \cdots$$

given by Proposition 2.17 (iii).

(iii) This is the stable version of Proposition 4.13, with each of the fixed point maps

$$\rho : [LV^\infty \wedge W^\infty \wedge X, L\mathbb{R}(\infty)^\infty \wedge LV^\infty \wedge W^\infty \wedge Y]_{\mathbb{Z}_2} \rightarrow [W^\infty \wedge X^{\mathbb{Z}_2}, W^\infty \wedge Y^{\mathbb{Z}_2}]$$

is an isomorphism.

(iv) By construction.

□

Proposition 4.11 has an analogue for the stable \mathbb{Z}_2 -equivariant homotopy groups:

Definition 4.31. (i) The *reduced stable \mathbb{Z}_2 -equivariant homotopy and cohomotopy* groups of a pointed \mathbb{Z}_2 -space X are

$$\tilde{\omega}_n^{\mathbb{Z}_2}(X) = \{S^n; X\}_{\mathbb{Z}_2}, \quad \tilde{\omega}_{\mathbb{Z}_2}^n(X) = \{X; S^n\}_{\mathbb{Z}_2} \quad (n \in \mathbb{Z}).$$

(ii) The *absolute stable \mathbb{Z}_2 -equivariant homotopy and cohomotopy* groups of a \mathbb{Z}_2 -space X are

$$\omega_n^{\mathbb{Z}_2}(X) = \tilde{\omega}_n^{\mathbb{Z}_2}(X^+), \quad \omega_{\mathbb{Z}_2}^n(X) = \tilde{\omega}_{\mathbb{Z}_2}^n(X^+) \quad (n \in \mathbb{Z}).$$

(iii) The *stable \mathbb{Z}_2 -equivariant homotopy groups of spheres* are

$$\omega_n^{\mathbb{Z}_2} = \omega_n^{\mathbb{Z}_2}(\{*\}) = \omega_{\mathbb{Z}_2}^{-n}(\{*\}) = \{S^n; S^0\}_{\mathbb{Z}_2} \quad (n \in \mathbb{Z}).$$

□

Stable \mathbb{Z}_2 -equivariant maps arise from double covers:

Proposition 4.32. (i) (Adams [3, p.511]) *For any semifree pointed \mathbb{Z}_2 -space the projection $f : X \rightarrow X/\mathbb{Z}_2$ induces a transfer $\mathbb{Z}[\mathbb{Z}_2]$ -module chain*

map

$$f^! : \dot{C}(X/\mathbb{Z}_2) \rightarrow \dot{C}(X) ; y \mapsto \sum_{x \in f^{-1}(y)} x$$

of the reduced singular chain complexes. If X is a finite pointed CW \mathbb{Z}_2 -complex the Umkehr map $f^!$ is induced by a stable \mathbb{Z}_2 -equivariant Umkehr map

$$F : (LV \oplus W)^\infty \wedge X/\mathbb{Z}_2 \rightarrow (LV \oplus W)^\infty \wedge X .$$

(ii) Let A be a pointed \mathbb{Z}_2 -space with a \mathbb{Z}_2 -equivariant map $\epsilon : A \rightarrow [0, \infty)$ such that $\epsilon^{-1}(0) = \{*\} \subset A$, with $*$ $\in A$ the base point. For any inner product space V there exists an open \mathbb{Z}_2 -equivariant embedding

$$e = (g, f) : LV \times X \hookrightarrow LV \times X/\mathbb{Z}_2$$

with a \mathbb{Z}_2 -equivariant Umkehr map

$$F : LV^\infty \wedge X \rightarrow LV^\infty \wedge X/\mathbb{Z}_2 .$$

(iii) (Adams [3, Thms. 5.3, 5.4, 5.5]) Let X be a semifree finite pointed CW \mathbb{Z}_2 -complex, let $f : X \rightarrow X/\mathbb{Z}_2$ be the projection, and let $F \in \{X/\mathbb{Z}_2; X\}_{\mathbb{Z}_2}$ be the stable \mathbb{Z}_2 -equivariant Umkehr map of (i). For any pointed finite CW complex Y the functions

$$\begin{aligned} \{X/\mathbb{Z}_2; Y\} &\rightarrow \{X; Y\}_{\mathbb{Z}_2} ; G \mapsto Gf , \\ \{Y; X/\mathbb{Z}_2\} &\rightarrow \{Y; X\}_{\mathbb{Z}_2} ; H \mapsto FH , \\ \varinjlim_V [V^\infty \wedge X, V^\infty \wedge Y]_{\mathbb{Z}_2} &\rightarrow \{X; Y\}_{\mathbb{Z}_2} \end{aligned}$$

are isomorphisms, with V running over all finite-dimensional inner product spaces (with the trivial \mathbb{Z}_2 -action).

Proof. (i) There exist finite-dimensional inner product spaces V, W and a \mathbb{Z}_2 -equivariant map

$$g : LV \oplus W \times X \rightarrow LV \oplus W$$

with the property that

$$e = (g, f) : LV \oplus W \times X \rightarrow LV \oplus W \times X/\mathbb{Z}_2 ; (v, w, x) \mapsto (g(v, w), f(x))$$

restricts to an open \mathbb{Z}_2 -equivariant embedding

$$e| : LV \oplus W \times (X \setminus \{x_0\}) \rightarrow LV \oplus W \times X/\mathbb{Z}_2$$

and as x approaches $*$ then $g(LV \oplus W \times \{x\})$ approaches $\{0\} \subset LV \oplus W$. The construction of Definition 3.9 gives a stable \mathbb{Z}_2 -equivariant Umkehr map

$$F : (LV \oplus W)^\infty \wedge X/\mathbb{Z}_2 \rightarrow (LV \oplus W)^\infty \wedge X$$

representing a stable \mathbb{Z}_2 -equivariant homotopy class $F \in \{X/\mathbb{Z}_2; X\}_{\mathbb{Z}_2}$ inducing the transfer chain map $F = f^! : \dot{C}(X/\mathbb{Z}_2) \rightarrow \dot{C}(X)$.

(ii) Define the semifree pointed \mathbb{Z}_2 -space $X = S(LV)^+ \wedge A$ and let

$$f : X \rightarrow X/\mathbb{Z}_2 = S(LV)^+ \wedge_{\mathbb{Z}_2} A$$

be the projection. The \mathbb{Z}_2 -equivariant map

$$g : LV \times X \rightarrow LV ; (u, (v, a)) \mapsto \epsilon(a)(v + \frac{u}{1 + \|u\|})$$

has the property that as $a \in A$ approaches the base point $*$ in A then $g(LV \times (S(LV)^+ \wedge \{a\}))$ approaches $0 \in LV$. Then

$$e = (g, f) : LV \times X \hookrightarrow LV \times X/\mathbb{Z}_2 ; (u, (v, a)) \mapsto (g(u, (v, a)), f(v, a))$$

is an open \mathbb{Z}_2 -equivariant embedding with \mathbb{Z}_2 -equivariant Umkehr map

$$F : LV^\infty \wedge X \rightarrow LV^\infty \wedge X/\mathbb{Z}_2 ;$$

$$(u, (v, a)) \mapsto \begin{cases} (\frac{u-v}{1-\|u-v\|}, (v, a)) & \text{if } (v, a) \in X, \|u-v\| < 1 \\ (\frac{u+v}{1-\|u+v\|}, (-v, a)) & \text{if } (v, a) \in X, \|u+v\| < 1 \\ * & \text{otherwise .} \end{cases}$$

(iii) See [3].

Example 4.33. The special case $Y = S^i$ of 4.32 (ii) gives identifications

$$\tilde{\omega}_i^{\mathbb{Z}_2}(X) = \tilde{\omega}_i(X/\mathbb{Z}_2) , \tilde{\omega}_{\mathbb{Z}_2}^i(X) = \tilde{\omega}^i(X/\mathbb{Z}_2)$$

for any finite pointed semifree CW \mathbb{Z}_2 -complex. □

A ' \mathbb{Z}_2 -spectrum' is the \mathbb{Z}_2 -equivariant analogue of a spectrum :

Definition 4.34. (i) A \mathbb{Z}_2 -spectrum $\underline{X} = \{X(V) | V\}$ is a sequence of pointed \mathbb{Z}_2 -spaces $X(V)$ indexed by finite-dimensional inner product \mathbb{Z}_2 -spaces V , with structure maps

$$(V^\perp)^\infty \wedge X(V) \rightarrow X(W)$$

defined whenever $V \subseteq W$, where $V^\perp \subseteq W$ is the orthogonal complement of V in W . For $n \in \mathbb{Z}$ let

$$\pi_n^{\mathbb{Z}_2}(\underline{X}) = \begin{cases} \varinjlim_V [\Sigma^n V^\infty, X(V)]_{\mathbb{Z}_2} & \text{if } n \geq 0 \\ \varinjlim_V [V^\infty, \Sigma^{-n} X(V)]_{\mathbb{Z}_2} & \text{if } n \leq -1. \end{cases}$$

(ii) A \mathbb{Z}_2 -spectrum \underline{X} is *connective* if

$$\pi_n^{\mathbb{Z}_2}(\underline{X}) = 0 \text{ for } n \leq -1.$$

In particular, this is the case if $\{W^\infty; X(V)\}_{\mathbb{Z}_2} = 0$ for every proper \mathbb{Z}_2 -invariant subspace $W \subset V$.

(iii) The \underline{X} -coefficient homology groups of a \mathbb{Z}_2 -space Y are

$$\underline{X}_n^{\mathbb{Z}_2}(Y) = \pi_n^{\mathbb{Z}_2}(\underline{X} \wedge Y^+) = \varinjlim_V [\Sigma^n V^\infty, X(V) \wedge Y^+]_{\mathbb{Z}_2}.$$

The *reduced \underline{X} -coefficient homology* groups of a pointed \mathbb{Z}_2 -space Y are

$$\tilde{\underline{X}}_n^{\mathbb{Z}_2}(Y) = \pi_n^{\mathbb{Z}_2}(\underline{X} \wedge Y).$$

If \underline{X} is connective then $\underline{X}_n^{\mathbb{Z}_2}(Y) = 0$ for $n \leq -1$, and similarly in the reduced case.

(iv) The *fixed point spectrum* of a \mathbb{Z}_2 -spectrum \underline{X} is the spectrum

$$\underline{X}^{\mathbb{Z}_2} = \{X(U)^{\mathbb{Z}_2} \mid U\},$$

with U running over finite-dimensional inner product spaces.

(v) The *homotopy fixed point \mathbb{Z}_2 -spectrum* of \underline{X} is the \mathbb{Z}_2 -spectrum

$$\underline{X}^{h\mathbb{Z}_2} = L\mathbb{R}(\infty)^\infty \wedge \underline{X} = \{L\mathbb{R}(\infty)^\infty \wedge X(V) \mid V\}$$

with

$$\begin{aligned} (\tilde{\underline{X}}^{h\mathbb{Z}_2})_n^{\mathbb{Z}_2}(Y) &= \varinjlim_V [\Sigma^n V^\infty, L\mathbb{R}(\infty)^\infty \wedge X(V) \wedge Y]_{\mathbb{Z}_2} \\ &= \varinjlim_V [\Sigma^n (V^\infty)^{\mathbb{Z}_2}, X(V)^{\mathbb{Z}_2} \wedge Y^{\mathbb{Z}_2}] \end{aligned}$$

(by Proposition 4.13). There are evident fixed point maps

$$\rho : \tilde{\underline{X}}_n^{\mathbb{Z}_2}(Y) \rightarrow (\tilde{\underline{X}}^{h\mathbb{Z}_2})_n^{\mathbb{Z}_2}(Y).$$

(vi) The *homotopy orbit \mathbb{Z}_2 -spectrum* of \underline{X} is the \mathbb{Z}_2 -spectrum

$$\underline{X}_{h\mathbb{Z}_2} = S(\infty)^+ \wedge \underline{X} = \{S(\infty)^+ \wedge X(V) \mid V\}$$

with

$$(\tilde{X}_{h\mathbb{Z}_2})_n^{\mathbb{Z}_2}(Y) = \varinjlim_V [\Sigma^n V^\infty, S(\infty)^+ \wedge X(V) \wedge Y]_{\mathbb{Z}_2} .$$

□

Proposition 4.35. *For any \mathbb{Z}_2 -spectrum \underline{X} and pointed \mathbb{Z}_2 -space Y there is a long exact sequence*

$$\cdots \rightarrow (\tilde{X}_{h\mathbb{Z}_2})_n^{\mathbb{Z}_2}(Y) \rightarrow \tilde{X}_n^{\mathbb{Z}_2}(Y) \xrightarrow{\rho} (\tilde{X}^{h\mathbb{Z}_2})_n^{\mathbb{Z}_2}(Y) \rightarrow (\tilde{X}_{h\mathbb{Z}_2})_{n-1}^{\mathbb{Z}_2}(Y) \rightarrow \cdots .$$

Proof. The \mathbb{Z}_2 -equivariant cofibration sequence of pointed \mathbb{Z}_2 -spaces

$$S(\infty)^+ \rightarrow S^0 \rightarrow L\mathbb{R}(\infty)^\infty \rightarrow \Sigma S(\infty)^+ \rightarrow \cdots$$

determines a cofibration sequence of \mathbb{Z}_2 -spectra

$$\underline{X}_{h\mathbb{Z}_2} \rightarrow \underline{X} \rightarrow \underline{X}^{h\mathbb{Z}_2} \rightarrow \Sigma \underline{X}_{h\mathbb{Z}_2} \rightarrow \cdots .$$

□

Example 4.36. (i) The special case $Y = S^0$ of Proposition 4.35 gives a long exact sequence

$$\cdots \rightarrow \pi_n^{\mathbb{Z}_2}(\underline{X}_{h\mathbb{Z}_2}) \rightarrow \pi_n^{\mathbb{Z}_2}(\underline{X}) \xrightarrow{\rho} \pi_n^{\mathbb{Z}_2}(\underline{X}^{h\mathbb{Z}_2}) \rightarrow \pi_{n-1}^{\mathbb{Z}_2}(\underline{X}_{h\mathbb{Z}_2}) \rightarrow \cdots .$$

(ii) The suspension \mathbb{Z}_2 -spectrum of a pointed \mathbb{Z}_2 -space X

$$\underline{X} = \{V^\infty \wedge X \mid V\}$$

is such that

$$\begin{aligned} \pi_n^{\mathbb{Z}_2}(\underline{X}) &= \tilde{\omega}_n^{\mathbb{Z}_2}(X) , \\ \pi_n^{\mathbb{Z}_2}(\underline{X}_{h\mathbb{Z}_2}) &= \tilde{\omega}_n(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) , \\ \pi_n^{\mathbb{Z}_2}(\underline{X}^{h\mathbb{Z}_2}) &= \tilde{\omega}_n(X^{\mathbb{Z}_2}) \quad (4.13) . \end{aligned}$$

By Proposition 4.29 the long exact sequence given by (i)

$$\begin{aligned} \cdots \rightarrow \tilde{\omega}_n(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) \rightarrow \tilde{\omega}_n^{\mathbb{Z}_2}(X) &\xrightarrow{\rho} \tilde{\omega}_n(X^{\mathbb{Z}_2}) \\ &\rightarrow \tilde{\omega}_{n-1}(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) \rightarrow \cdots \end{aligned}$$

breaks up into split short exact sequences

$$0 \rightarrow \tilde{\omega}_n(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) \rightarrow \tilde{\omega}_n^{\mathbb{Z}_2}(X) \xrightarrow{\rho} \tilde{\omega}_n(X^{\mathbb{Z}_2}) \rightarrow 0 ,$$

so that

$$\begin{aligned}\pi_n^{\mathbb{Z}_2}(\underline{X}) &= \pi_n^{\mathbb{Z}_2}(\underline{X}_{h\mathbb{Z}_2}) \oplus \pi_n^{\mathbb{Z}_2}(\underline{X}^{h\mathbb{Z}_2}) \\ &= \tilde{\omega}_n^{\mathbb{Z}_2}(X) = \tilde{\omega}_n(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) \oplus \tilde{\omega}_n(X^{\mathbb{Z}_2}).\end{aligned}$$

If the \mathbb{Z}_2 -action on X is trivial then

$$\pi_n^{\mathbb{Z}_2}(\underline{X}_{h\mathbb{Z}_2}) = \tilde{\omega}_n(P(\mathbb{R}(\infty))^+ \wedge X), \quad \pi_n^{\mathbb{Z}_2}(\underline{X}^{h\mathbb{Z}_2}) = \tilde{\omega}_n(X^{\mathbb{Z}_2}) = \tilde{\omega}_n(X).$$

If the \mathbb{Z}_2 -action on X is semifree then

$$\pi_n^{\mathbb{Z}_2}(\underline{X}_{h\mathbb{Z}_2}) = \tilde{\omega}_n(X/\mathbb{Z}_2), \quad \pi_n^{\mathbb{Z}_2}(\underline{X}^{h\mathbb{Z}_2}) = \tilde{\omega}_n(X^{\mathbb{Z}_2}) = 0.$$

(iii) The sphere \mathbb{Z}_2 -spectrum $\underline{S} = \{V^\infty\}$ is the suspension \mathbb{Z}_2 -spectrum of S^0 , with

$$\pi_n^{\mathbb{Z}_2}(\underline{S}) = \omega_n^{\mathbb{Z}_2}, \quad \pi_n^{\mathbb{Z}_2}(\underline{S}_{h\mathbb{Z}_2}) = \omega_n(P(\infty)), \quad \pi_n^{\mathbb{Z}_2}(\underline{S}^{h\mathbb{Z}_2}) = \omega_n.$$

The \mathbb{Z}_2 -spectrum $\underline{S} = \{V^\infty \mid V\}$ is connective, so that

$$\tilde{\omega}_n^{\mathbb{Z}_2}(X) = \tilde{\omega}_{\mathbb{Z}_2}^n(X) = 0 \text{ for } n < 0.$$

□

Definition 4.37. (i) A spectrum $\underline{A} = \{A(U) \mid U\}$ (with U running over finite-dimensional inner product spaces) determines a \mathbb{Z}_2 -spectrum

$$\underline{A} = \{A(V) \mid V\}$$

with $A(V) = A(|V|)$.

(ii) A \mathbb{Z}_2 -spectrum $\underline{X} = \{X(V) \mid V\}$ determines a spectrum

$$|\underline{X}| = \{X(U) \mid U\}.$$

(iii) A \mathbb{Z}_2 -spectrum \underline{X} is *split* if the fixed point spectrum $\underline{X}^{\mathbb{Z}_2}$ dominates $|\underline{X}|$, i.e. if there exists a map $\zeta : |\underline{X}| \rightarrow \underline{X}^{\mathbb{Z}_2}$ such that $j\zeta \simeq 1 : |\underline{X}| \rightarrow |\underline{X}|$, with $j : \underline{X}^{\mathbb{Z}_2} \rightarrow |\underline{X}|$ the inclusion.

□

Proposition 4.38. (Costenoble [9][Thms. 1.5,1.6]) *If \underline{X} is a split \mathbb{Z}_2 -spectrum then*

$$1 \wedge j\zeta : \underline{X}^{h\mathbb{Z}_2} = S(\infty)^+ \wedge \underline{X} \rightarrow |\underline{X}^{h\mathbb{Z}_2}| = S(\infty)^+ \wedge |\underline{X}|$$

is a homotopy equivalence of \mathbb{Z}_2 -spectra. Thus for any pointed \mathbb{Z}_2 -space Y which is semifree

$$(\tilde{X}^{h\mathbb{Z}_2})_*(Y) = |\tilde{X}^{h\mathbb{Z}_2}|_{\mathbb{Z}_2}^*(Y) = |\tilde{X}|_*(Y/\mathbb{Z}_2).$$

Furthermore, the natural map of \mathbb{Z}_2 -spectra

$$S(L\mathbb{R})^+ \wedge \underline{X}^{\mathbb{Z}_2} \rightarrow \underline{X}^{h\mathbb{Z}_2}$$

induces isomorphisms

$$\pi_*^{\mathbb{Z}_2}(S(L\mathbb{R})^+ \wedge \underline{X}^{\mathbb{Z}_2}) = \pi_*(\underline{X}^{\mathbb{Z}_2}) \cong \pi_*^{\mathbb{Z}_2}(\underline{X}^{h\mathbb{Z}_2})$$

and there is defined an exact sequence

$$\cdots \rightarrow |\tilde{X}|_n(Y/\mathbb{Z}_2) \rightarrow \tilde{X}_n^{\mathbb{Z}_2}(Y) \rightarrow (\tilde{X}^{\mathbb{Z}_2})_n^{\mathbb{Z}_2}(Y) \rightarrow |\tilde{X}|_{n-1}(Y/\mathbb{Z}_2) \rightarrow \cdots$$

□

Example 4.39. The sphere \mathbb{Z}_2 -spectrum $\underline{S} = \{V^\infty \mid V\}$ is split: the map

$$\zeta : |\underline{S}| \rightarrow \underline{S}^{\mathbb{Z}_2}$$

defined by

$$\zeta(U) = 1 : |\underline{S}(U)| = U^\infty \rightarrow \underline{S}^{\mathbb{Z}_2}(U) = U^\infty$$

is such that $j\zeta = 1 : |\underline{S}| \rightarrow |\underline{S}|$.

□

Definition 4.40. For any inner product space V regard $V \oplus V$ as an inner product \mathbb{Z}_2 -space via the *transposition involution* \mathbb{Z}_2 -action

$$T : V \oplus V \rightarrow V \oplus V ; (u, v) \mapsto (v, u) .$$

The \mathbb{Z}_2 -equivariant linear isomorphism

$$\kappa_V : LV \oplus V \cong V \oplus V ; (u, v) \mapsto (u + v, -u + v)$$

has inverse

$$\kappa_V^{-1} : V \oplus V \cong LV \oplus V ; (x, y) \mapsto \left(\frac{x - y}{2}, \frac{x + y}{2} \right)$$

and induces a \mathbb{Z}_2 -equivariant homeomorphism

$$\kappa_V : LV^\infty \wedge V^\infty \cong V^\infty \wedge V^\infty .$$

The restriction of κ_V is a \mathbb{Z}_2 -equivariant homeomorphism

$$\kappa_V| : (LV \setminus \{0\}) \times V \cong (V \oplus V) \setminus \Delta(V)$$

which induces a \mathbb{Z}_2 -equivariant homeomorphism

$$\kappa_V| : (LV \setminus \{0\})^\infty \wedge V^\infty \cong ((V \oplus V) \setminus \Delta(V))^\infty$$

where $(LV \setminus \{0\})^\infty \cong \Sigma S(LV)^+$. \square

Proposition 4.41. (i) *For any pointed \mathbb{Z}_2 -spaces X, Y and inner product spaces U, V there is defined a commutative braid of exact sequences of stable \mathbb{Z}_2 -equivariant homotopy groups*

$$\begin{array}{ccccc}
 & & \xrightarrow{\alpha_{LV}} & & \\
 A_1 & & \{X; S(LV)^+ \wedge Y\}_{\mathbb{Z}_2} & \xrightarrow{s_{LV}} & \{S(LU)^+ \wedge X; Y\}_{\mathbb{Z}_2} \\
 & \searrow & & \nearrow s_{LU}^* & \\
 & A_2 & & \{X; Y\}_{\mathbb{Z}_2} & \\
 & \nearrow & \xrightarrow{0_{LU}^*} & \searrow 0_{LV} & \\
 A_4 & & \{LU^\infty \wedge X; Y\}_{\mathbb{Z}_2} & \xrightarrow{\alpha_{LU}^*} & \{X; LV^\infty \wedge Y\}_{\mathbb{Z}_2} \\
 & \searrow & & \nearrow & \\
 & A_3 & & &
 \end{array}$$

with

$$\begin{aligned}
 A_1 &= \{\Sigma X; LV^\infty \wedge Y\}_{\mathbb{Z}_2}, \quad A_2 = \{\Sigma S(LU \oplus LV)^+ \wedge X; LV^\infty \wedge Y\}_{\mathbb{Z}_2}, \\
 A_3 &= \{S(LU \oplus LV)^+ \wedge X; LV^\infty \wedge Y\}_{\mathbb{Z}_2}, \quad A_4 = \{\Sigma S(LU)^+ \wedge X; Y\}_{\mathbb{Z}_2}.
 \end{aligned}$$

The morphism

$$0_{LV} : \{X; X \wedge X\}_{\mathbb{Z}_2} \rightarrow \{X; LV^\infty \wedge X \wedge X\}_{\mathbb{Z}_2} = \{V^\infty \wedge X; V^\infty \wedge LV^\infty \wedge X \wedge X\}_{\mathbb{Z}_2}$$

sends Δ_X to $0_{LV} \wedge \Delta_X = (\kappa_V^{-1} \wedge 1) \Delta_{V^\infty \wedge X}$.

(ii) For any pointed space X and any pointed \mathbb{Z}_2 -space Y

$$\{X; Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \oplus \{X; Y^{\mathbb{Z}_2}\},$$

with a direct sum system

$$\{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \xrightleftharpoons[\delta]{\gamma} \{X; Y\}_{\mathbb{Z}_2} \xrightleftharpoons[\sigma]{\rho} \{X; Y^{\mathbb{Z}_2}\}.$$

The map γ is induced by the projection $s_{L\mathbb{R}(\infty)} : S(\infty)^+ \rightarrow S^0$. The map δ is defined by sending a \mathbb{Z}_2 -equivariant map

$$F : U^\infty \wedge LV^\infty \wedge X \rightarrow U^\infty \wedge LV^\infty \wedge Y$$

to the stable \mathbb{Z}_2 -equivariant homotopy class of the relative difference of F and

$$\begin{aligned} \sigma\rho(F) : U^\infty \wedge LV^\infty \wedge X &\rightarrow U^\infty \wedge LV^\infty \wedge Y ; \\ (u, v, x) &\mapsto (w, v, y) \quad (F(u, 0, x) = (w, 0, y)) , \end{aligned}$$

that is

$$\begin{aligned} \delta(F) &= \delta(F, \sigma\rho(F)) \in \varinjlim_{U, V} \{U^\infty \wedge \Sigma S(LV)^+ \wedge X; U^\infty \wedge LV^\infty \wedge Y\}_{\mathbb{Z}_2} \\ &= \varinjlim_V \{X; S(LV)^+ \wedge Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} . \end{aligned}$$

(iii) For any inner product space U , any pointed space X and any pointed \mathbb{Z}_2 -space Y there is defined a long exact sequence

$$\begin{aligned} \dots &\longrightarrow \{\Sigma S(LU \oplus L\mathbb{R}(\infty))^+ \wedge X; L\mathbb{R}(\infty)^\infty \wedge Y\}_{\mathbb{Z}_2} \\ &\longrightarrow \{LU^\infty \wedge X; Y\}_{\mathbb{Z}_2} \xrightarrow{\rho} \{X; Y^{\mathbb{Z}_2}\} \\ &\longrightarrow \{S(LU \oplus L\mathbb{R}(\infty))^+ \wedge X; L\mathbb{R}(\infty)^\infty \wedge Y\}_{\mathbb{Z}_2} \longrightarrow \dots \end{aligned}$$

with ρ defined by the fixed points of the \mathbb{Z}_2 -action, and

$$\begin{aligned} \{S(LU \oplus L\mathbb{R}(\infty))^+ \wedge X; L\mathbb{R}(\infty)^\infty \wedge Y\}_{\mathbb{Z}_2} \\ = \{LU^\infty \wedge X; \Sigma S(LU \oplus L\mathbb{R}(\infty))^+ \wedge Y\}_{\mathbb{Z}_2} . \end{aligned}$$

Proof. (i) Immediate from the \mathbb{Z}_2 -equivariant cofibration sequences

$$\begin{aligned} S(LU)^+ &\rightarrow S^0 \rightarrow LU^\infty \rightarrow \Sigma S(LU)^+ \rightarrow \dots , \\ S(LV)^+ &\rightarrow S^0 \rightarrow LV^\infty \rightarrow \Sigma S(LV)^+ \rightarrow \dots . \end{aligned}$$

(ii) This is the \mathbb{Z}_2 -equivariant version of the braid of Proposition 3.3, noting that the fixed point function

$$\rho : \{X; L\mathbb{R}(\infty)^\infty \wedge Y\}_{\mathbb{Z}_2} \rightarrow \{X; Y^{\mathbb{Z}_2}\}$$

is an isomorphism by Proposition 4.30. The identification

$$0_{LV} \wedge \Delta_X = (\kappa_V^{-1} \wedge 1) \Delta_{V^\infty \wedge X}$$

is immediate from the commutative triangle of \mathbb{Z}_2 -equivariant maps

$$\begin{array}{ccc}
 V^\infty & \xrightarrow{0_{LV}} & LV^\infty \wedge V^\infty \\
 & \searrow \Delta_V & \nearrow \kappa_V \\
 & V^\infty \wedge V^\infty &
 \end{array}$$

(iii) The \mathbb{Z}_2 -equivariant cofibration sequence

$$S(\infty)^+ \xrightarrow{S L\mathbb{R}(\infty)} S^0 \xrightarrow{0_{L\mathbb{R}(\infty)}} L\mathbb{R}(\infty)^\infty$$

induces an exact sequence

$$\begin{aligned}
 \cdots \rightarrow \{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} &\xrightarrow{\gamma} \{X; Y\}_{\mathbb{Z}_2} \\
 &\xrightarrow{\rho} \{X; L\mathbb{R}(\infty)^\infty \wedge Y\}_{\mathbb{Z}_2} = \{X; Y^{\mathbb{Z}_2}\} \rightarrow \cdots
 \end{aligned}$$

□

Example 4.42. Let Y be a pointed \mathbb{Z}_2 -space.

(i) The splitting of Proposition 4.41 (ii)

$$\{X; Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \oplus \{X; Y^{\mathbb{Z}_2}\}$$

decomposes the stable \mathbb{Z}_2 -equivariant homotopy theory of Y according to the actual fixed point space $Y^{\mathbb{Z}_2}$ and the \mathbb{Z}_2 -homotopy orbit space $S(\infty)^+ \wedge_{\mathbb{Z}_2} Y$. In particular,

$$\tilde{\omega}_*^{\mathbb{Z}_2}(Y) = \tilde{\omega}_*(Y^{\mathbb{Z}_2}) \oplus \tilde{\omega}_*(S(\infty)^+ \wedge_{\mathbb{Z}_2} Y) .$$

(ii) If Y is semifree

$$\begin{aligned}
 Y^{\mathbb{Z}_2} &= \{*\} , \quad \tilde{\omega}_*(Y^{\mathbb{Z}_2}) = 0 , \quad S(\infty)^+ \wedge_{\mathbb{Z}_2} Y \simeq Y/\mathbb{Z}_2 , \\
 \tilde{\omega}_*^{\mathbb{Z}_2}(Y) &= \tilde{\omega}_*(S(\infty)^+ \wedge_{\mathbb{Z}_2} Y) = \tilde{\omega}_*(Y/\mathbb{Z}_2) .
 \end{aligned}$$

while if the \mathbb{Z}_2 -action on Y is trivial then

$$\begin{aligned}
 Y^{\mathbb{Z}_2} &= Y , \quad S(\infty)^+ \wedge_{\mathbb{Z}_2} Y = P(\infty)^+ \wedge Y , \\
 \tilde{\omega}_*^{\mathbb{Z}_2}(Y) &= \tilde{\omega}_*(Y) \oplus \tilde{\omega}_*(P(\infty)^+ \wedge Y) .
 \end{aligned}$$

□


Remark 4.43. The decomposition of the stable \mathbb{Z}_2 -equivariant homotopy groups of a \mathbb{Z}_2 -space X as a sum of ordinary stable homotopy groups of the fixed point space $X^{\mathbb{Z}_2}$ and the homotopy orbit space $S(\infty) \times_{\mathbb{Z}_2} X$ of the \mathbb{Z}_2 -action

$$\omega_*^{\mathbb{Z}_2}(X) = \omega_*(X^{\mathbb{Z}_2}) \oplus \omega_*(S(\infty) \times_{\mathbb{Z}_2} X)$$

is the special case $G = \mathbb{Z}_2$ of a general decomposition of the stable G -equivariant homotopy groups of a G -space for an arbitrary compact Lie (e.g. finite) group G – see Segal [56], tom Dieck [65], Hauschild [22], May et al. [41].

□

Example 4.44. (i) The braid of Proposition 4.41 (i) for $U = \mathbb{R}^{i-j}$, $X = S^j$, $V = \mathbb{R}(\infty)$ can be written




with

$$\begin{aligned} \tilde{\omega}_{i,j}(Y) &= \{S^j \wedge LS^{i-j}; Y\}_{\mathbb{Z}_2}, \\ A_1 &= \tilde{\omega}_{j+1}(Y^{\mathbb{Z}_2}), \quad A_2 = \tilde{\omega}_{i,j}(S(\infty)^+ \wedge Y), \\ A_3 &= \tilde{\omega}_{i-1,j-1}(S(\infty)^+ \wedge Y), \quad A_4 = \{\Sigma^{j+1}S(L\mathbb{R}^{i-j})^+; Y\}_{\mathbb{Z}_2}. \end{aligned}$$

For $i = j$ $\tilde{\omega}_{j,j}(Y) = \tilde{\omega}_j^{\mathbb{Z}_2}(Y)$.

(ii) The braid of Proposition 4.41 (i) for $U = \mathbb{R}$ can be written



with


$$\begin{aligned} A_1 &= \{\Sigma X; LV^\infty \wedge Y\}_{\mathbb{Z}_2}, \quad A_2 = \{\Sigma S(LV \oplus L\mathbb{R})^+ \wedge X; LV^\infty \wedge Y\}_{\mathbb{Z}_2}, \\ A_3 &= \{S(LV \oplus L\mathbb{R})^+ \wedge X; LV^\infty \wedge Y\}_{\mathbb{Z}_2}, \quad A_4 = \{\Sigma X; Y\}. \end{aligned}$$

The braid includes the exact sequence

$$\cdots \rightarrow \{\Sigma X; Y\}_{\mathbb{Z}_2} \xrightarrow{\alpha_{L\mathbb{R}}^*} \{L\mathbb{R}^\infty \wedge X; Y\}_{\mathbb{Z}_2} \xrightarrow{0_{L\mathbb{R}}^*} \{X; Y\}_{\mathbb{Z}_2} \xrightarrow{s_{L\mathbb{R}}^*} \{X; Y\} \rightarrow \cdots$$

with $s_{L\mathbb{R}}^* : \{X; Y\}_{\mathbb{Z}_2} \rightarrow \{S(L\mathbb{R})^+ \wedge X; Y\}_{\mathbb{Z}_2} = \{X; Y\}$ the forgetful map.

(iii) The special case $X = S^j \wedge LS^{i-j}$, $Y = S^0$, $V = \mathbb{R}(\infty)$ of the braid in (ii) is



(iv) Proposition 4.21 gives the bi-degree isomorphism

with

$$\begin{aligned}
\tilde{\omega}_i^{\mathbb{Z}_2}(X^{\mathbb{Z}_2}) &= \tilde{\omega}_i(X^{\mathbb{Z}_2}) \oplus \tilde{\omega}_i(P(\infty)^+ \wedge X^{\mathbb{Z}_2}) , \\
\tilde{\omega}_i^{\mathbb{Z}_2}(X) &= \tilde{\omega}_i(X^{\mathbb{Z}_2}) \oplus \tilde{\omega}_i(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) , \\
\tilde{\omega}_i^{\mathbb{Z}_2}(X/X^{\mathbb{Z}_2}) &= \tilde{\omega}_i((X/X^{\mathbb{Z}_2})/\mathbb{Z}_2) , \\
\tilde{\omega}_{\mathbb{Z}_2}^i(X^{\mathbb{Z}_2}) &= \tilde{\omega}^i(X^{\mathbb{Z}_2}) \oplus \tilde{\omega}^i(P(\infty)^+ \wedge X_{\mathbb{Z}_2}) , \\
\tilde{\omega}_{\mathbb{Z}_2}^i(X) &= \tilde{\omega}^i(X^{\mathbb{Z}_2}) \oplus \tilde{\omega}^i(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) , \\
\tilde{\omega}_{\mathbb{Z}_2}^i(X/X_{\mathbb{Z}_2}) &= \tilde{\omega}^i((X/X^{\mathbb{Z}_2})/\mathbb{Z}_2) .
\end{aligned}$$

Proof. These are \mathbb{Z}_2 -equivariant analogues of the Barratt-Puppe exact sequence. Proposition 4.41 (ii) (= Proposition 3.13 of Crabb [12]) gives the direct sum system

$$\tilde{\omega}_i^{\mathbb{Z}_2}(S(\infty)^+ \wedge X) \begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \tilde{\omega}_i^{\mathbb{Z}_2}(X) \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \tilde{\omega}_i(X^{\mathbb{Z}_2})$$

and Proposition 4.32 gives the identification

$$\tilde{\omega}_i^{\mathbb{Z}_2}(S(\infty)^+ \wedge X^{\mathbb{Z}_2}) = \tilde{\omega}_i(P(\infty)^+ \wedge X^{\mathbb{Z}_2}) .$$


Similarly for $\tilde{\omega}^i$.

□

Proposition 4.46. *For any space X and pointed \mathbb{Z}_2 -space Y there are defined direct sum systems*


$$\begin{aligned}
\{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} Y\} &\begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \{X; Y\}_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \{X; Y^{\mathbb{Z}_2}\} , \\
\{X; P(\infty)^+ \wedge Y^{\mathbb{Z}_2}\} &\begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \{X; Y^{\mathbb{Z}_2}\}_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \{X; Y^{\mathbb{Z}_2}\}
\end{aligned}$$

which fit together in a commutative braid of exact sequences



Proof. The direct sum systems are given by Proposition 4.41 (ii). \square

Example 4.47. In the special case $Y = A \wedge A$ for a pointed space A the commutative braid in Proposition 4.46 is given by



with $B = ((A \wedge A)/\Delta_A)/\mathbb{Z}_2$ and

$$\begin{aligned} \{X; A\}_{\mathbb{Z}_2} &= \{X; A\} \oplus \{X; P(\infty)^+ \wedge A\}, \\ \{X; A \wedge A\}_{\mathbb{Z}_2} &= \{X; A\} \oplus \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (A \wedge A)\}. \end{aligned}$$

\square

4.4 \mathbb{Z}_2 -equivariant bundles

We shall be mainly concerned with \mathbb{Z}_2 -equivariant U -bundles for finite-dimensional inner product spaces U .

Definition 4.48. (i) Let U be an inner product space, and X a \mathbb{Z}_2 -space. A \mathbb{Z}_2 -equivariant U -bundle ξ over X is a U -bundle

$$\xi : U \longrightarrow E(\xi) \xrightarrow{p} X$$

such that $E(\xi)$ is a \mathbb{Z}_2 -space, p is \mathbb{Z}_2 -equivariant, and the restrictions of $T : E(\xi) \rightarrow E(\xi)$ to the fibres

$$T|_{\xi_x} : \xi_x = p^{-1}(x) \rightarrow \xi_{Tx} = p^{-1}(Tx) = T\xi_x \quad (x \in X)$$

are isometries, with

$$\dim(\xi_x) = \dim(\xi_{T(x)}) = \dim(U)$$

and the Thom space $T(\xi)$ is a pointed \mathbb{Z}_2 -space. For $x \in X^{\mathbb{Z}_2}$ the fibre ξ_x is an inner product \mathbb{Z}_2 -space, and for finite-dimensional U the dimension of the fixed point subspace is a continuous function

$$\dim \xi_+ : X^{\mathbb{Z}_2} \rightarrow \mathbb{N} ; x \mapsto \dim(\xi_x)_+$$

which is constant on each component of $X^{\mathbb{Z}_2}$.

(ii) Let V be an inner product \mathbb{Z}_2 -space. The *trivial \mathbb{Z}_2 -equivariant $|V|$ -bundle* ϵ_V over a \mathbb{Z}_2 -space X is defined by

$$\epsilon_V : |V| \rightarrow E(\epsilon_V) = X \times V \rightarrow X ,$$

with

$$\begin{aligned} X \times V &\rightarrow X ; (x, v) \mapsto x , \quad (\epsilon_V)_x = V \quad (x \in X) , \\ T : E(\epsilon_V) &= X \times V \rightarrow X \times V ; (x, v) \mapsto (Tx, Tv) . \end{aligned}$$

(iii) The *Thom \mathbb{Z}_2 -spectrum* of a \mathbb{Z}_2 -equivariant U -bundle ξ is the suspension \mathbb{Z}_2 -spectrum of $T(\xi)$

$$\underline{T}(\xi) = \{T(\xi \oplus \epsilon_V) | V\}$$

with V running over finite-dimensional inner product \mathbb{Z}_2 -spaces, and

$$T(\xi \oplus \epsilon_V) = V^\infty \wedge T(\xi) .$$

For the trivial \mathbb{Z}_2 -equivariant V -bundle ϵ_V over X

$$\begin{aligned} E(\epsilon_V)^{\mathbb{Z}_2} &= V_+ \times X^{\mathbb{Z}_2}, \quad T(\epsilon_V) = V^\infty \wedge X^\infty, \\ T(\epsilon_V)^{\mathbb{Z}_2} &= (V_+)^{\infty} \wedge (X^{\mathbb{Z}_2})^\infty. \end{aligned}$$

□

Example 4.49. Let U be an inner product space.

(i) A \mathbb{Z}_2 -equivariant U -bundle ξ over $\{*\}$ is a \mathbb{Z}_2 -action on $E(\xi) = U$, so that

$$\xi = \epsilon_U = \epsilon_{U_+} \oplus \epsilon_{U_-}.$$

(ii) For a \mathbb{Z}_2 -equivariant U -bundle ξ over a \mathbb{Z}_2 -space X , forgetting the \mathbb{Z}_2 -structure determines a U -bundle ξ over X .

(iii) A U -bundle ξ over a space X can be regarded as \mathbb{Z}_2 -equivariant U -bundle ξ over X (with the trivial \mathbb{Z}_2 -action).

(iv) A U -bundle ξ over a space X determines a \mathbb{Z}_2 -equivariant U -bundle $L\xi$ over X with

$$E(L\xi) = E(\xi), \quad T : E(L\xi) \rightarrow E(L\xi); (x, u) \mapsto (x, -u),$$

such that

$$(L\xi)_+ = 0, \quad (L\xi)_- = \xi.$$

In particular, for the trivial U -bundle $\xi = \epsilon_U$ the construction gives the \mathbb{Z}_2 -equivariant U -bundle $L\xi = \epsilon_{LU}$.

□

Proposition 4.50. *Let ξ be a \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle over a \mathbb{Z}_2 -space X .*

(i) *Each fibre ξ_x ($x \in X^{\mathbb{Z}_2}$) is an inner product \mathbb{Z}_2 -space, and*

$$E(\xi)^{\mathbb{Z}_2} = \bigcup_{x \in X^{\mathbb{Z}_2}} (\xi_x)^{\mathbb{Z}_2}.$$

The subspaces

$$X(\xi, k) = \{x \in X^{\mathbb{Z}_2} \mid \dim((\xi_x)_+) = k\} \subseteq X^{\mathbb{Z}_2} \quad (0 \leq k \leq i)$$

are unions of components of $X^{\mathbb{Z}_2}$ such that

$$X^{\mathbb{Z}_2} = \prod_{k=0}^i X(\xi, k).$$

The restriction \mathbb{Z}_2 -equivariant bundle $(\xi, k) = \xi|_{X(\xi, k)}$ over $X(\xi, k)$ splits as a Whitney sum

$$(\xi, k) = (\xi, k)_+ \oplus L(\xi, k)_-$$

with

$$(\xi, k)_+ : X(\xi, k) \rightarrow BO(\mathbb{R}^k) , \quad (\xi, k)_- : X(\xi, k) \rightarrow BO(\mathbb{R}^{i-k})$$

nonequivariant bundles such that

$$\begin{aligned} E((\xi, k)_+) &= \bigcup_{x \in X(\xi, k)} (\xi_x)_+ , \quad E((\xi, k)_-) = \bigcup_{x \in X(\xi, k)} (\xi_x)_- , \\ E(\xi)^{\mathbb{Z}_2} &= \prod_{k=0}^i E((\xi, k)_+) , \quad T(\xi)^{\mathbb{Z}_2} = \bigvee_{k=0}^i T((\xi, k)_+) . \end{aligned}$$

(ii) For a trivial \mathbb{Z}_2 -equivariant bundle ϵ_V over a \mathbb{Z}_2 -space X

$$X(\epsilon_V, k) = \begin{cases} X^{\mathbb{Z}_2} & \text{if } k = \dim(V_+) \\ \emptyset & \text{if } k \neq \dim(V_+) . \end{cases}$$

(iii) For the Whitney sum $\xi \oplus \eta$ of \mathbb{Z}_2 -equivariant bundles ξ, η over a \mathbb{Z}_2 -space X

$$X(\xi \oplus \eta, k) = \coprod_{i+j=k} X(\xi, i) \cap X(\eta, j) .$$

In particular, if $\eta = \epsilon_V$

$$X(\xi \oplus \epsilon_V, k) = \begin{cases} X(\xi, k - \dim(V_+)) & \text{if } k \geq \dim(V_+) \\ \emptyset & \text{if } k < \dim(V_+) . \end{cases}$$

(iv) If $\xi \oplus \eta = \epsilon_V$ for some inner product \mathbb{Z}_2 -space V , then

$$(\xi_x)_+ \oplus (\eta_x)_+ = V_+ \quad (x \in X^{\mathbb{Z}_2}) ,$$

so that

$$\begin{aligned} X(\xi, k) &= \{x \in X^{\mathbb{Z}_2} \mid \dim((\xi_x)_+) = k\} \\ &= \{x \in X^{\mathbb{Z}_2} \mid \dim((\eta_x)_+) = \dim(V_+) - k\} \\ &= X(\eta, \dim(V_+) - k) . \end{aligned}$$

Proof. By construction. □

Proposition 4.51. Let U be a finite-dimensional inner product space, and let X be a \mathbb{Z}_2 -space, with $p : X \rightarrow X/\mathbb{Z}_2$ the projection.

- (i) For any U -bundle ξ over X/\mathbb{Z}_2 the pullback $p^*\xi$ is a \mathbb{Z}_2 -equivariant U -bundle over X , with the trivial \mathbb{Z}_2 -action on each fibre $p^*\xi_x = \xi_{p(x)}$ ($x \in X$).
- (ii) If X is a free \mathbb{Z}_2 -space the projection $p : X \rightarrow X/\mathbb{Z}_2$ is a double covering, and the function

$$\{U\text{-bundles over } X/\mathbb{Z}_2\} \rightarrow \{\mathbb{Z}_2\text{-equivariant } U\text{-bundles over } X\} ; \quad \xi \mapsto p^*\xi$$

is a bijection of the sets of isomorphism classes, with

$$\begin{aligned} E(p^*\xi)^{\mathbb{Z}_2} &= \emptyset, \quad T(p^*\xi)^{\mathbb{Z}_2} = \{\infty\}, \\ E(p^*\xi)/\mathbb{Z}_2 &= E(\xi), \quad T(p^*\xi)/\mathbb{Z}_2 = T(\xi). \end{aligned}$$

(iii) For any \mathbb{Z}_2 -equivariant U -bundle ξ over a \mathbb{Z}_2 -space X the function

$$\begin{aligned} &\{(\mathbb{Z}_2\text{-equivariant map } f : M \rightarrow X \text{ with } M \text{ } \mathbb{Z}_2\text{-free,} \\ &\quad \mathbb{Z}_2\text{-equivariant bundle map } b : \nu \rightarrow \xi)\} \\ &\rightarrow \{(g : M/\mathbb{Z}_2 \rightarrow S(\infty) \times_{\mathbb{Z}_2} X, \text{ bundle map } c : \nu/\mathbb{Z}_2 \rightarrow 0 \times_{\mathbb{Z}_2} \xi)\} \end{aligned}$$

is a bijection.

Proof. (i) By construction.

(ii) For any \mathbb{Z}_2 -equivariant U -bundle Ξ over X the \mathbb{Z}_2 -action on the total \mathbb{Z}_2 -space $E(\Xi)$ is free, and the quotient is the total space

$$E(\Xi)/\mathbb{Z}_2 = E(\xi)$$

of a U -bundle ξ over X/\mathbb{Z}_2 such that $\Xi = p^*\xi$.

□

Example 4.52. Let V be an inner product \mathbb{Z}_2 -space. The trivial \mathbb{Z}_2 -equivariant $|V|$ -bundle ϵ_V over a free \mathbb{Z}_2 -space X is the pullback $\epsilon_V = p^*(\epsilon_V/\mathbb{Z}_2)$ along the projection $p : X \rightarrow X/\mathbb{Z}_2$ of the $|V|$ -bundle ϵ_V/\mathbb{Z}_2 over X/\mathbb{Z}_2 with

$$V \rightarrow E(\epsilon_V/\mathbb{Z}_2) = E(\epsilon_V)/\mathbb{Z}_2 = X \times_{\mathbb{Z}_2} V \rightarrow X/\mathbb{Z}_2.$$

□

Proposition 4.53. (i) An m -dimensional \mathbb{Z}_2 -manifold M has a tangent \mathbb{Z}_2 -equivariant \mathbb{R}^m -bundle τ_M , namely the nonequivariant tangent \mathbb{R}^m -bundle τ_M with the \mathbb{Z}_2 -action $T : E(\tau_M) \rightarrow E(\tau_M)$ defined by the differentials of the \mathbb{Z}_2 -action $T : M \rightarrow M$.

(ii) The tangent \mathbb{R}^m -bundle of an m -dimensional free \mathbb{Z}_2 -manifold M is the pullback $\tau_M = p^*\tau_{M/\mathbb{Z}_2}$ along the projection $p : M \rightarrow M/\mathbb{Z}_2$ of τ_{M/\mathbb{Z}_2} , with

$$E(\tau_{M/\mathbb{Z}_2}) = E(\tau_M)/\mathbb{Z}_2, \quad \tau_{M/\mathbb{Z}_2} = \tau_M/\mathbb{Z}_2.$$

□

Example 4.54. Let U be a finite-dimensional inner product space.

(i) The \mathbb{Z}_2 -manifold LU has tangent \mathbb{Z}_2 -equivariant U -bundle

$$\tau_{LU} = \epsilon_{LU}.$$

(ii) The free \mathbb{Z}_2 -manifold $S(LU) \subset LU$ is such that there is defined a \mathbb{Z}_2 -equivariant homeomorphism

$$S(LU) \times \mathbb{R} \rightarrow LU \setminus \{0\} ; (u, x) \mapsto e^x u ,$$

so that the tangent \mathbb{Z}_2 -equivariant bundle $\tau_{S(LU)}$ is such that

$$\tau_{S(LU)} \oplus \epsilon_{\mathbb{R}} = \epsilon_{LU} .$$

(iii) The tangent bundle of the projective space $P(U) = S(LU)/\mathbb{Z}_2$

$$\tau_{P(U)} = \tau_{S(LU)}/\mathbb{Z}_2$$

As usual, it is possible to identify

$$\begin{aligned} P(U) &= G(\mathbb{R}, U) = \{K \subseteq U \mid \dim(K) = 1\} , \\ E(\tau_{S(LU)}) &= \{(x \in LU, y \in S(LU)) \mid \langle x, y \rangle = 1\} , \\ E(\tau_{P(U)}) &= E(\tau_{S(LU)})/\mathbb{Z}_2 = \{(K, x \in K^\perp) \mid K \subseteq U, \dim(K) = 1\} . \end{aligned}$$

□

Definition 4.55. Let U, V be finite-dimensional inner product spaces. The \mathbb{Z}_2 -equivariant embedding $S(LU) \subset S(LU \oplus LV)$ has normal \mathbb{Z}_2 -equivariant V -bundle

$$\nu_{S(LU) \subset S(LU \oplus LV)} = \epsilon_{LV} .$$

(i) The *Hopf V -bundle* over the projective space $P(U)$

$$H_V = \epsilon_{LV}/\mathbb{Z}_2 : V \rightarrow E(H_V) = S(LU) \times_{\mathbb{Z}_2} LV \rightarrow P(U)$$

is the normal V -bundle of the embedding $P(U) \subset P(U \oplus V)$

$$H_V = \nu_{P(U) \subset P(U \oplus V)} : P(U) \rightarrow BO(V) .$$

Also, for $V = U$

$$H_U = \tau_{P(U)} \oplus \epsilon_{\mathbb{R}} = \tau_{P(U \oplus \mathbb{R})} \mid : P(U) \rightarrow BO(U) .$$

(ii) The Thom space of H_V is the *stunted projective space*

$$T(H_V) = S(LU)^+ \wedge_{\mathbb{Z}_2} LV^\infty .$$

For $U = \mathbb{R}^m, V = \mathbb{R}^n$ write

$$P(\mathbb{R}^{m+n})/P(\mathbb{R}^n) = P(m+n, n) = S(L\mathbb{R}^m)^+ \wedge_{\mathbb{Z}_2} L\mathbb{R}^n ,$$

with $P(m, 0) = P(\mathbb{R}^m)^+$.

(iii) The *infinite stunted projective space* $P(\infty, n)$ is defined for any $n \in \mathbb{Z}$

to be the virtual Thom space (3.29) of the virtual bundle $\text{sgn}(n) H_{\mathbb{R}|n|}$ over $P(\mathbb{R}^m)$ for sufficiently large $m \geq 0$. For $n \geq 0$ this is the actual Thom space

$$P(\infty, n) = T(H_{\mathbb{R}^n}) = P(\mathbb{R}(\infty) \oplus \mathbb{R}^n)/P(\mathbb{R}^n) = S(\infty)^+ \wedge_{\mathbb{Z}_2} LS^n \\ (= P(\infty)^+ \text{ for } n = 0) ,$$

so that

$$\tilde{\omega}_j(P(\infty, n)) = \{S^j; S(\infty)^+ \wedge LS^n\}_{\mathbb{Z}_2} .$$

For $n < 0$ this is the ‘space’ $P(\infty, n)$ with

$$\tilde{\omega}_j(P(\infty, n)) = \{S^j \wedge LS^{-n}; S(\infty)^+\}_{\mathbb{Z}_2} .$$

□

Example 4.56. The Hopf \mathbb{R} -bundle $H_{\mathbb{R}}$ over $P(U)$ is such that

$$E(H_{\mathbb{R}}) = \{(K, x \in K) \mid K \subseteq U, \dim(K) = 1\} = S(LU) \times_{\mathbb{Z}_2} L\mathbb{R} , \\ S(H_{\mathbb{R}}) = \{(K, x \in K) \mid K \subseteq U, \dim(K) = 1, \|x\| = 1\} = S(LU) , \\ H_{\mathbb{R}} = \nu_{P(U) \subset P(U \oplus \mathbb{R})} : P(U) \rightarrow \varinjlim_p P(U \oplus \mathbb{R}^p) = BO(\mathbb{R}) , \\ \tau_{P(U)} \oplus \epsilon_{\mathbb{R}} = \tau_{P(U \oplus \mathbb{R})}| = H_U : P(U) \rightarrow BO(U) , \\ T(H_{\mathbb{R}}) = P(U \oplus \mathbb{R})/P(\mathbb{R}) = S(LU)^+ \wedge_{\mathbb{Z}_2} L\mathbb{R}^\infty ,$$

and for any $n \geq 1$

$$\nu_{P(U) \subset P(U \oplus \mathbb{R}^n)} = H_{\mathbb{R}^n} = \bigoplus_n H_{\mathbb{R}} : P(U) \rightarrow BO(\mathbb{R}^n) , \\ \tau_{P(U)} \oplus \epsilon_{\mathbb{R}} \oplus H_{\mathbb{R}^{n-1}} = \tau_{P(U \oplus \mathbb{R}^n)}| = H_{U \oplus \mathbb{R}^{n-1}} : P(U) \rightarrow BO(U \oplus \mathbb{R}^{n-1}) .$$

□

Definition 4.57. Let $\xi - \xi'$ be a virtual \mathbb{Z}_2 -equivariant bundle over a \mathbb{Z}_2 -space X , and let η be a \mathbb{Z}_2 -equivariant bundle over X such that $\xi' \oplus \eta = \epsilon_V$ for some finite-dimensional inner product \mathbb{Z}_2 -space V , so that $\xi - \xi' = \xi \oplus \eta - \epsilon_V$.

(i) The *virtual Thom \mathbb{Z}_2 -space* of $\xi - \xi'$ is the \mathbb{Z}_2 -spectrum

$$\underline{T}(\xi - \xi') = \{T(\xi - \xi')_U \mid U\}$$

defined by

$$T(\xi - \xi')_U = T(\xi \oplus \eta \oplus \epsilon_{U/V})$$

for finite-dimensional inner product \mathbb{Z}_2 -spaces U such that $V \subseteq U$.

(ii) Write

$$\omega_{\mathbb{Z}_2}^*(X; \xi - \xi') = \tilde{\omega}_{\mathbb{Z}_2}^*(\underline{T}(\xi - \xi')) , \\ \omega_*^{\mathbb{Z}_2}(X; \xi - \xi') = \tilde{\omega}_*^{\mathbb{Z}_2}(\underline{T}(\xi - \xi'))$$

so that

$$\begin{aligned}\omega_{\mathbb{Z}_2}^m(X; \xi - \xi') &= \{T(\xi \oplus \eta); \Sigma^m V^\infty\}_{\mathbb{Z}_2} , \\ \omega_m^{\mathbb{Z}_2}(X; \xi - \xi') &= \{\Sigma^m V^\infty; T(\xi \oplus \eta)\}_{\mathbb{Z}_2} .\end{aligned}$$

□

Proposition 4.58. (i) *Let X be a pointed \mathbb{Z}_2 -space and let $p : X \rightarrow X/\mathbb{Z}_2$ be the projection, so that there are induced morphisms*

$$\begin{aligned}p^* : \tilde{\omega}^m(X/\mathbb{Z}_2) &\rightarrow \tilde{\omega}_{\mathbb{Z}_2}^m(X) ; \\ (F : V^\infty \wedge X/\mathbb{Z}_2 &\rightarrow \Sigma^m V^\infty) \mapsto (F(1 \wedge p) : V^\infty \wedge X \rightarrow \Sigma^m V^\infty) , \\ p_* : \tilde{\omega}_m^{\mathbb{Z}_2}(X) &\rightarrow \tilde{\omega}_m(X/\mathbb{Z}_2) ; \\ (F : \Sigma^m V^\infty &\rightarrow V^\infty \wedge X) \mapsto ((1 \wedge p)F : \Sigma^m V^\infty \rightarrow V^\infty \wedge X/\mathbb{Z}_2) .\end{aligned}$$

If X is semifree then p^ is an isomorphism, and if in addition X is a finite CW \mathbb{Z}_2 -complex then p_* is also an isomorphism.*

(ii) *For a virtual \mathbb{Z}_2 -equivariant bundle $\xi - \xi'$ over a free \mathbb{Z}_2 -space X*

$$\omega_{\mathbb{Z}_2}^*(X; \xi - \xi') = \omega^*(X/\mathbb{Z}_2; \xi/\mathbb{Z}_2 - \xi'/\mathbb{Z}_2)$$

and if in addition X is a finite CW \mathbb{Z}_2 -complex then

$$\omega_*^{\mathbb{Z}_2}(X; \xi - \xi') = \omega_*(X/\mathbb{Z}_2; \xi/\mathbb{Z}_2 - \xi'/\mathbb{Z}_2) .$$

Proof. (i) For any finite-dimensional inner product \mathbb{Z}_2 -space V there is defined a V -bundle ϵ_V/\mathbb{Z}_2 over X/\mathbb{Z}_2 with

$$E(\epsilon_V/\mathbb{Z}_2) = V \times_{\mathbb{Z}_2} X .$$

For sufficiently large $n \geq 0$ there exists a bundle λ_V over X/\mathbb{Z}_2 such that

$$(\epsilon_V/\mathbb{Z}_2) \oplus \lambda_V = \epsilon_{\mathbb{R}^n} .$$

An element $G \in \omega_{\mathbb{Z}_2}^m(X)$ is represented by a \mathbb{Z}_2 -equivariant map $G : V^\infty \wedge X \rightarrow \Sigma^m V^\infty$, and the composite

$$\begin{aligned}H : (V^\infty \wedge_{\mathbb{Z}_2} X) \wedge_{X/\mathbb{Z}_2} T(\lambda_V) &= \Sigma^n X^+ \xrightarrow{G \wedge 1} \\ \Sigma^m (V^\infty \wedge_{\mathbb{Z}_2} X) \wedge_{X/\mathbb{Z}_2} T(\lambda_V) &= \Sigma^{m+n} X \rightarrow S^{m+n}\end{aligned}$$

is a stable map such that

$$\omega_{\mathbb{Z}_2}^m(X) \rightarrow \omega^m(X/\mathbb{Z}_2) ; G \mapsto H$$

is an inverse to p^* .

For X a finite CW \mathbb{Z}_2 -complex the stable \mathbb{Z}_2 -equivariant Umkehr map $F \in$

$\{X/\mathbb{Z}_2; X\}_{\mathbb{Z}_2}$ of Proposition 4.32 (ii) induces a morphism

$$\omega_m(X/\mathbb{Z}_2) \rightarrow \omega_m^{\mathbb{Z}_2}(X) ; G \mapsto FG$$

inverse to p_* , by 4.32 (iii).

(ii) Let η be a bundle over X such that $\xi'/\mathbb{Z}_2 \oplus \eta = \epsilon_V$ for some finite-dimensional inner product space V , so that $\xi' \oplus p^*\eta = \epsilon_V$,

$$\begin{aligned} \omega_{\mathbb{Z}_2}^m(X; \xi - \xi') &= \{T(\xi \oplus p^*\eta); \Sigma^m V^\infty\}_{\mathbb{Z}_2} , \\ \omega^m(X/\mathbb{Z}_2; \xi/\mathbb{Z}_2 - \xi'/\mathbb{Z}_2) &= \{T(\xi/\mathbb{Z}_2 \oplus \eta); \Sigma^m V^\infty\} , \\ \omega_m^{\mathbb{Z}_2}(X; \xi - \xi') &= \{\Sigma^m V^\infty; T(\xi \oplus \eta)\}_{\mathbb{Z}_2} , \\ \omega_m(X/\mathbb{Z}_2; \xi/\mathbb{Z}_2 - \xi'/\mathbb{Z}_2) &= \{\Sigma^m V^\infty; T(\xi/\mathbb{Z}_2 \oplus \eta)\} . \end{aligned}$$

The \mathbb{Z}_2 -action on $T(\xi \oplus p^*\eta)$ is semifree, with quotient

$$T(\xi \oplus p^*\eta)/\mathbb{Z}_2 = T(\xi/\mathbb{Z}_2 \oplus \eta) ,$$

so that by (i)

$$\begin{aligned} \omega_{\mathbb{Z}_2}^m(X; \xi - \xi') &= \{T(\xi \oplus p^*\eta); \Sigma^m V^\infty\}_{\mathbb{Z}_2} \\ &= \{T(\xi/\mathbb{Z}_2 \oplus \eta); \Sigma^m V^\infty\} = \omega^m(X/\mathbb{Z}_2; \xi/\mathbb{Z}_2 - \xi'/\mathbb{Z}_2) , \\ \omega_m^{\mathbb{Z}_2}(X; \xi - \xi') &= \{\Sigma^m V^\infty; T(\xi \oplus p^*\eta)\}_{\mathbb{Z}_2} \\ &= \{\Sigma^m V^\infty; T(\xi/\mathbb{Z}_2 \oplus \eta)\} = \omega_m(X/\mathbb{Z}_2; \xi/\mathbb{Z}_2 - \xi'/\mathbb{Z}_2) . \end{aligned}$$

□

Terminology 4.59 For $i \geq j \geq 0$ the i -dimensional inner product \mathbb{Z}_2 -space $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ has a j -dimensional fixed point space $V_+ = \mathbb{R}^j$. For any \mathbb{Z}_2 -space X write

$$\begin{aligned} \omega^{i,j}(X) &= \omega_{\mathbb{Z}_2}^0(X; -\epsilon_V) = \{X^+; V^\infty\}_{\mathbb{Z}_2} , \\ \omega_{i,j}(X) &= \omega_0^{\mathbb{Z}_2}(X; -\epsilon_V) = \{V^\infty; X^+\}_{\mathbb{Z}_2} \end{aligned}$$

exactly as in Crabb [12, pp. 28,29].

□

Example 4.60. If X is a free \mathbb{Z}_2 -space and $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ then

$$\begin{aligned} \omega^{i,j}(X) &= \omega_{\mathbb{Z}_2}^0(X; -\epsilon_V) = \omega^0(X/\mathbb{Z}_2; -\epsilon_V/2) , \\ \omega_{i,j}(X) &= \omega_0^{\mathbb{Z}_2}(X; -\epsilon_V) = \omega_0(X/\mathbb{Z}_2; -\epsilon_V/2) \end{aligned}$$

by Proposition 4.58, assuming X is a finite CW \mathbb{Z}_2 -complex for $\omega_{i,j}(X)$.

□

Example 4.61. The \mathbb{Z}_2 -equivariant cofibration sequence

$$S(\infty)^+ \rightarrow S^0 \rightarrow L\mathbb{R}(\infty)^\infty \rightarrow \Sigma S(\infty)^+ \rightarrow \dots$$

induces the exact sequence

$$\dots \longrightarrow \tilde{\omega}_j(P(\infty, j-i)) \longrightarrow \omega_{i,j} \xrightarrow{\rho} \omega_j \longrightarrow \tilde{\omega}_{j-1}(P(\infty, j-i)) \longrightarrow \dots$$

of Crabb [12, Proposition 4.6], with $\omega_{i,j} = \omega_{i,j}(\{*\})$.

□

Example 4.62. (i) For $i = j$

$$\begin{aligned} \omega^{i,i}(X) &= \omega_{\mathbb{Z}_2}^i(X) = \{X^+; S^i\}_{\mathbb{Z}_2} , \\ \omega_{i,i}(X) &= \omega_i^{\mathbb{Z}_2}(X) = \{S^i; X^+\}_{\mathbb{Z}_2} . \end{aligned}$$

(ii) For $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$, $V' = \mathbb{R}^{j'} \oplus L\mathbb{R}^{i'-j'}$,

$$\begin{aligned} \omega_{\mathbb{Z}_2}^n(X; \epsilon_V - \epsilon_{V'}) &= \{V^\infty \wedge X^+; (V' \oplus \mathbb{R}^n)^\infty\}_{\mathbb{Z}_2} = \omega^{n+i'-i, n+j'-j}(X) , \\ \omega_n^{\mathbb{Z}_2}(X; \epsilon_V - \epsilon_{V'}) &= \{(V' \oplus \mathbb{R}^n)^\infty; V^\infty \wedge X^+\}_{\mathbb{Z}_2} = \omega_{n+i'-i, n+j'-j}(X) . \end{aligned}$$

□

Proposition 4.63. *Let ξ be a \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle over a \mathbb{Z}_2 -space X , so that $0 \times \xi$ is a \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle over the free \mathbb{Z}_2 -space $S(\infty) \times X$ with*

$$E(0 \times \xi) = S(\infty) \times E(\xi) , \quad T(0 \times \xi) = S(\infty)^+ \wedge T(\xi) ,$$

and $0 \times_{\mathbb{Z}_2} \xi$ is an \mathbb{R}^i -bundle over $S(\infty) \times_{\mathbb{Z}_2} X$ with

$$E(0 \times_{\mathbb{Z}_2} \xi) = S(\infty) \times_{\mathbb{Z}_2} E(\xi) , \quad T(0 \times_{\mathbb{Z}_2} \xi) = S(\infty)^+ \wedge_{\mathbb{Z}_2} T(\xi) .$$

(i) *For any inner product \mathbb{Z}_2 -space V the stable \mathbb{Z}_2 -equivariant homotopy groups $\omega_*^{\mathbb{Z}_2}(X; \xi - \epsilon_V)$ fit into a long exact sequence*

$$\begin{aligned} \dots \rightarrow \omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) &\rightarrow \omega_m^{\mathbb{Z}_2}(X; \xi - \epsilon_V) \\ \xrightarrow{\rho} \omega_m(T(\xi - \epsilon_V)^{\mathbb{Z}_2}) &\rightarrow \omega_{m-1}^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \rightarrow \dots \end{aligned}$$

with ρ the fixed point map, and

$$\begin{aligned} \omega_m^{\mathbb{Z}_2}(X; \xi - \epsilon_V) &= \{(V \oplus \mathbb{R}^m)^\infty; T(\xi)\}_{\mathbb{Z}_2} , \\ \omega_m(T(\xi - \epsilon_V)^{\mathbb{Z}_2}) &= \{(V_+ \oplus \mathbb{R}^m)^\infty; T(\xi)^{\mathbb{Z}_2}\} \\ &= \bigoplus_{k=0}^i \omega_{m+\dim(V_+)}(X(\xi, k); (\xi, k)_+) . \end{aligned}$$

Furthermore, if X is a finite CW \mathbb{Z}_2 -complex

$$\omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) = \omega_m(S(\infty) \times_{\mathbb{Z}_2} X; 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) .$$

(ii) For $V = \{0\}$ the sequence in (i) breaks up into split short exact sequences: the stable \mathbb{Z}_2 -equivariant homotopy groups $\omega_*^{\mathbb{Z}_2}(X; \xi)$ split as

$$\begin{aligned} \omega_m^{\mathbb{Z}_2}(X; \xi) &= \tilde{\omega}_m^{\mathbb{Z}_2}(T(\xi)) \\ &= \tilde{\omega}_m(S(\infty)^+ \wedge_{\mathbb{Z}_2} T(\xi)) \oplus \tilde{\omega}_m(T(\xi)^{\mathbb{Z}_2}) \\ &= \omega_m(S(\infty) \times_{\mathbb{Z}_2} X; 0 \times_{\mathbb{Z}_2} \xi) \oplus \bigoplus_{k=0}^i \omega_m(X(\xi, k); (\xi, k)_+) . \end{aligned}$$

Proof. These are special cases of Proposition 4.30 (ii)+(iii), combined with Proposition 4.58. \square

The nonequivariant bundle constructions of §3.2 all have \mathbb{Z}_2 -equivariant analogues, including classifying spaces. We shall only be considering \mathbb{Z}_2 -equivariant bundles over reasonable \mathbb{Z}_2 -spaces, such as finite CW \mathbb{Z}_2 -complexes.

For an inner product \mathbb{Z}_2 -space V a subspace $W \subseteq |V|$ is \mathbb{Z}_2 -invariant (i.e. $TW = W$) if and only if

$$W = W_+ \oplus W_- \subseteq |V| = V_+ \oplus V_-$$

with

$$W_+ = W \cap V_+ \subseteq V_+ , \quad W_- = W \cap V_- \subseteq V_- .$$

Definition 4.64. (i) Given a finite-dimensional inner product space U and a finite-dimensional inner product \mathbb{Z}_2 -space V with $\dim(U) \leq \dim(V)$ let $G(U, V)$ be the Grassmann \mathbb{Z}_2 -manifold of subspaces $W \subseteq |V|$ isomorphic to U , with the \mathbb{Z}_2 -action

$$T : G(U, V) \rightarrow G(U, V) ; W \mapsto TW .$$

Nonequivariantly $G(U, V) = G(U, |V|)$. The fixed point space is

$$\begin{aligned} G(U, V)^{\mathbb{Z}_2} &= \{W \mid W = TW \in G(U, V)\} \\ &= \coprod_{\substack{k=\max(0, \dim(U)-\dim(V_-)) \\ \min(\dim(U), \dim(V_+))}}^{\min(\dim(U), \dim(V_+))} \{W \mid W = TW \in G(U, V), \dim(W_+) = k\} , \\ &= \coprod_{k=\max(0, \dim(U)-\dim(V_-))}^{\min(\dim(U), \dim(V_+))} G(\mathbb{R}^k, V_+) \times G(\mathbb{R}^{\dim(U)-k}, V_-) . \end{aligned}$$

(ii) The *canonical \mathbb{Z}_2 -equivariant U -bundle* over $G(U, V)$

$$\xi(U, V) : U \longrightarrow E(\xi(U, V)) \longrightarrow G(U, V)$$

has

$$\begin{aligned} E(\xi(U, V)) &= O(U, |V|) \times_{O(U)} U \\ &= \{(W, x) \mid W \in G(U, V), x \in W\} , \\ T : E(\xi(U, V)) &\rightarrow E(\xi(U, V)) ; (W, x) \mapsto (TW, Tx) , \end{aligned}$$

with

$$\begin{aligned} G(U, V)(\xi(U, V), k) &= G(\mathbb{R}^k, V_+) \times G(\mathbb{R}^{\dim(U)-k}, V_-) , \\ E(\xi(U, V))^{\mathbb{Z}_2} &= \{(W, x) \mid W = TW \in G(U, V), x \in W^{\mathbb{Z}_2}\} \\ &= \coprod_{k=\max(0, \dim(U)-\dim(V_-))}^{\min(\dim(U), \dim(V_+))} E(\xi(\mathbb{R}^k, V_+)) . \end{aligned}$$

The Thom \mathbb{Z}_2 -space

$$T(\xi(U, V)) = O(U, |V|)^+ \wedge_{O(U)} U^\infty$$

is written $MO^{\mathbb{Z}_2}(U, V)$, and is such that

$$MO^{\mathbb{Z}_2}(U, V)^{\mathbb{Z}_2} = \bigvee_{k=\max(0, \dim(U)-\dim(V_-))}^{\min(\dim(U), \dim(V_+))} T(\xi(\mathbb{R}^k, V_+)) .$$

(iii) For a finite-dimensional inner product \mathbb{Z}_2 -space V the *classifying space* for \mathbb{Z}_2 -equivariant $|V|$ -bundles is the \mathbb{Z}_2 -space

$$BO^{\mathbb{Z}_2}(V) = \varinjlim_{p,q} G(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q) .$$

Note that nonequivariantly $BO^{\mathbb{Z}_2}(V) = BO(|V|)$. The *canonical \mathbb{Z}_2 -equivariant $|V|$ -bundle* over $BO^{\mathbb{Z}_2}(V)$ is

$$\begin{aligned} \xi(V) &= \varinjlim_{p,q} \xi(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q) : \\ |V| \rightarrow EO^{\mathbb{Z}_2}(V) &= \varinjlim_{p,q} E(\xi(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q)) \rightarrow BO^{\mathbb{Z}_2}(V) , \end{aligned}$$

with Thom \mathbb{Z}_2 -space

$$T(\xi(V)) = MO^{\mathbb{Z}_2}(V) = \varinjlim_{p,q} MO^{\mathbb{Z}_2}(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q) .$$

(iv) The *universal Thom \mathbb{Z}_2 -spectrum* is

$$\underline{MO}^{\mathbb{Z}_2} = \{MO^{\mathbb{Z}_2}(V) \mid V\} .$$

□

Proposition 4.65. *Let V be a finite-dimensional inner product \mathbb{Z}_2 -space.*

(i) *The \mathbb{Z}_2 -equivariant $|V|$ -bundles over X*

$$\xi : |V| \longrightarrow E(\xi) \longrightarrow X$$

are classified by the \mathbb{Z}_2 -equivariant homotopy classes of \mathbb{Z}_2 -equivariant maps

$$f : X \rightarrow G(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q) \subset BO^{\mathbb{Z}_2}(V) \quad (p, q \text{ large}) ,$$

with

$$\xi = f^*\xi(V) , \quad E(\xi) = f^*E(\xi(V)) = \{(x, y) \in X \times V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q \mid y \in f(x)\} .$$

The function

$$\begin{aligned} [X, BO^{\mathbb{Z}_2}(V)]_{\mathbb{Z}_2} &\rightarrow \\ \{ \text{isomorphism classes of } \mathbb{Z}_2\text{-equivariant } |V|\text{-bundles over } X \} &; \\ (f : X \rightarrow BO^{\mathbb{Z}_2}(V)) &\mapsto f^*\xi(V) \end{aligned}$$

is a bijection.

(ii) *For any \mathbb{Z}_2 -equivariant $|V|$ -bundle $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$ there exist a finite-dimensional inner product \mathbb{Z}_2 -space W and a \mathbb{Z}_2 -equivariant $|W|$ -bundle $\eta : X \rightarrow BO^{\mathbb{Z}_2}(W)$ such that*

$$\xi \oplus \eta = \epsilon_{V \oplus W} : X \rightarrow BO^{\mathbb{Z}_2}(W) .$$

(iii) *The forgetful function*

$$\{\mathbb{Z}_2\text{-equivariant } |V|\text{-bundles over } X\} \rightarrow \{|V|\text{-bundles over } X\}$$

corresponds to the homeomorphism

$$BO^{\mathbb{Z}_2}(V) = \varinjlim_{p,q} G(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q) \rightarrow BO(|V|) = \varinjlim_{p,q} G(|V|, |V \oplus \mathbb{R}^{p+q}|)$$

which forgets the \mathbb{Z}_2 -action.

(iv) *If X is a free CW \mathbb{Z}_2 -complex and $p : X \rightarrow X/\mathbb{Z}_2$ is the projection, the function*

$$[X/\mathbb{Z}_2, BO(|V|)] \rightarrow [X, BO^{\mathbb{Z}_2}(V)]_{\mathbb{Z}_2} ; \quad \xi \mapsto p^*\xi$$

is a bijection, so that

$$\{\mathbb{Z}_2\text{-equivariant } |V|\text{-bundles over } X\} = \{|V|\text{-bundles over } X/\mathbb{Z}_2\} .$$

(v) For $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ and any $p, q \geq 0$ the fixed point set of $G(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q)$ is

$$G(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q)^{\mathbb{Z}_2} = \prod_{k=\max(0, j-q)}^{\min(i, j+p)} G(\mathbb{R}^k, \mathbb{R}^{j+p}) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j+q}) .$$

The canonical \mathbb{Z}_2 -equivariant $|V|$ -bundle $\xi(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q)$ restricts to

$$\begin{aligned} \xi(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q) &= \prod_{k=\max(0, j-q)}^{\min(i, j+p)} \xi(\mathbb{R}^k, \mathbb{R}^{j+p}) \oplus L\xi(\mathbb{R}^{i-k}, \mathbb{R}^{i-j+q}) : \\ &\prod_{k=\max(0, j-q)}^{\min(i, j+p)} G(\mathbb{R}^k, \mathbb{R}^{j+p}) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j+q}) \rightarrow BO^{\mathbb{Z}_2}(V) \end{aligned}$$

so that

$$\begin{aligned} T(\xi(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q))^{\mathbb{Z}_2} &= T(\xi(|V|, V \oplus \mathbb{R}^p \oplus L\mathbb{R}^q))^{\mathbb{Z}_2} \\ &= \bigvee_{k=\max(0, j-q)}^{\min(i, j+p)} T(\xi(\mathbb{R}^k, \mathbb{R}^{j+p})) \wedge G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j+q})^+ , \\ \xi(V) &= \coprod_{k=0}^i (\xi(V, k)_+ \oplus \xi(V, k)_-) : \\ BO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2} &= \varinjlim_{p, q} \left(\prod_{k=\max(0, j-q)}^{\min(i, j+p)} G(\mathbb{R}^k, \mathbb{R}^{j+p}) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j+q}) \right) \\ &= \prod_{k=0}^i BO(\mathbb{R}^k) \times BO(\mathbb{R}^{i-k}) \rightarrow BO^{\mathbb{Z}_2}(V) , \\ BO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2}(\xi(V)) &= \varinjlim_{p, q} \left(\prod_{k=\max(0, j-q)}^{\min(i, j+p)} G(\mathbb{R}^k, \mathbb{R}^{j+p}) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j+q}) \right) , \\ BO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2}(\xi(V), k) &= BO(\mathbb{R}^k) \times BO(\mathbb{R}^{i-k}) , \\ T(\xi(V))^{\mathbb{Z}_2} &= MO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2} = \bigvee_{k=0}^i MO(k) \wedge BO(\mathbb{R}^{i-k})^+ . \end{aligned}$$

(vi) If the \mathbb{Z}_2 -action on X is trivial then

$$[X, BO^{\mathbb{Z}_2}(V)]_{\mathbb{Z}_2} = [X, BO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2}] = [X, \prod_{k=0}^i BO(\mathbb{R}^k) \times BO(\mathbb{R}^{i-k})] ,$$

and for any $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$

$$X(\xi, k) = \xi^{-1}(BO(\mathbb{R}^k) \times BO(\mathbb{R}^{i-k})) \subseteq X .$$

(vii) The two inclusions

$$\begin{aligned}\iota_1 & : BO(|V|) = BO(|V|) \times \{*\} \subset BO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2} , \\ \iota_2 & : BO(|V|) = \{*\} \times BO(|V|) \subset BO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2}\end{aligned}$$

correspond to the two constructions of a \mathbb{Z}_2 -equivariant $|V|$ -bundle over X from a nonequivariant $|V|$ -bundle $\xi : X \rightarrow BO(|V|)$, with $\iota_1\xi = \xi$, $\iota_2\xi = L\xi$. \square

In particular, for a \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$ ($i = \dim(V)$) over a \mathbb{Z}_2 -space X

(i) the \mathbb{Z}_2 -action on $Y = X \setminus X^{\mathbb{Z}_2}$ is free, so that

$$\xi|_Y = p^*(\xi|_{Y/\mathbb{Z}_2}) : Y \rightarrow BO^{\mathbb{Z}_2}(V)$$

is the \mathbb{Z}_2 -equivariant pullback along the double cover projection $p : Y \rightarrow Y/\mathbb{Z}_2$ of a nonequivariant \mathbb{R}^i -bundle $\xi|_{Y/\mathbb{Z}_2} : Y/\mathbb{Z}_2 \rightarrow BO(V)$,

(ii) the fixed point space decomposes as

$$X^{\mathbb{Z}_2} = \coprod_{k=0}^i X(\xi, k) ,$$

with each $X(\xi, k)$ a union of components,

$$(\xi, k) = (\xi, k)_+ \oplus L(\xi, k)_- : X(\xi, k) \rightarrow BO(\mathbb{R}^k) \times BO(\mathbb{R}^{i-k}) \subset BO^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2} ,$$

and

$$T(\xi)^{\mathbb{Z}_2} = \bigvee_{k=0}^i T((\xi, k)_+) .$$

Proposition 4.66. (i) The universal Thom \mathbb{Z}_2 -spectrum $\underline{MO}^{\mathbb{Z}_2}$ is split, and such that

$$|\underline{MO}^{\mathbb{Z}_2}| = \underline{MO} ,$$

the universal Thom spectrum.

(ii) The homotopy orbit \mathbb{Z}_2 -spectrum of $\underline{MO}^{\mathbb{Z}_2}$ is such that

$$(\underline{MO}^{\mathbb{Z}_2})_{h\mathbb{Z}_2} \simeq S(\infty)^+ \wedge \underline{MO} .$$

(iii) The homotopy fixed point \mathbb{Z}_2 -spectrum of $\underline{MO}^{\mathbb{Z}_2}$ is such that the map

$$S(L\mathbb{R})^+ \wedge (\underline{MO}^{\mathbb{Z}_2})^{\mathbb{Z}_2} \rightarrow (\underline{MO}^{\mathbb{Z}_2})^{h\mathbb{Z}_2}$$

induces isomorphisms of reduced generalized homology groups for any pointed \mathbb{Z}_2 -space X

$$\begin{aligned}
(S(L\mathbb{R})^+ \wedge (\underline{MO}^{\mathbb{Z}_2})^{\mathbb{Z}_2})_*(X) &= \bigoplus_{k=0}^{\infty} \widetilde{MO}_k(X^{\mathbb{Z}_2} \wedge BO^+) \\
&\cong (\widetilde{MO}^{\mathbb{Z}_2})_*^{h\mathbb{Z}_2}(X)
\end{aligned}$$

with an exact sequence

$$\begin{aligned}
\cdots \rightarrow \widetilde{MO}_*(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) &\rightarrow \widetilde{MO}_*^{\mathbb{Z}_2}(X) \\
&\rightarrow \bigoplus_{k=0}^{\infty} \widetilde{MO}_k(X^{\mathbb{Z}_2} \wedge BO^+) \rightarrow \widetilde{MO}_{*-1}(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) \rightarrow \cdots
\end{aligned}$$

Proof. (i) For each inner product space U the canonical \mathbb{Z}_2 -equivariant U -bundle $\xi(U)$ over $BO^{\mathbb{Z}_2}(U)$ is classified as a nonequivariant U -bundle by a map

$$\zeta(U) : BO^{\mathbb{Z}_2}(U) \rightarrow BO(|U|) \subset BO^{\mathbb{Z}_2}(U)$$

with a corresponding map of Thom spaces

$$\zeta(U) : |\underline{MO}^{\mathbb{Z}_2}(U)| = MO^{\mathbb{Z}_2}(U) = MO(U) \rightarrow (\underline{MO}^{\mathbb{Z}_2})^{\mathbb{Z}_2}(U) = MO^{\mathbb{Z}_2}(U)^{\mathbb{Z}_2}.$$

The corresponding map of spectra

$$\zeta : |\underline{MO}^{\mathbb{Z}_2}| \rightarrow (\underline{MO}^{\mathbb{Z}_2})^{\mathbb{Z}_2}$$

is such that $\zeta j \simeq 1$, with $j : (\underline{MO}^{\mathbb{Z}_2})^{\mathbb{Z}_2} \rightarrow |\underline{MO}^{\mathbb{Z}_2}|$ the inclusion.

(ii)+(iii) Immediate from (i) and Proposition 4.38. □

Proposition 4.67. (Conner and Floyd [8, 22.1]) (The \mathbb{Z}_2 -equivariant tubular neighbourhood theorem).

A \mathbb{Z}_2 -equivariant embedding $M \subset N$ of a closed m -dimensional \mathbb{Z}_2 -manifold M^m in a closed n -dimensional \mathbb{Z}_2 -manifold N^n has a normal \mathbb{Z}_2 -equivariant \mathbb{R}^{n-m} -bundle

$$\nu_{M \subset N} : M \rightarrow BO^{\mathbb{Z}_2}(\mathbb{R}^{n-m})$$

such that

$$\tau_M \oplus \nu_{M \subset N} = \tau_N|_M : M \rightarrow BO^{\mathbb{Z}_2}(\mathbb{R}^n),$$

with a codimension 0 \mathbb{Z}_2 -equivariant embedding $E(\nu_{M \subset N}) \subset N$ and a \mathbb{Z}_2 -equivariant Umkehr map

$$\alpha : N^\infty \rightarrow N/(N \setminus E(\nu_{M \subset N})) = T(\nu_{M \subset N}).$$

□

Proposition 4.68. Let M be an m -dimensional \mathbb{Z}_2 -manifold.

(i) There exists a \mathbb{Z}_2 -equivariant embedding $M \subset V \oplus \mathbb{R}^m$ for some finite-

dimensional inner product \mathbb{Z}_2 -space V .

(ii) The embedding in (i) extends to a \mathbb{Z}_2 -equivariant embedding $E(\nu_M) \subset V \oplus \mathbb{R}^m$ of the normal \mathbb{Z}_2 -equivariant $|V|$ -bundle

$$\nu_M = \nu_{M \subset V \oplus \mathbb{R}^m} : M \rightarrow BO^{\mathbb{Z}_2}(V) ,$$

with

$$\tau_M \oplus \nu_M = \epsilon_{V \oplus \mathbb{R}^m} : M \rightarrow BO^{\mathbb{Z}_2}(V \oplus \mathbb{R}^m) .$$

The \mathbb{Z}_2 -equivariant Pontryagin-Thom Umkehr map

$$\alpha_M : (V \oplus \mathbb{R}^m)^\infty = \Sigma^m V^\infty \rightarrow M/\text{cl.}(M \setminus E(\nu_M)) = T(\nu_M)$$

represents an element

$$\alpha_M \in \omega_0^{\mathbb{Z}_2}(M; -\tau_M) = \omega_m^{\mathbb{Z}_2}(M; \nu_M - \epsilon_V) = \{(V \oplus \mathbb{R}^m)^\infty; T(\nu_M)\}_{\mathbb{Z}_2} .$$

Proof. (i) See Mostow [45], the generalization of the nonequivariant embedding theorem of Whitney [72].

(ii) Combine (i) and the tubular neighbourhood theorem (4.67). \square

Definition 4.69. Let V be a finite-dimensional inner product \mathbb{Z}_2 -space.

(i) (Costenoble [10, p.156]) A \mathbb{Z}_2 -equivariant $|V|$ -bundle ξ over a \mathbb{Z}_2 -space X is V -restricted if for some $n \geq 0$ there exists a \mathbb{Z}_2 -equivariant \mathbb{R}^n -bundle η over X such that

$$\xi \oplus \eta \cong \epsilon_{V \oplus \mathbb{R}^n} .$$

(ii) A \mathbb{Z}_2 -equivariant m -dimensional manifold M is V -restricted if for some $n \geq 0$ there exists a \mathbb{Z}_2 -equivariant embedding $M \subset V \oplus \mathbb{R}^n$, in which case the normal \mathbb{Z}_2 -equivariant $|V|$ -bundle $\nu_M = \nu_{M \subset V \oplus \mathbb{R}^n}$ is such that

$$\tau_M \oplus \nu_M = \epsilon_{V \oplus \mathbb{R}^n} ,$$

so that ν_M is a V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundle. \square

Proposition 4.70. (i) If $0 \leq j \leq k \leq i$ the trivial \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle $\epsilon_{\mathbb{R}^k \oplus L\mathbb{R}^{i-k}}$ over any \mathbb{Z}_2 -space X is $\mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ -restricted.

(ii) If ξ is V -restricted and ξ' is V' -restricted, then $\xi \oplus \xi'$ is $V \oplus V'$ -restricted.

(iii) The pullback of a V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundle ξ over a \mathbb{Z}_2 -space X along a \mathbb{Z}_2 -equivariant map $f : Y \rightarrow X$ is a V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundle $f^*\xi$ over a \mathbb{Z}_2 -space Y .

(iv) If X is a space with trivial \mathbb{Z}_2 -action and ξ is a \mathbb{Z}_2 -equivariant $|V|$ -bundle over X with the trivial \mathbb{Z}_2 -action (i.e. a nonequivariant $|V|$ -bundle) then ξ is

V-restricted.

(v) If X is a space with a free \mathbb{Z}_2 -action then every \mathbb{Z}_2 -equivariant $|V|$ -bundle ξ over X is *V*-restricted.

(vi) The canonical \mathbb{Z}_2 -equivariant $|V|$ -bundle $\xi = \xi(|V|, V \oplus \mathbb{R}^p)$ over the Grassmann \mathbb{Z}_2 -manifold $G(|V|, V \oplus \mathbb{R}^p)$ is *V*-restricted, for any $p \geq 0$.

Proof. (i) It is immediate from the identity

$$(\mathbb{R}^k \oplus L\mathbb{R}^{i-k}) \oplus L\mathbb{R}^{k-j} = (\mathbb{R}^j \oplus L\mathbb{R}^{i-j}) \oplus \mathbb{R}^{k-j}$$

that

$$\epsilon_{\mathbb{R}^k \oplus L\mathbb{R}^{i-k}} \oplus \epsilon_{L\mathbb{R}^{k-j}} = \epsilon_{\mathbb{R}^j \oplus L\mathbb{R}^{i-j}} \oplus \epsilon_{\mathbb{R}^{k-j}} : X \rightarrow BO^{\mathbb{Z}_2}(\mathbb{R}^k \oplus L\mathbb{R}^{i-j}) .$$

(ii) If $\xi \oplus \eta = \epsilon_{V \oplus \mathbb{R}^p}$ and $\xi' \oplus \eta' = \epsilon_{V' \oplus \mathbb{R}^{p'}}$ then

$$(\xi \oplus \xi') \oplus (\eta \oplus \eta') = \epsilon_{(V \oplus V') \oplus \mathbb{R}^{p+p'}} : X \rightarrow BO^{\mathbb{Z}_2}(V \oplus V' \oplus \mathbb{R}^{p+p'}) .$$

(iii) If $\xi \oplus \eta = \epsilon_{V \oplus \mathbb{R}^p}$ over X then

$$f^*\xi \oplus f^*\eta = \epsilon_{V \oplus \mathbb{R}^p} : Y \rightarrow BO^{\mathbb{Z}_2}(V \oplus \mathbb{R}^p) .$$

(iv) If $\xi \oplus \eta = \epsilon_{\mathbb{R}^p}$ nonequivariantly then

$$\xi \oplus (\epsilon_V \oplus \eta) = \epsilon_{V \oplus \mathbb{R}^p} : X \rightarrow BO^{\mathbb{Z}_2}(V \oplus \mathbb{R}^p) .$$

(v) Let $p : X \rightarrow X/\mathbb{Z}_2$ be the covering projection, and let ξ_0 be the nonequivariant $|V|$ -bundle over X/\mathbb{Z}_2 with

$$E(\xi_0) = E(\xi)/\mathbb{Z}_2 .$$

Regarded as a \mathbb{Z}_2 -equivariant $|V|$ -bundle via the trivial \mathbb{Z}_2 -action ξ_0 is *V*-restricted by (iv), and $\xi = p^*\xi_0$ is *V*-restricted by (iii).

(vi) The canonical \mathbb{Z}_2 -equivariant \mathbb{R}^p -bundle $\eta = \xi(\mathbb{R}^p, V \oplus \mathbb{R}^p)$ over $G(|V|, V \oplus \mathbb{R}^p)$ is such that

$$\xi \oplus \eta = \epsilon_{V \oplus \mathbb{R}^p} : G(|V|, V \oplus \mathbb{R}^p) \rightarrow BO^{\mathbb{Z}_2}(V \oplus \mathbb{R}^p) .$$

□

Example 4.71. For any $p \geq 0$, $q \geq 1$ the canonical \mathbb{Z}_2 -equivariant $|V|$ -bundle $\xi = \xi(|V|, V \oplus \mathbb{R}^p \oplus \mathbb{R}^q)$ over the Grassmann \mathbb{Z}_2 -manifold $G(|V|, V \oplus \mathbb{R}^p \oplus \mathbb{R}^q)$ is not *V*-restricted.

□

Proposition 4.72. *Let $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ be an i -dimensional inner product \mathbb{Z}_2 -space, and let ξ be a V -restricted \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle over a \mathbb{Z}_2 -space X .*

(i) *The spaces*

$$X(\xi, k) = \{x \in X^{\mathbb{Z}_2} \mid \dim((\xi_x)_+) = k\} \subseteq X^{\mathbb{Z}_2}$$

are such that

$$X(\xi, k) = \emptyset \text{ for } k > i \text{ and } k < j ,$$

with

$$X^{\mathbb{Z}_2} = \coprod_{k=j}^i X(\xi, k) .$$

(ii) *The identity $\xi \oplus \eta = \epsilon_{V \oplus \mathbb{R}^p}$ determines identities*

$$\begin{aligned} (\xi, k)_- \oplus (\eta, j - k + p)_- &= \epsilon_{\mathbb{R}^{i-j}} : \\ X(\xi, k) &= X(\eta, j - k + p) \rightarrow BO(\mathbb{R}^{i-j}) \quad (j \leq k \leq i) , \end{aligned}$$

and hence maps

$$X(\xi, k) \rightarrow G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) = G(\mathbb{R}^{k-j}, \mathbb{R}^{i-j})$$

such that

$$\begin{aligned} (\xi, k)_- : X(\xi, k) &\rightarrow G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) \subset BO(\mathbb{R}^{i-k}) , \\ (\eta, j - k + p)_- &= (\xi, k)_-^\perp : X(\xi, k) \rightarrow G(\mathbb{R}^{k-j}, \mathbb{R}^{i-j}) \subset BO(\mathbb{R}^{k-j}) . \end{aligned}$$

□

Definition 4.73. (i) The *classifying space* for V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundles is the \mathbb{Z}_2 -space

$$bo^{\mathbb{Z}_2}(V) = \varinjlim_p G(|V|, V \oplus \mathbb{R}^p) \subset BO^{\mathbb{Z}_2}(V) .$$

Note that nonequivariantly $bo^{\mathbb{Z}_2}(V) = BO(|V|)$.

(ii) The *canonical V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundle* over $bo^{\mathbb{Z}_2}(V)$ is

$$\begin{aligned} \Xi(V) &= \varinjlim_p \xi(V, V \oplus \mathbb{R}^p) : \\ |V| \rightarrow eo^{\mathbb{Z}_2}(V) &= \varinjlim_p E(\xi(V, V \oplus \mathbb{R}^p)) \rightarrow bo^{\mathbb{Z}_2}(V) . \end{aligned}$$

(iii) The *universal restricted Thom \mathbb{Z}_2 -spectrum* is

$$\underline{mo}^{\mathbb{Z}_2} = \{mo^{\mathbb{Z}_2}(V) \mid V\}$$

with

$$mo^{\mathbb{Z}_2}(V) = T(\Xi(V)) = \varinjlim_p MO^{\mathbb{Z}_2}(V, V \oplus \mathbb{R}^p) .$$

□

Proposition 4.74. *Let $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ be an i -dimensional inner product \mathbb{Z}_2 -space.*

(i) *The V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundles over X*

$$\Xi : |V| \longrightarrow E(\Xi) \longrightarrow X$$

are classified by the \mathbb{Z}_2 -equivariant homotopy classes of \mathbb{Z}_2 -equivariant maps

$$f : X \rightarrow G(|V|, V \oplus \mathbb{R}^p) \subset bo^{\mathbb{Z}_2}(V) \text{ (} p \text{ large) ,}$$

with

$$\Xi = f^*\Xi(V) , \quad E(\Xi) = f^*E(\Xi(V)) = \{(x, y) \in X \times V \oplus \mathbb{R}^p \mid y \in f(x)\} .$$

The function

$$\begin{aligned} [X, bo^{\mathbb{Z}_2}(V)]_{\mathbb{Z}_2} &\rightarrow \\ \{ \text{isomorphism classes of } V\text{-restricted } \mathbb{Z}_2\text{-equivariant } |V|\text{-bundles over } X \} ; \\ (f : X \rightarrow bo^{\mathbb{Z}_2}(V)) &\mapsto f^*\Xi(V) \end{aligned}$$

is a bijection.

(ii) *For $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ and any $p \geq 0$ the fixed point set of $G(|V|, V \oplus \mathbb{R}^p)$ is*

$$G(|V|, V \oplus \mathbb{R}^p)^{\mathbb{Z}_2} = \prod_{k=j}^i G(\mathbb{R}^k, \mathbb{R}^{j+p}) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) .$$

The canonical V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundle $\xi(|V|, V \oplus \mathbb{R}^p)$ restricts to

$$\begin{aligned} \xi(|V|, V \oplus \mathbb{R}^p) &= \prod_{k=j}^i \xi(\mathbb{R}^k, \mathbb{R}^{j+p}) \oplus L\xi(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) : \\ \prod_{k=j}^i G(\mathbb{R}^k, \mathbb{R}^{j+p}) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) &\rightarrow BO^{\mathbb{Z}_2}(V) \end{aligned}$$

so that

$$\begin{aligned}
T(\xi(|V|, V \oplus \mathbb{R}^p))^{\mathbb{Z}_2} &= T(\xi(|V|, V \oplus \mathbb{R}^p))^{\mathbb{Z}_2} \\
&= \bigvee_{k=j}^i T(\xi(\mathbb{R}^k, \mathbb{R}^{j+p})) \wedge G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j})^+, \\
|\Xi(V)| &= \coprod_{k=j}^i \Xi(k)_+ \oplus \Xi(k)_- : \\
bo^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2} &= \varinjlim_p \left(\coprod_{k=j}^i G(\mathbb{R}^k, \mathbb{R}^{j+p}) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) \right) \\
&= \coprod_{k=j}^i BO(\mathbb{R}^k) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) \rightarrow bo^{\mathbb{Z}_2}(V), \\
bo^{\mathbb{Z}_2}(V)(\Xi(V), k) &= BO(\mathbb{R}^k) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j}) \quad (j \leq k \leq i), \\
T(\Xi(V))^{\mathbb{Z}_2} &= mo^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2} = \bigvee_{k=j}^i MO(k) \wedge G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j})^+.
\end{aligned}$$

(iv) If the \mathbb{Z}_2 -action on X is trivial then

$$[X, bo^{\mathbb{Z}_2}(V)]_{\mathbb{Z}_2} = [X, bo^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2}] = [X, \coprod_{k=j}^i BO(\mathbb{R}^k) \times G(\mathbb{R}^{i-k}, \mathbb{R}^{i-j})].$$

(v) Let X be a free \mathbb{Z}_2 -space, and let $p : X \rightarrow X/\mathbb{Z}_2$ be the covering projection. By Proposition 4.65 (iv) a \mathbb{Z}_2 -equivariant $|V|$ -bundle over X is of the type $\xi = p^*\xi_0$ for a nonequivariant $|V|$ -bundle

$$\xi_0 : X/\mathbb{Z}_2 \rightarrow G(|V|, \mathbb{R}^p) \subset BO(|V|) \quad (p \text{ large}).$$

The composite \mathbb{Z}_2 -equivariant map

$$\xi : X \xrightarrow{p} X/\mathbb{Z}_2 \xrightarrow{\xi_0} G(|V|, \mathbb{R}^p) \subset G(|V|, V \oplus \mathbb{R}^p) \subset bo^{\mathbb{Z}_2}(V)$$

classifies ξ , so that ξ is V -restricted. The forgetful maps

$$[X, bo^{\mathbb{Z}_2}(V)]_{\mathbb{Z}_2} \rightarrow [X, BO^{\mathbb{Z}_2}(V)]_{\mathbb{Z}_2} \rightarrow [X/\mathbb{Z}_2, BO(V)]$$

are thus bijections. □

Remark 4.75. It is immediate from Proposition 4.74 that a \mathbb{Z}_2 -equivariant $|V|$ -bundle ξ over a \mathbb{Z}_2 -space X is V -restricted if and only if the classifying \mathbb{Z}_2 -equivariant map $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$ can be compressed (up to \mathbb{Z}_2 -equivariant homotopy) into $bo^{\mathbb{Z}_2}(V) \subset BO^{\mathbb{Z}_2}(V)$. □

Example 4.76. (i) For any $p, q \geq 0$ the Grassmann \mathbb{Z}_2 -manifold of 1-dimensional subspaces $\mathbb{R} \subset \mathbb{R}^p \oplus L\mathbb{R}^{q+1}$ is the $(p+q)$ -dimensional real projective space

$$G(\mathbb{R}, \mathbb{R}^p \oplus L\mathbb{R}^{q+1}) = P(\mathbb{R}^p \oplus L\mathbb{R}^{q+1})$$

with the \mathbb{Z}_2 -action inherited from $\mathbb{R}^p \oplus L\mathbb{R}^{q+1}$, and the canonical \mathbb{Z}_2 -equivariant \mathbb{R} -bundle $\xi(L\mathbb{R}, \mathbb{R}^p \oplus L\mathbb{R}^{q+1})$ such that

$$\begin{aligned} S(\xi(\mathbb{R}, \mathbb{R}^p \oplus L\mathbb{R}^{q+1})) &= S(\mathbb{R}^p \oplus L\mathbb{R}^{q+1}) = S^{p-1} * LS^q, \\ T(\xi(\mathbb{R}, \mathbb{R}^p \oplus L\mathbb{R}^{q+1})) &= P(\mathbb{R}^{p+1} \oplus L\mathbb{R}^{q+1}). \end{aligned}$$

(ii) The classifying space for \mathbb{Z}_2 -equivariant \mathbb{R} -bundles

$$BO^{\mathbb{Z}_2}(L\mathbb{R}) = \varinjlim_{p,q} G(\mathbb{R}, \mathbb{R}^p \oplus L\mathbb{R}^{q+1})$$

is the infinite-dimensional projective \mathbb{Z}_2 -space of 1-dimensional subspaces $\mathbb{R} \subset \mathbb{R}^p \oplus L\mathbb{R}^{q+1}$ (p, q large), with the canonical \mathbb{Z}_2 -equivariant \mathbb{R} -bundle $\xi(L\mathbb{R})$ such that

$$\begin{aligned} S(\xi(L\mathbb{R})) &= S(\mathbb{R}(\infty) \oplus L\mathbb{R}(\infty)) = S(\mathbb{R}(\infty)) * S(\infty), \\ T(\xi(L\mathbb{R})) &= P(\mathbb{R}(\infty) \oplus L\mathbb{R}(\infty)). \end{aligned}$$

The classifying space for \mathbb{Z}_2 -equivariant \mathbb{R} -bundles over \mathbb{Z}_2 -spaces with trivial \mathbb{Z}_2 -action is

$$BO^{\mathbb{Z}_2}(L\mathbb{R})^{\mathbb{Z}_2} = \varinjlim_p G(\mathbb{R}, \mathbb{R}^p) \sqcup \varinjlim_q G(\mathbb{R}, L\mathbb{R}^{q+1}) = BO(\mathbb{R}) \sqcup BO(\mathbb{R}).$$

(iii) The classifying space for $L\mathbb{R}$ -restricted \mathbb{Z}_2 -equivariant \mathbb{R} -bundles

$$bo^{\mathbb{Z}_2}(L\mathbb{R}) = \varinjlim_p G(\mathbb{R}, \mathbb{R}^p \oplus L\mathbb{R})$$

is the infinite-dimensional projective \mathbb{Z}_2 -space of 1-dimensional subspaces $\mathbb{R} \subset \mathbb{R}^p \oplus L\mathbb{R}$ (p large), with the canonical $L\mathbb{R}$ -restricted \mathbb{Z}_2 -equivariant \mathbb{R} -bundle $\Xi(L\mathbb{R})$ such that

$$\begin{aligned} S(\Xi(L\mathbb{R})) &= S(\mathbb{R}(\infty) \oplus L\mathbb{R}) = S(\mathbb{R}(\infty)) * S(L\mathbb{R}), \\ T(\Xi(L\mathbb{R})) &= P(\mathbb{R}(\infty) \oplus L\mathbb{R}). \end{aligned}$$

The classifying space for $L\mathbb{R}$ -restricted \mathbb{Z}_2 -equivariant \mathbb{R} -bundles over \mathbb{Z}_2 -spaces with trivial \mathbb{Z}_2 -action is

$$bo^{\mathbb{Z}_2}(\mathbb{R})^{\mathbb{Z}_2} = \varinjlim_p G(\mathbb{R}, \mathbb{R}^p) \sqcup G(\mathbb{R}, L\mathbb{R}) = BO(\mathbb{R}) \sqcup \{*\}.$$

(iv) For any $q \geq 1$ let $\xi_{\mathbb{R}} : P(\mathbb{R}^q) \rightarrow BO(\mathbb{R})$ be the canonical \mathbb{R} -bundle, with $S(\xi_{\mathbb{R}}) = S^{q-1}$. The \mathbb{Z}_2 -equivariant \mathbb{R} -bundle $L\xi_{\mathbb{R}} : P(\mathbb{R}^q) \rightarrow BO^{\mathbb{Z}_2}(\mathbb{R})$ is an $L\mathbb{R}$ -restricted \mathbb{Z}_2 -equivariant $L\mathbb{R}$ -bundle for $q = 1$ (when $L\xi_{\mathbb{R}} = \epsilon_{L\mathbb{R}}$ over

$P(\mathbb{R}) = \{*\}$, but not for $q \geq 2$.

□

Proposition 4.77. (i) *The universal restricted Thom \mathbb{Z}_2 -spectrum $\underline{mo}^{\mathbb{Z}_2}$ is split, and such that*

$$|\underline{mo}^{\mathbb{Z}_2}| = \underline{MO} ,$$

the universal Thom spectrum.

(ii) *The homotopy orbit \mathbb{Z}_2 -spectrum of $\underline{mo}^{\mathbb{Z}_2}$ is such that*

$$(\underline{mo}^{\mathbb{Z}_2})_{h\mathbb{Z}_2} \simeq S(\infty)^+ \wedge \underline{mo} .$$

(iii) *The homotopy fixed point \mathbb{Z}_2 -spectrum of $\underline{mo}^{\mathbb{Z}_2}$ is such that the map*

$$S(L\mathbb{R})^+ \wedge (\underline{mo}^{\mathbb{Z}_2})^{\mathbb{Z}_2} \rightarrow (\underline{mo}^{\mathbb{Z}_2})^{h\mathbb{Z}_2}$$

induces isomorphisms of reduced generalized homology groups for any pointed \mathbb{Z}_2 -space X

$$\begin{aligned} (S(L\mathbb{R})^+ \wedge (\underline{mo}^{\mathbb{Z}_2})^{\mathbb{Z}_2})_*(X) &= \bigoplus_{k=0}^{\infty} \widetilde{MO}_k(X^{\mathbb{Z}_2} \wedge BO(*-k)^+) \\ &\cong (\widetilde{MO}^{\mathbb{Z}_2})_*^{h\mathbb{Z}_2}(X) \end{aligned}$$

with an exact sequence

$$\begin{aligned} \cdots \rightarrow \widetilde{MO}_*(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) \rightarrow \widetilde{mo}_*^{\mathbb{Z}_2}(X) \\ \rightarrow \bigoplus_{k=0}^{\infty} \widetilde{MO}_k(X^{\mathbb{Z}_2} \wedge BO(*-k)^+) \rightarrow \widetilde{MO}_{*-1}(S(\infty)^+ \wedge_{\mathbb{Z}_2} X) \rightarrow \cdots \end{aligned}$$

Proof. As in the unrestricted case (4.66).

□

The stable cohomotopy Thom and Euler classes (3.50) have \mathbb{Z}_2 -equivariant versions:

Definition 4.78. (i) *The stable \mathbb{Z}_2 -equivariant cohomotopy Thom class of a \mathbb{Z}_2 -equivariant V -bundle ξ over X*

$$u^{\mathbb{Z}_2}(\xi) = 1 \in \omega_{\mathbb{Z}_2}^0(D(\xi), S(\xi); -\xi) = \omega_{\mathbb{Z}_2}^0(X)$$

is represented by the pointed \mathbb{Z}_2 -map $1 : X^+ \rightarrow S^0$ sending X to the non-base point.

(ii) *The stable \mathbb{Z}_2 -equivariant cohomotopy Euler class of a \mathbb{Z}_2 -equivariant V -bundle ξ over X*

$$\gamma^{\mathbb{Z}_2}(\xi) = z^* u^{\mathbb{Z}_2}(\xi) \in \omega_{\mathbb{Z}_2}^0(X; -\xi) = \{T(\eta); (V \oplus W)^\infty\}_{\mathbb{Z}_2}$$

is represented by the \mathbb{Z}_2 -equivariant map

$$T(\eta) \xrightarrow{z} T(\xi \oplus \eta) = (V \oplus W)^\infty \wedge X^+ \longrightarrow (V \oplus W)^\infty$$

with $\xi \oplus \eta = \epsilon_{V \oplus W}$ and

$$z : E(\eta) \rightarrow E(\xi \oplus \eta) ; (w, x) \mapsto ((0, w), x) .$$

□

Example 4.79. Let V be a finite-dimensional inner product \mathbb{Z}_2 -space. The stable \mathbb{Z}_2 -equivariant cohomotopy Euler class of the trivial \mathbb{Z}_2 -equivariant V -bundle $\epsilon_V : X \rightarrow BO^{\mathbb{Z}_2}(V)$ over a \mathbb{Z}_2 -space X

$$\gamma^{\mathbb{Z}_2}(\epsilon_V) = 0_V \in \omega_{\mathbb{Z}_2}^0(X; -\epsilon_V) = \{X^+; V^\infty\}_{\mathbb{Z}_2}$$

is represented by the pointed \mathbb{Z}_2 -map $0_V : X^+ \rightarrow V^\infty$ sending X to $0_V \in V^\infty$.

□

Example 4.80. (i) The special case

$$X = (\mathbb{R}^j \oplus L\mathbb{R}^{i-j})^\infty = \Sigma^j(LS^{i-j}) , Y = S^0$$

of Proposition 4.41 (i) is the exact sequence

$$\dots \longrightarrow \omega_{i+1,j} \xrightarrow{b} \omega_{i,j} \longrightarrow \omega_i \longrightarrow \omega_{i,j-1} \longrightarrow \dots$$

of Crabb [12, Lemma (4.3)], with

$$\begin{aligned} \omega_i &= \omega_i(\text{pt.}) = \{S^i; S^0\} = \pi_i^S , \\ \omega_{i,j} &= \omega_{i,j}(\text{pt.}) = \{\Sigma^j(LS^{i-j}); S^0\}_{\mathbb{Z}_2} \\ &= \begin{cases} 0 & \text{if } i < 0 \text{ and } j < 0 \\ \tilde{\omega}_j(P(\infty, j-i)) & \text{if } j < -1 \\ \tilde{\omega}_j(P(\infty, j-i)) \oplus \omega_j & \text{if } i \leq j \end{cases} \end{aligned}$$

with $P(\infty, j-i) = P(\infty)/P(\mathbb{R}^{j-i})$ the infinite stunted projective space (4.55), and

$$b = \gamma^{\mathbb{Z}_2}(\epsilon_{L\mathbb{R}}) = 1 \in \omega_{-1,0} = \{S^0; LS^1\}_{\mathbb{Z}_2} = \omega_0 = \mathbb{Z}$$

the \mathbb{Z}_2 -equivariant Euler class of $\epsilon_{L\mathbb{R}}$, represented by $0 : S^0 \rightarrow LS^1$ ([12, Remarks 4.7]).

(ii) The degree defines an isomorphism

$$\omega_0 = \{S^0; S^0\} \xrightarrow{\cong} \mathbb{Z} ; (G : W^\infty \rightarrow W^\infty) \mapsto \text{degree}(G) .$$

By Example 4.26 there is defined an isomorphism

$$\text{bi-degree} : \omega_{0,0} = \{S^0; S^0\}_{\mathbb{Z}_2} \xrightarrow{\cong} \omega_0(P(\infty)) \oplus \omega_0 = \mathbb{Z} \oplus \mathbb{Z} .$$

□

Definition 4.81. An element $c \in O(V, U \oplus V)$ is a linear isometry $c : V \rightarrow U \oplus V$. The \mathbb{Z}_2 -equivariant adjoint of a map $c : X \rightarrow O(V, U \oplus V)$ is the \mathbb{Z}_2 -equivariant pointed map

$$F_c : LV^\infty \wedge X^+ \rightarrow (LU \oplus LV)^\infty ; (v, x) \mapsto c(x)(v) .$$

□

Proposition 4.82. (i) Let V, W be finite-dimensional inner product \mathbb{Z}_2 -spaces. For a stably trivial \mathbb{Z}_2 -equivariant V -bundle ξ and a \mathbb{Z}_2 -equivariant $V \oplus W$ -bundle isomorphism $\delta\xi : \xi \oplus \epsilon_W \cong \epsilon_{V \oplus W}$

$$\gamma^{\mathbb{Z}_2}(\xi) \in \omega_{\mathbb{Z}_2}^0(X; -\xi) = \{X^+; V^\infty\}_{\mathbb{Z}_2}$$

is the stable \mathbb{Z}_2 -equivariant homotopy class of the adjoint of $\delta\xi$

$$\gamma^{\mathbb{Z}_2}(\xi) : V^\infty \wedge X^+ \rightarrow (V \oplus W)^\infty ; (v, x) \mapsto \delta\xi(x)(v) ,$$

exactly as in the nonequivariant case (3.53).

(ii) If $s : Y \rightarrow S(\xi|_Y)$ is a \mathbb{Z}_2 -equivariant section of $p_{S(\xi|_Y)} : S(\xi|_Y) \rightarrow Y$ (for some $Y \subseteq X$) then

$$\gamma^{\mathbb{Z}_2}(\xi) \in \ker(\omega_{\mathbb{Z}_2}^0(X; -\xi) \rightarrow \omega_{\mathbb{Z}_2}^0(Y; -\xi|_Y)) = \text{im}(\omega_{\mathbb{Z}_2}^0(X, Y; -\xi) \rightarrow \omega_{\mathbb{Z}_2}^0(X; -\xi))$$

and there is defined a rel Y Euler class $\gamma^{\mathbb{Z}_2}(\xi, s) \in \omega_{\mathbb{Z}_2}^0(X, Y; -\xi)$ with image $\gamma^{\mathbb{Z}_2}(\xi) \in \omega_{\mathbb{Z}_2}^0(X; -\xi)$.

(iii) The rel Y Euler classes of \mathbb{Z}_2 -equivariant sections $s_0, s_1 : Y \rightarrow S(\xi|_Y)$ which agree on a subspace $Z \subseteq Y$ are such that

$$\begin{aligned} \gamma^{\mathbb{Z}_2}(\xi, s_0) - \gamma^{\mathbb{Z}_2}(\xi, s_1) &\in \ker(\omega_{\mathbb{Z}_2}^0(X, Y; -\xi) \rightarrow \omega_{\mathbb{Z}_2}^0(X, Z; -\xi)) \\ &= \text{im}(\omega_{\mathbb{Z}_2}^{-1}(Y, Z; -\xi) \rightarrow \omega_{\mathbb{Z}_2}^0(X, Y; -\xi)) \end{aligned}$$

and there is a \mathbb{Z}_2 -equivariant difference class

$$\delta(s_0, s_1) \in \omega_{\mathbb{Z}_2}^{-1}(Y, Z; -\xi) = \{\Sigma(Y/Z); V^\infty\}_{\mathbb{Z}_2}$$

with image

$$\gamma^{\mathbb{Z}_2}(\xi, s_0) - \gamma^{\mathbb{Z}_2}(\xi, s_1) \in \omega_{\mathbb{Z}_2}^0(X, Y; -\xi) = \{X/Y; V^\infty\}_{\mathbb{Z}_2} .$$

□

Example 4.83. (i) Let V be a finite-dimensional inner product space. Given a V -bundle $\xi : X \rightarrow BO(V)$ let $\eta : X \rightarrow BO(W)$ be a W -bundle such that

$$\xi \oplus \eta = \epsilon_{V \oplus W} : X \rightarrow BO(V \oplus W) ,$$

so that

$$L\xi \oplus L\eta = \epsilon_{LV \oplus LW} : X \rightarrow BO^{\mathbb{Z}_2}(V \oplus W) .$$

The \mathbb{Z}_2 -equivariant Euler class of $L\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$

$$\gamma^{\mathbb{Z}_2}(L\xi) \in \omega_{\mathbb{Z}_2}^0(X; -L\xi) = \{T(L\eta); (LV \oplus LW)^\infty\}_{\mathbb{Z}_2}$$

is represented by the composite \mathbb{Z}_2 -equivariant map

$$T(L\eta) \xrightarrow{z} T(L\xi \oplus L\eta) = (LV \oplus LW)^\infty \wedge X^+ \rightarrow (LV \oplus LW)^\infty$$

with

$$z : E(L\eta) \rightarrow E(L\xi \oplus L\eta) = (LV \oplus LW) \times X ; (v, x) \mapsto ((0, v), x) .$$

(ii) For $\xi = \epsilon_V$ can take $W = \{0\}$, $\eta = 0$ in (i), so that

$$\gamma^{\mathbb{Z}_2}(\epsilon_{LV}) \in \omega_{\mathbb{Z}_2}^0(X; -\epsilon_{LV}) = \{X^+; LV^\infty\}_{\mathbb{Z}_2}$$

is represented by the \mathbb{Z}_2 -equivariant map

$$X^+ \rightarrow LV^\infty ; x \mapsto 0 , \infty \mapsto \infty .$$

If V is non-zero then ϵ_{LV} does not admit a \mathbb{Z}_2 -equivariant section, and $\gamma^{\mathbb{Z}_2}(\epsilon_{LV})$ is non-zero – see Example 4.80 above.

□

4.5 \mathbb{Z}_2 -equivariant S -duality

We recall the \mathbb{Z}_2 -equivariant S -duality theory of Wirthmüller [75]. (The theory is for G -spaces with G a compact Lie group, although we shall only be concerned with the case $G = \mathbb{Z}_2$). The theory deals with the stable \mathbb{Z}_2 -equivariant homotopy groups

$$\{X; Y\}_{\mathbb{Z}_2} = \varinjlim_U [U^\infty \wedge X, U^\infty \wedge Y]_{\mathbb{Z}_2}$$

as in Definition 4.24, with the direct limit running over all the finite-dimensional inner product \mathbb{Z}_2 -spaces U .

Definition 4.84. Let X, Y be pointed \mathbb{Z}_2 -spaces, and let $U - V$ be a formal difference of finite-dimensional inner product \mathbb{Z}_2 -spaces.

(i) Define *cap* products

$$\begin{aligned} \tilde{\omega}_0^{\mathbb{Z}_2}(X \wedge Y; \epsilon_U - \epsilon_V) \otimes \tilde{\omega}_{\mathbb{Z}_2}^i(X) &\rightarrow \tilde{\omega}_{-i}^{\mathbb{Z}_2}(Y; \epsilon_U - \epsilon_V) ; \\ (\sigma : V^\infty \rightarrow U^\infty \wedge X \wedge Y) \otimes (f : X \rightarrow S^i) &\mapsto ((1 \wedge f)\sigma : V^\infty \rightarrow \Sigma^i U^\infty \wedge Y) . \end{aligned}$$

(ii) An element $\sigma \in \tilde{\omega}_0^{\mathbb{Z}_2}(X \wedge Y; \epsilon_U - \epsilon_V)$ is a \mathbb{Z}_2 -equivariant *S-duality* if the products

$$\sigma \otimes - : \tilde{\omega}_{\mathbb{Z}_2}^i(X) \rightarrow \tilde{\omega}_{-i}^{\mathbb{Z}_2}(Y; \epsilon_U - \epsilon_V) \quad (i \in \mathbb{Z})$$

are isomorphisms. □

Proposition 4.85. (Wirthmüller [75])

(i) If $\sigma \in \tilde{\omega}_0^{\mathbb{Z}_2}(X \wedge Y; \epsilon_U - \epsilon_V)$ is a \mathbb{Z}_2 -equivariant *S-duality* there are induced isomorphisms

$$\begin{aligned} \sigma : \{X \wedge A; B\}_{\mathbb{Z}_2} &\xrightarrow{\cong} \{V^\infty \wedge A; U^\infty \wedge B \wedge Y\}_{\mathbb{Z}_2} ; F \mapsto (F \wedge 1_Y)(\sigma \wedge 1_A) , \\ \sigma : \{A \wedge Y; B\}_{\mathbb{Z}_2} &\xrightarrow{\cong} \{V^\infty \wedge A; U^\infty \wedge X \wedge B\}_{\mathbb{Z}_2} ; G \mapsto (1_X \wedge G)(1_A \wedge \sigma) \end{aligned}$$

for any pointed CW \mathbb{Z}_2 -complexes A, B .

(ii) For any finite pointed CW \mathbb{Z}_2 -complex X there exist a finite-dimensional inner product \mathbb{Z}_2 -space V , a finite pointed CW \mathbb{Z}_2 -complex Y and a \mathbb{Z}_2 -equivariant map $\sigma : V^\infty \rightarrow X \wedge Y$ such that $\sigma \in \tilde{\omega}_0^{\mathbb{Z}_2}(X \wedge Y; -\epsilon_V)$ is a \mathbb{Z}_2 -equivariant *S-duality*, and for any pointed CW \mathbb{Z}_2 -complex B

$$\{X; B\}_{\mathbb{Z}_2} \cong \{V^\infty; B \wedge Y\}_{\mathbb{Z}_2} , \quad \{Y; B\}_{\mathbb{Z}_2} \cong \{V^\infty; B \wedge X\}_{\mathbb{Z}_2} .$$

□

Example 4.86. Let X be a pointed CW complex, let Y a finite pointed CW \mathbb{Z}_2 -complex, and let $i : Y^{\mathbb{Z}_2} \rightarrow Y$ be the inclusion. For any stable \mathbb{Z}_2 -equivariant map $F : X \rightarrow Y$ the fixed point stable map $G = \rho(F) : X \rightarrow Y^{\mathbb{Z}_2}$ is such that F and $\sigma(G) = iG : X \rightarrow Y$ agree on the fixed points, with the relative difference \mathbb{Z}_2 -equivariant map

$$\delta(F, \sigma(G)) : \Sigma S(\infty)^+ \wedge X \rightarrow L\mathbb{R}(\infty)^\infty \wedge Y$$

\mathbb{Z}_2 -equivariantly S -dual (by 4.85 (ii), with $V = \mathbb{R}(\infty)$) to a stable \mathbb{Z}_2 -equivariant map

$$\delta'(F, \sigma(G)) : X \rightarrow S(\infty)^+ \wedge Y$$

such that

$$F - \sigma(G) = \delta'(F, \sigma(G)) \in \text{im}(\delta) = \ker(\rho)$$

in the direct sum system of Proposition 4.44

$$\{X; S(\infty)^+ \wedge Y\}_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \{X; Y\}_{\mathbb{Z}_2} \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \{X; Y^{\mathbb{Z}_2}\} .$$

□

Proposition 4.87. *Let V be a finite-dimensional inner product \mathbb{Z}_2 -space, and let M be an m -dimensional V -restricted \mathbb{Z}_2 -manifold, so that there exists a \mathbb{Z}_2 -equivariant embedding $M \subset V \oplus \mathbb{R}^m$ with a \mathbb{Z}_2 -equivariant normal $|V|$ -bundle ν_M such that*

$$\tau_M \oplus \nu_M = \epsilon_{V \oplus \mathbb{R}^m} .$$

The composite of the \mathbb{Z}_2 -equivariant Pontryagin-Thom map $\alpha : (V \oplus \mathbb{R}^m)^\infty \rightarrow T(\nu_M)$ and the diagonal map $\Delta : T(\nu_M) \rightarrow M^+ \wedge T(\nu_M)$ is a \mathbb{Z}_2 -equivariant map

$$\sigma = \Delta\alpha : (V \oplus \mathbb{R}^m)^\infty \rightarrow M^+ \wedge T(\nu_M)$$

such that

$$\begin{aligned} \sigma \in \omega_0^{\mathbb{Z}_2}(M \times M, 0 \times -\tau_M) &= \omega_m^{\mathbb{Z}_2}(M \times M, 0 \times (\nu_M - \epsilon_V)) \\ &= \omega_0^{\mathbb{Z}_2}(M^+ \wedge T(\nu_M); -\epsilon_{V \oplus \mathbb{R}^m}) \\ &= \{(V \oplus \mathbb{R}^m)^\infty; M^+ \wedge T(\nu_M)\}_{\mathbb{Z}_2} \end{aligned}$$

is a \mathbb{Z}_2 -equivariant S -duality between M^+ and $T(\nu_M)$ inducing Poincaré duality isomorphisms

$$\sigma \cap - : \omega_{\mathbb{Z}_2}^*(M) \cong \omega_{m-*}^{\mathbb{Z}_2}(M, \nu_M - \epsilon_V) .$$

The isomorphism $\omega_{\mathbb{Z}_2}^0(M) \cong \omega_m^{\mathbb{Z}_2}(M; \nu_M - \epsilon_V)$ sends $1 \in \omega_{\mathbb{Z}_2}^0(M) = \{M^+; S^0\}_{\mathbb{Z}_2}$ to

$$\alpha \in \omega_0^{\mathbb{Z}_2}(M, -\tau_M) = \omega_m^{\mathbb{Z}_2}(M, \nu_M - \epsilon_V) = \{(V \oplus \mathbb{R}^m)^\infty; T(\nu_M)\}_{\mathbb{Z}_2} .$$

□

Example 4.88. Let M be an m -dimensional free \mathbb{Z}_2 -manifold, with a \mathbb{Z}_2 -equivariant embedding $M \subset V \oplus \mathbb{R}^m$ as in 4.87, and let $p : M \rightarrow M/\mathbb{Z}_2$ be the double covering projection, so that

$$\begin{aligned}\tau_M &= p^* \tau_{M/\mathbb{Z}} = \epsilon_{V \oplus \mathbb{R}^m} - \nu_M = p^*(\epsilon_{V \oplus \mathbb{R}^m}/\mathbb{Z}_2) - p^*(\nu_M/\mathbb{Z}_2), \\ \tau_{M/\mathbb{Z}} &= \epsilon_{V \oplus \mathbb{R}^m}/\mathbb{Z}_2 - \nu_M/\mathbb{Z}_2.\end{aligned}$$

The normal bundle of an embedding $M/\mathbb{Z}_2 \subset U \oplus \mathbb{R}^m$ (for some finite-dimensional inner product space U) is such that

$$\nu_{M/\mathbb{Z}_2} = \epsilon_{U \oplus \mathbb{R}^m} - \tau_{M/\mathbb{Z}_2} = \epsilon_U - \epsilon_V/\mathbb{Z}_2 + \nu_M/\mathbb{Z}_2,$$

and

$$\begin{aligned}\alpha \in \omega_0^{\mathbb{Z}_2}(M, -\tau_M) &= \{(V \oplus \mathbb{R}^m)^\infty; T(\nu_M)\}_{\mathbb{Z}_2} \\ &= \omega_0(M/\mathbb{Z}_2, -\tau_{M/\mathbb{Z}_2}) = \{(U \oplus \mathbb{R}^m)^\infty; T(\nu_{M/\mathbb{Z}_2})\}\end{aligned}$$

is the Pontryagin-Thom map of $M/\mathbb{Z}_2 \subset U \oplus \mathbb{R}^m$. The \mathbb{Z}_2 -equivariant S -duality between M^+ and $T(\nu_M)$ can also be regarded as a nonequivariant S -duality between $(M/\mathbb{Z}_2)^+$ and $T(\nu_{M/\mathbb{Z}_2})$

$$\begin{aligned}\sigma \in \omega_0^{\mathbb{Z}_2}(M \times M, 0 \times -\tau_M) &= \omega_m^{\mathbb{Z}_2}(M \times T(\nu_M), -\epsilon_V) \\ &= \omega_0(M/\mathbb{Z}_2 \times M/\mathbb{Z}_2, 0 \times_{\mathbb{Z}_2} -\tau_M) \\ &= \omega_m((M/\mathbb{Z}_2)^+ \wedge T(\nu_{M/\mathbb{Z}_2}); -\epsilon_U),\end{aligned}$$

with the \mathbb{Z}_2 -equivariant Poincaré duality isomorphisms

$$\sigma \cap - : \omega_{\mathbb{Z}_2}^*(M) \cong \omega_{m-*}^{\mathbb{Z}_2}(M, \nu_M - \epsilon_V)$$

regarded as nonequivariant Poincaré duality isomorphisms

$$\sigma \cap - : \omega^*(M/\mathbb{Z}_2) \cong \omega_{m-*}(M/\mathbb{Z}_2, \nu_{M/\mathbb{Z}_2} - \epsilon_U).$$

□

Example 4.89. (i) For any finite-dimensional inner product space V the inclusion $S(LV) \subset LV$ is a \mathbb{Z}_2 -equivariant embedding with trivial normal \mathbb{Z}_2 -equivariant \mathbb{R} -bundle $\nu_{S(LV) \subset LV} = \epsilon_{\mathbb{R}}$. The composite of

$$\alpha_{LV} : LV^\infty \rightarrow T(\epsilon_{\mathbb{R}}) = \Sigma S(LV)^+ ; [t, u] \mapsto (t, u)$$

and the diagonal map $\Delta : \Sigma S(LV)^+ \rightarrow S(LV)^+ \wedge \Sigma S(LV)^+$

$$\sigma_{LV} = \Delta \alpha_{LV} : LV^\infty \rightarrow S(LV)^+ \wedge \Sigma S(LV)^+$$

represents a \mathbb{Z}_2 -equivariant S -duality

$$\sigma_{LV} \in \omega_0(S(LV)^+ \wedge \Sigma S(LV)^+; -\epsilon_{LV}).$$

Thus for any pointed CW \mathbb{Z}_2 -complexes A, B there is defined an S -duality isomorphism

$$\{\Sigma S(LV)^+ \wedge A; LV^\infty \wedge B\}_{\mathbb{Z}_2} \rightarrow \{A; S(LV)^+ \wedge B\}_{\mathbb{Z}_2}; F \mapsto (1 \wedge F)(\Delta \alpha_{LV} \wedge 1).$$

The \mathbb{Z}_2 -equivariant S -duality map σ_{LV} induces a $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$\sigma_{LV} : \dot{C}^{cell}(LV^\infty) \rightarrow C^{cell}(S(LV)) \otimes_{\mathbb{Z}} SC^{cell}(S(LV))$$

with adjoint an isomorphism of $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes

$$\begin{aligned} \sigma_{LV} &: \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(SC^{cell}(S(LV)), \dot{C}^{cell}(LV^\infty)) \\ &\xrightarrow{\cong} \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(\dot{C}^{cell}(LV^\infty), C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}^{cell}(LV^\infty)) = C^{cell}(S(LV)). \end{aligned}$$

(ii) For any $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes D, E the isomorphism of (i) determines a chain level \mathbb{Z}_2 -equivariant S -duality isomorphism

$$\begin{aligned} \sigma_{LV} &: \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(SC^{cell}(S(LV)) \otimes_{\mathbb{Z}} D, \dot{C}^{cell}(LV^\infty) \otimes_{\mathbb{Z}} E) \\ &\xrightarrow{\cong} \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(D, C^{cell}(S(LV)) \otimes_{\mathbb{Z}} E). \end{aligned}$$

(iii) The \mathbb{Z}_2 -equivariant S -duality isomorphism

$$\sigma_{LV} : \{\Sigma S(LV)^+ \wedge A; LV^\infty \wedge B\}_{\mathbb{Z}_2} \xrightarrow{\cong} \{A; S(LV)^+ \wedge B\}_{\mathbb{Z}_2}$$

induces the chain level \mathbb{Z}_2 -equivariant S -duality isomorphism of (ii) with $D = C(A)$, $E = C(B)$. A stable \mathbb{Z}_2 -equivariant map $F : \Sigma S(LV)^+ \wedge A \rightarrow LV^\infty \wedge B$ as in 4.85 (ii) induces a $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$f : SC^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(A) \rightarrow \dot{C}^{cell}(LV^\infty) \otimes_{\mathbb{Z}} \dot{C}(B).$$

The S -dual stable \mathbb{Z}_2 -equivariant map

$$G = (1 \wedge F)(\Delta \alpha_{LV} \wedge 1) : A \rightarrow S(LV)^+ \wedge B$$

induces a $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$g : \dot{C}(A) \rightarrow C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(B)$$

which is just the adjoint of f , i.e. f and g correspond under the isomorphism of (ii). Thus there is defined a commutative diagram

$$\begin{array}{ccc}
\{\Sigma S(LV)^+ \wedge A; LV^\infty \wedge B\}_{\mathbb{Z}_2} & \xrightarrow{\text{Hurewicz}} & H' \\
\sigma_{LV} \downarrow \cong & & \sigma_{LV} \downarrow \cong \\
\{A, S(LV)^+ \wedge B\}_{\mathbb{Z}_2} & \xrightarrow{\text{Hurewicz}} & H''
\end{array}$$

where

$$\begin{aligned}
H' &= H_0(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(SC^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(A), C^{cell}(LV^\infty) \otimes_{\mathbb{Z}} \dot{C}(B))) , \\
H'' &= H_0(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(\dot{C}(A), C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(B))) .
\end{aligned}$$

(iv) For $A = B = S^0$ in (iii) there is defined a commutative diagram of isomorphisms

$$\begin{array}{ccc}
\{\Sigma S(LV)^+; LV^\infty\}_{\mathbb{Z}_2} & \xrightarrow{\text{Hurewicz}} & H' \\
\sigma_{LV} \downarrow \cong & \cong & \sigma_{LV} \downarrow \cong \\
\{S^0; S(LV)^+\}_{\mathbb{Z}_2} & \xrightarrow{\text{Hurewicz}} & H''
\end{array}$$

with

$$\begin{aligned}
&\{\Sigma S(LV)^+; LV^\infty\}_{\mathbb{Z}_2} \rightarrow H' = \mathbb{Z} ; \\
&(F' : \Sigma S(LV)^+ \rightarrow LV^\infty) \mapsto \text{semidegree}(F') = (F')_*(1)/2 , \\
&\{S^0; S(LV)^+\}_{\mathbb{Z}_2} \rightarrow H'' = H_0(P(V)) = \mathbb{Z} ; \\
&(F'' : S^0 \rightarrow S(LV)^+) \mapsto (F'')_*(1)/2 .
\end{aligned}$$

□

4.6 \mathbb{Z}_2 -equivariant bordism

A \mathbb{Z}_2 -space M is a union

$$M = M^{\mathbb{Z}_2} \cup (M \setminus M^{\mathbb{Z}_2})$$

of a space with trivial \mathbb{Z}_2 -action and a space with free \mathbb{Z}_2 -action.

Example 4.90. If ξ is a vector bundle over a space X then the \mathbb{Z}_2 -equivariant bundle $L\xi$ over X is such that

$$E(L\xi)^{\mathbb{Z}_2} = X , \quad E(L\xi) \setminus E(L\xi)^{\mathbb{Z}_2} = S(L\xi) \times (0, \infty)$$

with a free \mathbb{Z}_2 -action on $S(L\xi)$.

□

If M is a closed \mathbb{Z}_2 -manifold then $M \setminus M^{\mathbb{Z}_2}$ is an open codimension 0 submanifold, and the fixed point set $M^{\mathbb{Z}_2}$ is a union of closed submanifolds of different dimensions, with corresponding \mathbb{Z}_2 -equivariant normal bundles. More precisely:

Proposition 4.91. *Let M be a closed m -dimensional \mathbb{Z}_2 -manifold.*

(i) *The fixed point set of M is a disjoint union*

$$M^{\mathbb{Z}_2} = \coprod_{k=0}^m M(k)$$

of closed k -dimensional manifolds, such that

$$\begin{aligned} M(k) &= M(\tau_M, k) = \{x \in M^{\mathbb{Z}_2} \mid \dim((\tau_M)_x)_+ = k\} , \\ \tau_{M(k)} &= (\tau_M, k)_+ : M(k) \rightarrow BO(\mathbb{R}^k) , \\ \nu_{M(k) \subset M} &= (\tau_M, k)_- : M(k) \rightarrow BO(\mathbb{R}^{m-k}) , \\ \coprod_{k=0}^m E(L\nu_{M(k) \subset M}) &\subset M \end{aligned}$$

with each $E(L\nu_{M(k) \subset M}) \subset M$ a \mathbb{Z}_2 -equivariant embedding.

(ii) *The exterior of the fixed point set*

$$M' = \text{cl.}(M \setminus \coprod_{k=0}^m D(L\nu_{M(k) \subset M}))$$

is an m -dimensional free \mathbb{Z}_2 -manifold with boundary

$$\partial M' = \coprod_{k=0}^m S(L\nu_{M(k) \subset M}) .$$

The interior $M' \setminus \partial M'$ is homeomorphic to $M \setminus M^{\mathbb{Z}_2}$, such that

$$M = \coprod_{k=0}^m D(L\nu_{M(k) \subset M}) \cup_{\partial M'} M' .$$

(iii) *The \mathbb{Z}_2 -equivariant Umkehr map*

$$F : M^\infty \rightarrow M/M' = T(\nu_{M^{\mathbb{Z}_2} \subset M}) = \bigvee_{k=0}^m T(L\nu_{M(k) \subset M})$$

restricts to the identity on the fixed point set

$$\begin{aligned} F| &= 1 : (M^\infty)^{\mathbb{Z}_2} = (M^{\mathbb{Z}_2})^\infty \\ &\rightarrow T(\nu_{M^{\mathbb{Z}_2} \subset M})^{\mathbb{Z}_2} = (M^{\mathbb{Z}_2})^\infty = \bigvee_{k=0}^m M(k)^\infty . \end{aligned}$$

□

A closed m -dimensional \mathbb{Z}_2 -manifold M can thus be viewed as

$$M = \coprod_{k=0}^m D(L\nu(k)) \cup_{\partial M'} M' , \quad \nu(k) = \nu_{M(k) \subset M} ,$$

consisting of :

- (i) a collection $\{M(k) \mid 0 \leq k \leq m\}$ of closed k -dimensional manifolds,
- (ii) a collection $\{\nu(k) : M(k) \rightarrow BO(\mathbb{R}^{m-k}) \mid 0 \leq k \leq m\}$ of \mathbb{R}^{m-k} -bundles, so that each $S(L\nu(k))$ is an $(m-1)$ -dimensional free \mathbb{Z}_2 -manifold,
- (iii) an m -dimensional free \mathbb{Z}_2 -manifold M' with boundary $\partial M' = \coprod_{k=0}^m S(L\nu(k))$.

Proposition 4.92. *Let $M^m \subset N^n$ be a \mathbb{Z}_2 -equivariant embedding of a closed m -dimensional \mathbb{Z}_2 -manifold M in a closed n -dimensional \mathbb{Z}_2 -manifold N with normal \mathbb{Z}_2 -equivariant \mathbb{R}^{n-m} -bundle $\nu_{M \subset N} : M \rightarrow BO^{\mathbb{Z}_2}(\mathbb{R}^{n-m})$ and $E(\nu_{M \subset N}) \subset N$.*

(i) *The fixed point sets of M and N are such that*

$$M^{\mathbb{Z}_2} = M \cap N^{\mathbb{Z}_2} = \coprod_{k=0}^m M(k) \subseteq N^{\mathbb{Z}_2} = \coprod_{\ell=0}^n N(\ell) .$$

For $k \leq \ell$ the k -dimensional manifold defined by

$$\begin{aligned} M(k, \ell) &= M(k) \cap N(\ell) \\ &= \{x \in M^{\mathbb{Z}_2} \mid \dim((\tau_M)_x)_+ = k, \dim((\nu_{M \subset N})_x)_+ = \ell - k\} \end{aligned}$$

is a union of components of $M(k)$ (possibly empty), with

$$\begin{aligned} \tau_{M(k, \ell)} &= (\tau_M|)_+ : M(k, \ell) \rightarrow BO(\mathbb{R}^k) , \\ \nu_{M(k, \ell) \subset N(\ell)} &= (\nu_{M \subset N}|)_+ : M(k, \ell) \rightarrow BO(\mathbb{R}^{\ell-k}) \end{aligned}$$

(ii) *The \mathbb{Z}_2 -equivariant Pontryagin-Thom Umkehr map of $N \subset M$*

$$\alpha : N^\infty \rightarrow N/(N \setminus E(\nu_{M \subset N})) = T(\nu_{M \subset N}) .$$

restricts on the fixed point sets to

$$\alpha| = \bigvee_{k \leq \ell} \alpha(k, \ell) : (N^\infty)^{\mathbb{Z}_2} = \bigvee_{\ell} N(\ell)^\infty \rightarrow \bigvee_{k \leq \ell} T(\nu_{M(k, \ell) \subset N(\ell)})$$

with

$$\alpha(k, \ell) : N(\ell)^\infty \rightarrow T(\nu_{M(k, \ell) \subset N(\ell)})$$

the nonequivariant Pontryagin-Thom Umkehr map of $M(k, \ell) \subset N(\ell)$.

□

Proposition 4.93. *Let $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$, and let M be an m -dimensional \mathbb{Z}_2 -manifold, with a \mathbb{Z}_2 -equivariant embedding $M \subset V \oplus \mathbb{R}^m$.*

(i) *The normal \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle $\nu_M : M \rightarrow BO^{\mathbb{Z}_2}(V)$ is such that*

$$\tau_M \oplus \nu_M = \epsilon_{V \oplus \mathbb{R}^m} : M \rightarrow BO^{\mathbb{Z}_2}(V \oplus \mathbb{R}^m) ,$$

so that ν_M is V -restricted, $\nu_M : M \rightarrow bo^{\mathbb{Z}_2}(V)$.

(ii) *The embedding $M \subset V \oplus \mathbb{R}^m$ restricts to an embedding of the fixed point sets*

$$M^{\mathbb{Z}_2} = \prod_{k=\max(0, m-i+j)}^m M(k) \subset (V \oplus \mathbb{R}^m)^{\mathbb{Z}_2} = \mathbb{R}^{j+m}$$

with

$$\nu_M = \nu_{M \subset V \oplus \mathbb{R}^m} : M \rightarrow G(\mathbb{R}^i, V \oplus \mathbb{R}^m) \subset bo^{\mathbb{Z}_2}(V) ,$$

$$M(k) = M(\tau_M, k) = M(\nu_M, j - k + m) ,$$

$$\nu_M|_{M(k)} = (\nu_M, j - k + m) = (\nu_M, j - k + m)_+ \times L(\nu_M, j - k + m)_- :$$

$$M(k) \rightarrow G(\mathbb{R}^{j-k+m}, \mathbb{R}^{j+m}) \times G(\mathbb{R}^{i-j+k-m}, \mathbb{R}^{i-j}) \subset bo^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2} ,$$

$$\tau_{M(k)} = (\tau_M, k)_+ = (\nu_M, j - k + m)_+^\perp :$$

$$M(k) \rightarrow G(\mathbb{R}^k, \mathbb{R}^{j+m}) \subset BO(\mathbb{R}^k) ,$$

$$\nu_{M(k) \subset M} = (\tau_M, k)_- = (\nu_M, j - k + m)_-^\perp :$$

$$M(k) \rightarrow G(\mathbb{R}^{m-k}, \mathbb{R}^{i-j}) \subset BO(\mathbb{R}^{m-k}) ,$$

$$\nu_{M(k) \subset V \oplus \mathbb{R}^m} = \nu_{M(k) \subset \mathbb{R}^{j+m}} \times L\nu_{\mathbb{R}^{j+m} \subset V \oplus \mathbb{R}^m}|_{M(k)} =$$

$$(\nu_M, j - k + m)_+ \times L\epsilon_{\mathbb{R}^{i-j}} :$$

$$M(k) \rightarrow BO(\mathbb{R}^{j-k+m}) \times BO(\mathbb{R}^{i-j}) \subset bo^{\mathbb{Z}_2}(V \oplus \mathbb{R}^{m-k}) ,$$

$$\tau_M|_{M(k)} = (\tau_M, k)_+ \oplus (\tau_M, k)_- :$$

$$M(k) \rightarrow G(\mathbb{R}^k, \mathbb{R}^{j+m}) \times G(\mathbb{R}^{m-k}, \mathbb{R}^{i-j}) \subset BO(\mathbb{R}^m) .$$

(iii) *By the \mathbb{Z}_2 -equivariant tubular neighbourhood theorem there is an extension of $M \subset V \oplus \mathbb{R}^m$ to a codimension 0 \mathbb{Z}_2 -equivariant embedding $E(\nu_M) \subset V \oplus \mathbb{R}^m$. The \mathbb{Z}_2 -equivariant Pontryagin-Thom Umkehr map of $M \subset V \oplus \mathbb{R}^m$*

$$\alpha : (V \oplus \mathbb{R}^m)^\infty \rightarrow V \oplus \mathbb{R}^m / (V \oplus \mathbb{R}^m \setminus E(\nu_M)) = D(\nu_M) / S(\nu_M) = T(\nu_M)$$

restricts on the fixed point sets to the one-point union of the nonequivariant Pontryagin-Thom Umkehr maps of $M(k) \subset \mathbb{R}^{j+m}$

$$\alpha(k) : S^{j+m} \rightarrow T(\nu_{M(k) \subset \mathbb{R}^{j+m}}) = T((\nu_M, j - k + m)_+) ,$$

that is

$$\begin{aligned} \alpha| &= \bigvee_{k=m-i+j}^m \alpha(k) : \\ ((V \oplus \mathbb{R}^m)^\infty)^{\mathbb{Z}_2} &= S^{j+m} \rightarrow T(\nu_M)^{\mathbb{Z}_2} = \bigvee_{k=m-i+j}^m T((\nu_M, j - k + m)_+) . \end{aligned}$$

□

Example 4.94. An inner product \mathbb{Z}_2 -space $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ is a \mathbb{Z}_2 -equivariant tubular neighborhood of the \mathbb{Z}_2 -equivariant submanifold $V_+ = \mathbb{R}^j \subset V$ with

$$\nu_{V_+ \subset V} = \epsilon_{L\mathbb{R}^{i-j}} , \quad E(\nu_{V_+ \subset V}) = V ,$$

and the \mathbb{Z}_2 -equivariant Pontryagin-Thom Umkehr map is

$$\alpha = 1 : V^\infty = S^j \wedge LS^{i-j} \rightarrow T(\nu_{V_+ \subset V}) = S^j \wedge LS^{i-j} .$$

□

Example 4.95. Let N be an n -dimensional manifold with an embedding $N \subset \mathbb{R}^{j+2n}$, so that

$$\begin{aligned} \tau_N &: N \rightarrow G(\mathbb{R}^n, \mathbb{R}^{j+2n}) \subset BO(\mathbb{R}^n) , \\ \nu_N &= \nu_{N \subset \mathbb{R}^{j+2n}} = \tau_N^\perp : N \rightarrow G(\mathbb{R}^{j+n}, \mathbb{R}^{j+2n}) \subset BO(\mathbb{R}^{j+n}) , \\ \tau_N \oplus \nu_N &= \epsilon_{\mathbb{R}^{j+2n}} : N \rightarrow BO(\mathbb{R}^{j+2n}) . \end{aligned}$$

The $2n$ -dimensional \mathbb{Z}_2 -manifold $M = N \times N$ with

$$T : M \rightarrow M ; (x, y) \mapsto (y, x)$$

is such that

$$\begin{aligned}
M &\subset V \oplus \mathbb{R}^{2n} \text{ with } V = \mathbb{R}^j \oplus L\mathbb{R}^{j+2n}, \\
M^{\mathbb{Z}_2} &= M(n) = \{(x, x) \mid x \in N\} = N^n, \\
\tau_M &= \tau_N \times L\tau_N : M \rightarrow G(\mathbb{R}^n, \mathbb{R}^{j+2n}) \times G(\mathbb{R}^n, \mathbb{R}^{j+2n}) \subset BO^{\mathbb{Z}_2}(\mathbb{R}^{2n}), \\
\tau_M|_N &= \tau_N \oplus \nu_{\Delta: N \subset M} = \tau_N \oplus L\tau_N : \\
M(n) &= N \rightarrow BO(\mathbb{R}^n) \times BO(\mathbb{R}^n) \subset BO^{\mathbb{Z}_2}(\mathbb{R}^{2n})^{\mathbb{Z}_2}, \\
\nu_M &= \nu_{M \subset V \oplus \mathbb{R}^{2n}} = \nu_N \times \nu_N : M \rightarrow G(\mathbb{R}^{2j+2n}, V \oplus \mathbb{R}^{2n}) \subset bo^{\mathbb{Z}_2}(V), \\
M(n) &= M(\tau_M, n) = M(\nu_M, j+n) = N, \\
\nu_M|_{M(n)} &= (\nu_M, j+n) = (\nu_M, j+n)_+ \times L(\nu_M, j+n)_- = \nu_N \times \nu_N : \\
M(n) &= N \rightarrow G(\mathbb{R}^{j+n}, \mathbb{R}^{j+2n}) \times G(\mathbb{R}^{j+n}, \mathbb{R}^{j+2n}) \subset bo^{\mathbb{Z}_2}(V)^{\mathbb{Z}_2}, \\
\tau_{M(n)} &= (\tau_M, n)_+ = (\nu_M, j+n)_+^\perp = \tau_N : \\
M(n) &= N \rightarrow G(\mathbb{R}^n, \mathbb{R}^{j+2n}) \subset BO(\mathbb{R}^n), \\
\nu_{M(n) \subset M} &= (\tau_M, n)_- = (\nu_M, j+n)_-^\perp = \tau_N : \\
M(n) &= N \rightarrow G(\mathbb{R}^n, \mathbb{R}^{j+2n}) \subset BO(\mathbb{R}^n), \\
\nu_{M(n) \subset V \oplus \mathbb{R}^{j+2n}} &= \nu_{M(n) \subset \mathbb{R}^{j+2n}} \times L\nu_{\mathbb{R}^{j+2n} \subset V \oplus \mathbb{R}^{2n}}|_{M(n)} = \nu_N \times L\epsilon_{\mathbb{R}^{j+2n}} : \\
M(n) &= N \rightarrow BO(\mathbb{R}^{j+n}) \times BO(\mathbb{R}^{j+2n}) \subset bo^{\mathbb{Z}_2}(V \oplus \mathbb{R}^n), \\
\tau_M|_{M(n)} &= (\tau_M, n)_+ \oplus (\tau_M, n)_- = \tau_N \oplus \tau_N : \\
M(n) &= N \rightarrow G(\mathbb{R}^n, \mathbb{R}^{j+2n}) \times G(\mathbb{R}^n, \mathbb{R}^{j+2n}) \subset BO(\mathbb{R}^{2n}).
\end{aligned}$$

□

Definition 4.96. Let ξ be a \mathbb{Z}_2 -equivariant $|V|$ -bundle over a \mathbb{Z}_2 -space X , for some finite-dimensional inner product \mathbb{Z}_2 -space V .

(i) An m -dimensional \mathbb{Z}_2 -equivariant normal map

$$(f, b) : (M, \nu_M - \epsilon_{V \oplus W}) \rightarrow (X, \xi - \epsilon_V)$$

consists of an m -dimensional \mathbb{Z}_2 -manifold M with a \mathbb{Z}_2 -equivariant embedding $M \subset \mathbb{R}^m \oplus V \oplus W$ for some finite-dimensional inner product \mathbb{Z}_2 -space W and a \mathbb{Z}_2 -equivariant bundle map over f

$$b : \nu_M = \nu_{M \subset \mathbb{R}^m \oplus V \oplus W} \rightarrow \xi \oplus \epsilon_W,$$

or equivalently a bundle map

$$b : \tau_M \oplus f^*\xi \oplus \epsilon_W \rightarrow \epsilon_{\mathbb{R}^m \oplus V \oplus W},$$

with

$$\tau_M \oplus \nu_M = \epsilon_{\mathbb{R}^m \oplus V \oplus W}, \quad \epsilon_{\mathbb{R}^m} - \tau_M = f^*(\xi - \epsilon_V).$$

Let $V \oplus W = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$. The k -dimensional part of the fixed point set $M^{\mathbb{Z}_2}$ is

$$\begin{aligned}
M(k) &= \{x \in M^{\mathbb{Z}_2} \mid \dim((\tau_M)_x)_+ = k\} \\
&= \{x \in M^{\mathbb{Z}_2} \mid \dim((\nu_M)_x)_+ = j - k + m\} \\
&= \{x \in M^{\mathbb{Z}_2} \mid \dim((\xi \oplus \epsilon_W)_{f(x)})_+ = j - k + m\} \\
&= f^{-1}(X(\xi \oplus \epsilon_W, j - k + m)) \quad (0 \leq k \leq m),
\end{aligned}$$

and (f, b) restricts to nonequivariant normal maps

$$\begin{aligned}
(f, b)| &= (f(k), b(k)) : (M(k), \nu_{M(k) \subset \mathbb{R}^{m+j}}) \\
&\rightarrow (X(\xi \oplus \epsilon_W, j - k + m), (\xi \oplus \epsilon_W, j - k + m)_+)
\end{aligned}$$

such that

$$\begin{aligned}
\nu_{M(k) \subset M} &= (\tau_M, k)_- : M(k) \xrightarrow{f(k)} X(\xi \oplus \epsilon_W, j - k + m) \\
&\xrightarrow{(\xi \oplus \epsilon_W, j - k + m)_+^\perp} \\
G(\mathbb{R}^{i-j+k-m}, \mathbb{R}^{i-j}) &= G(\mathbb{R}^{m-k}, \mathbb{R}^{i-j}) \subset BO(\mathbb{R}^{m-k}).
\end{aligned}$$

(ii) Let $\Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V)$ be the bordism group of m -dimensional \mathbb{Z}_2 -equivariant normal maps $(f, b) : (M, \nu_M - \epsilon_{V \oplus W}) \rightarrow (X, \xi - \epsilon_V)$. Use the \mathbb{Z}_2 -equivariant Pontryagin-Thom map

$$\alpha(f, b) : (\mathbb{R}^m \oplus V \oplus W)^\infty \xrightarrow{\alpha_M} T(\nu_M) \xrightarrow{T(b)} T(\xi \oplus \epsilon_W)$$

to define the \mathbb{Z}_2 -equivariant Pontryagin-Thom map

$$\begin{aligned}
PT^{\mathbb{Z}_2} : \Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) &\rightarrow \omega_m^{\mathbb{Z}_2}(X; \xi - \epsilon_V) \\
&= \{(V \oplus \mathbb{R}^m)^\infty; T(\xi)\}_{\mathbb{Z}_2} \\
&= \{(V \oplus W \oplus \mathbb{R}^m)^\infty; T(\xi \oplus \epsilon_W)\}_{\mathbb{Z}_2}; \\
((f, b) : (M, \nu_M - \epsilon_{V \oplus W}) &\rightarrow (X, \xi - \epsilon_V)) \mapsto \alpha(f, b)
\end{aligned}$$

(iii) Let $\Omega_{m,k}^{\mathbb{Z}_2}(X, \xi - \epsilon_V)$ ($j \leq k \leq i$) be the bordism group of m -dimensional \mathbb{Z}_2 -equivariant normal maps $(f, b) : (M, \nu_M - \epsilon_{V \oplus W}) \rightarrow (X, \xi - \epsilon_V)$ with k -dimensional fixed point set $M^{\mathbb{Z}_2} = M(k)$.

(iv) Let $\Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V)$ be the bordism group of m -dimensional \mathbb{Z}_2 -equivariant normal maps $(f, b) : (M, \nu_M - \epsilon_{V \oplus W}) \rightarrow (X, \xi - \epsilon_V)$ with free \mathbb{Z}_2 -action on M , i.e. $M^{\mathbb{Z}_2} = \emptyset$. Let $c \in [M/\mathbb{Z}_2, P(\infty)] = H^1(M; \mathbb{Z}_2)$ classify the double cover $M \rightarrow M/\mathbb{Z}_2$, and let $\bar{c} : M \rightarrow S(\infty)$ be a \mathbb{Z}_2 -equivariant map such that $\bar{c}/\mathbb{Z}_2 = c$. Use the composite \mathbb{Z}_2 -equivariant map

$$\begin{aligned} \alpha_c(f, b) : (V \oplus W \oplus \mathbb{R}^m)^\infty &\xrightarrow{\alpha_M} T(\nu_M) \xrightarrow{\Delta} M^+ \wedge T(\nu_M) \\ &\xrightarrow{\bar{c} \wedge T(b)} S(\infty)^+ \wedge T(\xi \oplus \epsilon_W) \end{aligned}$$

to define the free \mathbb{Z}_2 -equivariant Pontryagin-Thom map

$$\begin{aligned} PT^{\mathbb{Z}_2\text{-free}} : \Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) &\rightarrow \\ \omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) &= \{(V \oplus \mathbb{R}^m)^\infty; S(\infty)^+ \wedge T(\xi)\}_{\mathbb{Z}_2} ; \\ (f, b) &\mapsto \alpha_c(f, b) . \end{aligned}$$

(v) Let $\Omega_m^{\mathbb{Z}_2}(X)$ be the bordism group of m -dimensional \mathbb{Z}_2 -manifolds M with a \mathbb{Z}_2 -equivariant map $f : M \rightarrow X$.

□

Proposition 4.97. *Let $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ be an i -dimensional inner product \mathbb{Z}_2 -space, and let $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$ be a \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle over a \mathbb{Z}_2 -space X , so that*

$$X^{\mathbb{Z}_2} = \prod_{k=0}^i X(\xi, k) .$$

(i) *Define the \mathbb{Z}_2 -equivariant space*

$$F(\xi, V) = \varinjlim_{W, m} \bigcup_{x \in X} O(\xi_x \oplus W, \mathbb{R}^m \oplus V \oplus W)$$

with W running over finite-dimensional inner product \mathbb{Z}_2 -spaces, such that the projection

$$p : F(\xi, V) \rightarrow X ; (x, v) \mapsto x$$

has contractible point inverses (so it is a nonequivariant homotopy equivalence). The fixed point set is

$$F(\xi, V)^{\mathbb{Z}_2} = \bigcup_{x \in X^{\mathbb{Z}_2}} \varinjlim_{W, m} (O((\xi_x)_+ \oplus W_+, \mathbb{R}^m \oplus V_+ \oplus W_+) \times O((\xi_x)_- \oplus W_-, V_- \oplus W_-)) .$$

The map

$$p : F_-(\xi, V) = \bigcup_{x \in X^{\mathbb{Z}_2}} \varinjlim_{W_-} O((\xi_x)_- \oplus W_-, V_- \oplus W_-) \rightarrow X^{\mathbb{Z}_2} ; (x, v) \mapsto x$$

restricts to fibre bundle projections

$$p(k) = p| : F_-(\xi, V, k) = p^{-1}(X(\xi, k)) \rightarrow X(\xi, k)$$

with fibre the $(k - j)$ -connected stable Stiefel manifold

$$\varinjlim_N O(\mathbb{R}^{i-k+N}, \mathbb{R}^{i-j+N}) .$$

The projection onto the second factor defines a map $F(\xi, V)^{\mathbb{Z}_2} \rightarrow F_-(\xi, V)$ with contractible point inverses, which is thus a homotopy equivalence.

(ii) The pullback construction along

$$p(k)^*(\xi, k)^\perp : F_-(\xi, V, k) \rightarrow X(\xi, k) \rightarrow G(\mathbb{R}^k, \mathbb{R}^k) = G(\mathbb{R}^{k-j}, \mathbb{R}^k) \subset BO(\mathbb{R}^{k-j})$$

defines morphisms

$$\partial_\Omega(k) : \Omega_{m-k+j}(F_-(\xi, V, k), p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) \rightarrow \Omega_{m-1}^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V)$$

with

$$\begin{aligned} \partial_\Omega(k)((g, c) : (N, \nu_N - \epsilon_{\mathbb{R}^{\dim(\nu_N)}})) &\rightarrow (F_-(\xi, V, k), p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) \\ &= \left((g, c)^! : S(\nu_N)^! \rightarrow S(p(k)^*(\xi, k)^\perp) \right), \end{aligned}$$

such that there is defined a commutative diagram

$$\begin{array}{ccc} \Omega_{m-k+j}(F_-(\xi, V, k), p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) & \xrightarrow[\cong]{PT} & \omega_{m-k+j}(F_-(\xi, V, k); p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) \\ \downarrow \partial_\Omega(k) & & \downarrow \partial_\omega(k) \\ \Omega_{m-1}^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) & \xrightarrow{PT^{\mathbb{Z}_2\text{-free}}} & \omega_{m-1}^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \end{array}$$

with $\partial_\omega(k) = PT^{\mathbb{Z}_2\text{-free}} \partial_\Omega(k) (PT)^{-1}$.

(iii) The \mathbb{Z}_2 -free Pontryagin-Thom maps

$$PT^{\mathbb{Z}_2\text{-free}} : \Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) \rightarrow \omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V))$$

are isomorphisms.

Proof. (i) By construction.

(ii) By construction, noting that the nonequivariant Pontryagin-Thom map PT is an isomorphism (Proposition 3.34).

(iii) The nonequivariant Pontryagin-Thom map

$$PT : \Omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) \rightarrow \omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V))$$

is an isomorphism. An element $(N, g, c) \in \Omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V))$ is an m -dimensional normal map

$$(g, c) : (N, \nu_N) \rightarrow (S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) .$$

The pullback along g of the double covering projection

$$S(\infty) \times X \rightarrow S(\infty) \times_{\mathbb{Z}_2} X$$

is a double cover $\bar{N} = g^*(S(\infty) \times X)$ of N with a V -restricted \mathbb{Z}_2 -equivariant normal map

$$(\bar{g}, \bar{c}) : (\bar{N}, \nu_{\bar{N}}) \rightarrow (X, \xi - \epsilon_V) .$$

The maps

$$\begin{aligned} \Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) &\rightarrow \Omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) ; \\ &\quad (M, f, b) \rightarrow (M, f, b)/\mathbb{Z}_2 , \\ \Omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) &\rightarrow \Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) ; \\ &\quad (N, g, c) \mapsto (\bar{N}, \bar{g}, \bar{c}) \end{aligned}$$

are inverse isomorphisms. Furthermore, the \mathbb{Z}_2 -action on $S(\infty) \times X$ is free, so by Proposition 4.58 (ii)

$$\omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) = \omega_m(S(\infty) \times_{\mathbb{Z}_2} X; (0 \times (\xi - \epsilon_V))/\mathbb{Z}_2) .$$

□

Proposition 4.98. *Let $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ be an i -dimensional inner product \mathbb{Z}_2 -space, and let $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$ be a \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle over a \mathbb{Z}_2 -space X .*

(i) *The various bordism groups fit into an exact sequence (of the type pioneered by Conner and Floyd [8, §28] and Stong [64])*

$$\begin{array}{c} \dots \longrightarrow \Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) = \Omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) \\ \xrightarrow{\quad \gamma \quad} \Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) \\ \\ \xrightarrow{\quad \rho \quad} \bigoplus_{k=j}^{j+m} \Omega_{m-k+j}(F_-(\xi, V, k), p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) \\ \partial_\Omega = \bigoplus_k \partial_\Omega(k) \\ \xrightarrow{\quad \quad \quad} \Omega_{m-1}^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) \longrightarrow \dots \end{array}$$

with ρ the fixed point map.

(ii) *The various stable homotopy groups fit into an exact sequence*

$$\begin{array}{c}
\vdots \\
\downarrow \\
\omega_m^{\mathbb{Z}_2}(S(\infty) \times X, 0 \times (\xi - \epsilon_V)) \\
\parallel \\
\omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) \\
\parallel \\
\omega_m(S(\infty) \times_{\mathbb{Z}_2} F(\xi, V), 0 \times_{\mathbb{Z}_2} p^*(\xi - \epsilon_V)) \\
\downarrow \gamma \\
\omega_m^{\mathbb{Z}_2}(F(\xi, V), p^*(\xi - \epsilon_V)) \\
\downarrow \rho \\
\omega_m(F(\xi, V)^{\mathbb{Z}_2}, p^*(\xi - \epsilon_V)) \\
\parallel \\
\bigoplus_{k=j}^{j+m} \omega_{m-k+j}(F_-(\xi, V, k), p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) \\
\downarrow \partial_\omega = \bigoplus_k \partial_\omega(k) \\
\omega_{m-1}^{\mathbb{Z}_2}(S(\infty) \times X, 0 \times (\xi - \epsilon_V)) \\
\downarrow \\
\vdots
\end{array}$$

with ρ the fixed point map.

Proof. (i) Let $\Omega_m^{\mathbb{Z}_2, \mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V)$ be the relative bordism group of \mathbb{Z}_2 -manifolds with boundary $(M, \partial M)$ with the \mathbb{Z}_2 -action on ∂M free, designed to fit into the exact sequence

$$\begin{array}{c}
\cdots \longrightarrow \Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) \xrightarrow{\gamma} \Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) \\
\longrightarrow \Omega_m^{\mathbb{Z}_2, \mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) \xrightarrow{\partial} \Omega_{m-1}^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) \longrightarrow \cdots
\end{array}$$

The fixed point map

$$\begin{aligned} \rho &: \Omega_m^{\mathbb{Z}_2, \mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) \rightarrow \\ &\bigoplus_{k=j}^{j+m} \Omega_{m-k+j}(F_-(\xi, V, k), p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) ; (M, \partial M) \mapsto M^{\mathbb{Z}_2} \end{aligned}$$

is an isomorphism, since $(M, \partial M)$ is bordant to a \mathbb{Z}_2 -equivariant tubular neighbourhood normal bundle of $M^{\mathbb{Z}_2} \subset M$ (4.67).

(ii) This is a special case of Proposition 4.30 (ii). \square

Proposition 4.99. *The maps*

$$\begin{aligned} \Omega_m^{\mathbb{Z}_2}(X) &\rightarrow \varinjlim_V \Omega_m^{\mathbb{Z}_2}(X \times BO^{\mathbb{Z}_2}(V), 0 \times (\xi(V) - \epsilon_V)) ; \\ (f : M \rightarrow X) &\mapsto (f \times \nu_M : (M, \nu_M - \epsilon_V) \rightarrow (X \times BO^{\mathbb{Z}_2}(V), \xi(V) - \epsilon_V)) \end{aligned}$$

are isomorphisms, with ν_M the normal bundle of a \mathbb{Z}_2 -equivariant embedding $M \subset \mathbb{R}^m \oplus V$ and V a sufficiently high-dimensional inner product \mathbb{Z}_2 -space. Thus there are defined \mathbb{Z}_2 -equivariant Pontryagin-Thom maps

$$PT^{\mathbb{Z}_2} : \Omega_m^{\mathbb{Z}_2}(X) \rightarrow MO_m^{\mathbb{Z}_2}(X) = \varinjlim_V \omega_m^{\mathbb{Z}_2}(X \times BO^{\mathbb{Z}_2}(V), 0 \times (\xi(V) - \epsilon_V)) .$$

Proof. By the classification of \mathbb{Z}_2 -equivariant bundles in Proposition 4.65. \square

Remark 4.100. It is well-known that the \mathbb{Z}_2 -equivariant Pontryagin-Thom maps $PT^{\mathbb{Z}_2} : \Omega_m^{\mathbb{Z}_2}(X) \rightarrow MO_m^{\mathbb{Z}_2}(X)$ are not isomorphisms (e.g. Sinha [59]). However, the \mathbb{Z}_2 -equivariant Pontryagin-Thom maps for the V -restricted Thom \mathbb{Z}_2 -spectrum $mo_*^{\mathbb{Z}_2}$

$$pt^{\mathbb{Z}_2} : \Omega_m^{\mathbb{Z}_2}(X) \rightarrow mo_m^{\mathbb{Z}_2}(X)$$

are isomorphisms (e.g. Costenoble [10]). This isomorphism will now be generalized to the Thom space of any V -restricted \mathbb{Z}_2 -equivariant $|V|$ -bundle. \square

Proposition 4.101. *Let $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ be an i -dimensional inner product \mathbb{Z}_2 -space, and let $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$ be a \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle over a \mathbb{Z}_2 -space X , so that*

$$X^{\mathbb{Z}_2} = \prod_{k=0}^i X(\xi, k) .$$

(i) *The Pontryagin-Thom maps define an isomorphism of exact sequences*

$$\begin{array}{ccc}
\vdots & & \vdots \\
\downarrow & & \downarrow \\
\Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2\text{-free}}} & \omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
\parallel & & \parallel \\
\Omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) & \xrightarrow[\cong]{PT} & \omega_m(S(\infty) \times_{\mathbb{Z}_2} X; 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) \\
\downarrow \gamma & & \downarrow \\
\Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2}} & \omega_m^{\mathbb{Z}_2}(F(\xi, V); p^*(\xi - \epsilon_V)) \\
\downarrow \rho & & \downarrow \rho \\
& & \omega_m^{\mathbb{Z}_2}(L\mathbb{R}(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
& & \parallel \\
\bigoplus_{k=j}^{j+m} \Omega_{m-k+j}(F_-(\xi, V, k), p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) & \xrightarrow[\cong]{PT} & \bigoplus_{k=j}^{j+m} \omega_{m-k+j}(F_-(\xi, V, k); p(k)^*((\xi, k)_+ - \epsilon_{\mathbb{R}^k})) \\
\downarrow \partial_\Omega & & \downarrow \partial_\omega \\
\Omega_{m-1}^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2\text{-free}}} & \omega_{m-1}^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
\downarrow & & \downarrow \\
\vdots & & \vdots
\end{array}$$

with ρ the fixed point map, and ∂_Ω defined by pullbacks along

$$p(k)^*(\xi, k)_-^\perp : F_-(\xi, V, k) \rightarrow BO(\mathbb{R}^{k-j})$$

with

$$\begin{aligned}
& \partial_\Omega((f, b) : (M, \nu_M - \epsilon_{V \oplus W}) \rightarrow (X, \xi - \epsilon_V)) = \\
& \bigoplus_{k=j}^{j+m} \left((f(j-k+m), b(j-k+m))^\dagger : S(\nu_{M(j-k+m) \subset M}) \rightarrow S((\xi, j-k+m)_-^\perp) \right) \\
& \in \text{im} \left(\bigoplus_{k=j}^{j+m} \Omega_{m-1}(X(\xi, k), (\xi, k)_+ - \epsilon_{\mathbb{R}^k}) \rightarrow \Omega_{m-1}(P(V_-) \times X^{\mathbb{Z}_2}, 0 \times \xi) \right. \\
& \quad \left. \rightarrow \Omega_{m-1}(S(LV_-) \times_{\mathbb{Z}_2} X, e_2(\xi)) \right).
\end{aligned}$$

(i)' If $\xi : X \rightarrow BO^{\mathbb{Z}_2}(V)$ is V -restricted there is defined an exact sequence

$$\begin{array}{c}
\vdots \\
\downarrow \\
\omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
\parallel \\
\omega_m(S(\infty) \times_{\mathbb{Z}_2} X; 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) \\
\downarrow \\
\omega_m^{\mathbb{Z}_2}(X; \xi - \epsilon_V) \\
\downarrow \rho \\
\omega_m^{\mathbb{Z}_2}(L\mathbb{R}(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
\parallel \\
\bigoplus_{k=j}^i \omega_{m-k+j}(X(\xi, k); (\xi, k)_+ - \epsilon_{\mathbb{R}^k}) \\
\downarrow \partial_\omega \\
\omega_{m-1}^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
\downarrow \\
\vdots
\end{array}$$

with ρ the fixed point map.

(ii) If $X(\xi, k) = \emptyset$ for $k \neq p$ then

$$\Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) = \Omega_{m, j+m-p}^{\mathbb{Z}_2}(X, \xi - \epsilon_V)$$

and the Pontryagin-Thom maps define an isomorphism of exact sequences

$$\begin{array}{ccc}
\vdots & & \vdots \\
\downarrow & & \downarrow \\
\Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2\text{-free}}} & \omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times \xi - \epsilon_V) \\
\downarrow \gamma & & \downarrow \\
\Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2}} & \omega_m^{\mathbb{Z}_2}(X; \xi - \epsilon_V) \\
\downarrow \rho & & \downarrow \rho \\
\Omega_{j+m-p}(X(\xi, p), (\xi, p)_+ - \epsilon_{\mathbb{R}^p}) & \xrightarrow[\cong]{PT} & \omega_{j+m-p}(X(\xi, p); (\xi, p)_+ - \epsilon_{\mathbb{R}^p}) \\
\downarrow \partial_\Omega & & \downarrow \partial_\omega \\
\Omega_{m-1}^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2\text{-free}}} & \omega_{m-1}^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
\downarrow & & \downarrow \\
\vdots & & \vdots
\end{array}$$

The Pontrjagin-Thom maps PT , $PT^{\mathbb{Z}_2\text{-free}}$ are isomorphisms by 3.34, 4.97, and the Pontrjagin-Thom maps $PT^{\mathbb{Z}_2}$ are isomorphisms by the 5-lemma.

(iii) If $X(\xi, k) = \emptyset$ for $k \neq j$ then

$$\Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) = \Omega_{m,m}^{\mathbb{Z}_2}(X, \xi - \epsilon_V)$$

and the Pontryagin-Thom maps define an isomorphism of direct sum systems

$$\begin{array}{ccc}
\Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2\text{-free}}} & \omega_m^{\mathbb{Z}_2}(S(\infty) \times X; 0 \times (\xi - \epsilon_V)) \\
\parallel & & \parallel \\
\Omega_m(S(\infty) \times_{\mathbb{Z}_2} X, 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) & \xrightarrow[\cong]{PT} & \omega_m(S(\infty) \times_{\mathbb{Z}_2} X; 0 \times_{\mathbb{Z}_2} (\xi - \epsilon_V)) \\
\delta \uparrow \downarrow \gamma & & \delta \uparrow \downarrow \gamma \\
\Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2}} & \omega_m^{\mathbb{Z}_2}(X; \xi - \epsilon_V) \\
\sigma \uparrow \downarrow \rho & & \sigma \uparrow \downarrow \rho \\
\Omega_m(F_-(\xi, V, j), p(j)^*((\xi, j)_+ - \epsilon_{\mathbb{R}^j})) & \xrightarrow[\cong]{PT} & \omega_m(X(\xi, j); (\xi, j)_+ - \epsilon_{\mathbb{R}^j})
\end{array}$$

An m -dimensional V -restricted \mathbb{Z}_2 -equivariant normal map

$$(f, b) : (M, \nu_M - \epsilon_{V \oplus W}) \rightarrow (X, \xi - \epsilon_V)$$

is a disjoint union $(f, b) = (f_1, b_1) \sqcup (f_2, b_2)$ of \mathbb{Z}_2 -equivariant normal maps

$$\begin{aligned} (f_1, b_1) &: (M_1, \nu_{M_1} - \epsilon_{V \oplus W}) \rightarrow (X \setminus X(\xi, j), (\xi - \epsilon_V)|) , \\ (f_2, b_2) &: (M_2, \nu_{M_2} - \epsilon_{V \oplus W}) \rightarrow (X(\xi, j), (\xi, j)_+) \end{aligned}$$

with a free \mathbb{Z}_2 -action on $M_1 = M \setminus M^{\mathbb{Z}_2}$ and a trivial \mathbb{Z}_2 -action on $M_2 = M^{\mathbb{Z}_2}$, and the isomorphism

$$(\delta \rho) : \Omega_m^{\mathbb{Z}_2}(X, \xi - \epsilon_V) \xrightarrow{\cong} \Omega_m^{\mathbb{Z}_2\text{-free}}(X, \xi - \epsilon_V) \oplus \Omega_m(X(\xi, j), (\xi, j)_+ - \epsilon_{\mathbb{R}^j})$$

is given by $(f, b) \mapsto ((f_1, b_1), (f_2, b_2))$.

□

Definition 4.102. Let $V = \mathbb{R}^j \oplus L\mathbb{R}^{i-j}$ be a finite-dimensional inner product \mathbb{Z}_2 -space.

(i) An m -dimensional \mathbb{Z}_2 -manifold M is V -framed if it is V -restricted and the normal \mathbb{Z}_2 -equivariant bundle of an embedding $M \subset V \oplus \mathbb{R}^{m+n}$ is trivialized,

$$\nu_M \cong \epsilon_{V \oplus \mathbb{R}^n} ,$$

so that the \mathbb{Z}_2 -equivariant Pontryagin-Thom Umkehr map

$$\alpha_M : (V \oplus \mathbb{R}^{m+n})^\infty = \Sigma^{m+n} V^\infty \rightarrow T(\nu_M) = \Sigma^n V^\infty \wedge M^+$$

represents an element

$$\alpha_M \in \{\Sigma^{m+n} V^\infty; \Sigma^n V^\infty \wedge M^+\}_{\mathbb{Z}_2} = \omega_m^{\mathbb{Z}_2}(M) .$$

(ii) The m -dimensional framed \mathbb{Z}_2 -equivariant bordism group of a \mathbb{Z}_2 -space X

$$\Omega_m^{\mathbb{Z}_2\text{-fr}}(X) = \Omega_m^{\mathbb{Z}_2}(X, 0) = \Omega_{m,m}^{\mathbb{Z}_2}(X, 0) \quad (\xi = \epsilon_V)$$

is the bordism group of m -dimensional V -framed \mathbb{Z}_2 -manifolds M with a \mathbb{Z}_2 -equivariant map $f : M \rightarrow X$, with the \mathbb{Z}_2 -equivariant framed Pontryagin-Thom map

$$PT^{\mathbb{Z}_2} : \Omega_m^{\mathbb{Z}_2\text{-fr}}(X) \rightarrow \omega_m^{\mathbb{Z}_2}(X) ; ((f, b) : M \rightarrow X) \mapsto f_* \alpha_M .$$

□

Proposition 4.103. The Pontryagin-Thom maps define isomorphisms of direct sum systems

$$\begin{array}{ccc}
\Omega_m^{\mathbb{Z}_2\text{-free-fr}, V}(X) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2\text{-free}}} & \omega_m^{\mathbb{Z}_2}(S(\infty) \times X) \\
\parallel & & \parallel \\
\Omega_m^{\text{fr}}(S(\infty) \times_{\mathbb{Z}_2} X) & \xrightarrow[\cong]{PT} & \omega_m(S(\infty) \times_{\mathbb{Z}_2} X) \\
\delta \uparrow \downarrow \gamma & & \delta \uparrow \downarrow \gamma \\
\Omega_m^{\mathbb{Z}_2\text{-fr}}(X) & \xrightarrow[\cong]{PT^{\mathbb{Z}_2}} & \omega_m^{\mathbb{Z}_2}(X) \\
\sigma \uparrow \downarrow \rho & & \sigma \uparrow \downarrow \rho \\
\Omega_m^{\text{fr}}(X^{\mathbb{Z}_2}) & \xrightarrow[\cong]{PT} & \omega_m(X^{\mathbb{Z}_2})
\end{array}$$

□

Example 4.104. The Pontryagin-Thom construction identifies $\omega_m = \{S^m; S^0\}$ with the cobordism group Ω_m^{fr} of m -dimensional framed manifolds, with

$$\omega_0 = \Omega_0^{\text{fr}} = A(*) = \mathbb{Z}$$

the Burnside ring of finite sets. Similarly, for any finite-dimensional inner product space V the \mathbb{Z}_2 -equivariant Pontryagin-Thom construction identifies $\omega_{m,0} = \{S^m; S^0\}_{\mathbb{Z}_2}$ with the cobordism group $\Omega_m^{\mathbb{Z}_2\text{-fr}, V}$ of m -dimensional V -framed \mathbb{Z}_2 -manifolds $M \subset V \oplus \mathbb{R}^{m+n}$, with

$$\omega_{0,0} = \Omega_0^{\mathbb{Z}_2\text{-fr}, V} = A(\mathbb{Z}_2) = \mathbb{Z}[\mathbb{Z}_2]$$

the Burnside ring $A(\mathbb{Z}_2)$ of finite \mathbb{Z}_2 -sets, with an isomorphism

$$\omega_{0,0} \xrightarrow{\cong} A(\mathbb{Z}_2) = \mathbb{Z}[T]/(1 - T^2) ; S \mapsto \frac{|S| + |S^{\mathbb{Z}_2}|}{2} + \frac{|S| - |S^{\mathbb{Z}_2}|}{2} T .$$

The maps ρ, σ in the direct sum system

$$\omega_0(P(\infty)) = \mathbb{Z} \begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \omega_{0,0} = A(\mathbb{Z}_2) \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \omega_0 = \mathbb{Z}$$

are defined by

$$\begin{aligned}
\rho : \omega_{0,0} = A(\mathbb{Z}_2) &\rightarrow \omega_0 = \mathbb{Z} ; S \mapsto |S^{\mathbb{Z}_2}| , \\
\sigma : \omega_0 = \mathbb{Z} &\rightarrow \omega_{0,0} = A(\mathbb{Z}_2) ; n \mapsto \text{sign}(n)\{1, 2, \dots, |n|\}
\end{aligned}$$

corresponding to

$$\begin{aligned} A(\mathbb{Z}_2) &= \mathbb{Z}[T]/(1 - T^2) \rightarrow \omega_0 = \mathbb{Z} ; a + bT \mapsto a - b , \\ \sigma : \omega_0 = \mathbb{Z} &\rightarrow A(\mathbb{Z}_2) ; n \mapsto n . \end{aligned}$$

An element in $\omega_0(P(\infty)) = \ker(\rho)$ is represented by a finite free \mathbb{Z}_2 -set S , with an isomorphism

$$\ker(\rho) \xrightarrow{\cong} \mathbb{Z} ; S \mapsto \frac{|S|}{2}$$

and injection

$$s_{L\mathbb{R}} : \ker(\rho) \rightarrow A(\mathbb{Z}_2) ; S \mapsto \frac{|S|}{2}(1 + T) .$$

□

Remark 4.105. For any \mathbb{Z}_2 -space X let $\Omega_m^{\mathbb{Z}_2}(X)$ be the bordism group of m -dimensional \mathbb{Z}_2 -manifolds M with a \mathbb{Z}_2 -equivariant map $f : M \rightarrow X$, as in Definition 4.96 (v). Since for every m -dimensional manifold M there exist a finite-dimensional inner product \mathbb{Z}_2 -space V and a \mathbb{Z}_2 -equivariant embedding $M \subset V \oplus \mathbb{R}^m$ with a V -restricted normal bundle $\nu_M : M \rightarrow bo^{\mathbb{Z}_2}(V)$ it is possible to identify

$$\Omega_m^{\mathbb{Z}_2}(X) = \varinjlim_V \Omega_m^{\mathbb{Z}_2}(X \times bo^{\mathbb{Z}_2}(V), 0 \times (\Xi(V) - \epsilon_V)) .$$

□

Proposition 4.106. (Costenoble [10, p.156]). (i) *The \mathbb{Z}_2 -equivariant Pontryagin-Thom maps*

$$PT^{\mathbb{Z}_2} : \Omega_m^{\mathbb{Z}_2}(X) \rightarrow mo_m(X) = \varinjlim_V \omega_m(X \times bo^{\mathbb{Z}_2}(V), 0 \times \Xi(V) - \epsilon_V)$$

are isomorphisms, for any reasonable \mathbb{Z}_2 -space X .

(ii) *The MO -homology groups are such that*

$$\begin{aligned} MO_*^{\mathbb{Z}_2}(X) &= \varinjlim_k mo_{*+k}^{\mathbb{Z}_2}(X \times D(L\mathbb{R}^k), X \times S(L\mathbb{R}^k)) \\ &= \varinjlim_k \widetilde{mo}_{*+k}^{\mathbb{Z}_2}(X^+ \wedge (L\mathbb{R}^k)^\infty) \end{aligned}$$

for any reasonable \mathbb{Z}_2 -space X .

□

Remark 4.107. See Wasserman [69] for an explicit \mathbb{Z}_2 -equivariant transversality theorem, which given that for any \mathbb{Z}_2 -equivariant \mathbb{R}^i -bundle ξ over a

\mathbb{Z}_2 -space X and n -dimensional inner product \mathbb{Z}_2 -space V with sufficiently large $\dim(V_+)$, $\dim(V_-)$ the group $[V^\infty, T(\xi)]_{\mathbb{Z}_2}$ is isomorphic to the bordism group of \mathbb{Z}_2 -equivariant normal maps $(f, b) : (M, \nu_M) \rightarrow (X, \xi)$ with $M \subset V$ an m -dimensional \mathbb{Z}_2 -equivariant submanifold with $m = n - i$, $f : M \rightarrow X$ a \mathbb{Z}_2 -equivariant map and

$$b : \nu_M = \nu_{M \subset V} \rightarrow \xi$$

a \mathbb{Z}_2 -equivariant bundle map over f .

□

Chapter 5

The geometric Hopf invariant

We shall now construct the geometric Hopf invariant of a stable map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$.

5.1 The Q -groups

In the first instance we recall the definition of the various \mathbb{Z}_2 -hypercohomology Q -groups required for the quadratic construction.

Let A be a ring with an involution

$$A \rightarrow A ; a \mapsto \bar{a} ,$$

a function such that

$$\overline{a+b} = \bar{a} + \bar{b} , \overline{ab} = \bar{b}\bar{a} , \bar{\bar{a}} = a , \bar{1} = 1 \in A .$$

The two main examples are:

1. A commutative ring A with the identity involution

$$A \rightarrow A ; a \mapsto \bar{a} = a .$$

2. A group ring $A = \mathbb{Z}[\pi]$ with the involution

$$\mathbb{Z}[\pi] \rightarrow \mathbb{Z}[\pi] ; \sum_{g \in \pi} n_g g \mapsto \sum_{g \in \pi} n_g w(g) g^{-1}$$

for a group morphism $w : \pi \rightarrow \mathbb{Z}_2 = \{\pm 1\}$.

The involution on A can be used to define the dual of a (left) A -module M to be the A -module

$$M^* = \text{Hom}_A(M, A), \quad A \times M^* \rightarrow M^*; (f, a) \mapsto (x \mapsto f(x)\bar{a}),$$

with a duality morphism

$$\text{Hom}_A(M, N) \rightarrow \text{Hom}_A(N^*, M^*); f \mapsto (f^* : g \mapsto gf).$$

The involution can also be used to define a right A -module structure on the additive group of an A -module M

$$M \times A \rightarrow M; (x, a) \mapsto \bar{a}x.$$

For A -modules M, N there is defined a \mathbb{Z} -module

$$M \otimes_A N = M \otimes_{\mathbb{Z}} N / \{ax \otimes y - x \otimes \bar{a}y \mid a \in A, x \in M, y \in N\}$$

and the natural \mathbb{Z} -module morphism

$$M \otimes_A N \rightarrow \text{Hom}_A(M^*, N); x \otimes y \mapsto (f \mapsto f(x)y)$$

is an isomorphism for f.g. projective M . In particular, for $N = A$ and a f.g. projective A -module M there is defined a natural isomorphism

$$M \rightarrow M^{**}; x \mapsto (f \mapsto f(x)).$$

Given A -module chain complexes C, D let $C \otimes_A D, \text{Hom}_A(C, D)$ be the \mathbb{Z} -module chain complexes with

$$\begin{aligned} (C \otimes_A D)_n &= \sum_{p+q=n} C_p \otimes_A D_q, \quad d(x \otimes y) = x \otimes d_D(y) + (-)^q d_C(x) \otimes y, \\ \text{Hom}_A(C, D)_n &= \sum_{q-p=n} \text{Hom}_A(C_p, D_q), \quad d(f) = d_D f + (-)^q f d_C. \end{aligned}$$

A cycle $f \in \text{Hom}_A(C, D)_0$ is a chain map $f : C \rightarrow D$, and $H_0(\text{Hom}_A(C, D))$ is the abelian group of chain homotopy classes of chain maps $f : C \rightarrow D$.

The *suspension* of an A -module chain complex C is the A -module chain complex SC with

$$d_{SC} = d_C : (SC)_r = C_{r-1} \rightarrow (SC)_{r-1} = C_{r-2}.$$

Let C^{-*} be the A -module chain complex with

$$(C^{-*})_r = C^{-r} = \text{Hom}_A(C_{-r}, A), \quad d_{C^{-*}} = (d_C)^*.$$

The natural \mathbb{Z} -module chain map

$$C \otimes_A D \rightarrow \text{Hom}_A(C^{-*}, D) ; x \otimes y \mapsto (f \mapsto \overline{f(x)}y)$$

is an isomorphism if C is a bounded f.g. projective A -module chain complex, in which case a homology class $f \in H_n(C \otimes_A D) = H_0(\text{Hom}_A(C^{n-*}, D))$ is a chain homotopy class of A -module chain maps $f : C^{n-*} \rightarrow D$, with

$$d_{C^{n-*}} = (-)^r d_C^* : (C^{n-*})_r = C^{n-r} \rightarrow (C^{n-*})_{r-1} = C^{n-r+1} .$$

For an A -module chain complex C let the generator $T \in \mathbb{Z}_2$ act on the \mathbb{Z} -module chain complex $C \otimes_A C$ by the signed transposition

$$T : C_p \otimes_A C_q \rightarrow C_q \otimes_A C_p ; x \otimes y \mapsto (-)^{pq} y \otimes x ,$$

so that $C \otimes_A C$ is a $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complex. If C is a bounded f.g. projective A -module chain complex transposition corresponds to duality under the natural isomorphism $C \otimes_A C \cong \text{Hom}_A(C^{-*}, C)$, with

$$T : \text{Hom}_A(C_p^*, C_q) \rightarrow \text{Hom}_A(C_q^*, C_p) ; f \mapsto (-)^{pq} f^* .$$

Definition 5.1. For $-\infty \leq i \leq j \leq \infty$ let $W[i, j]$ be the $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complex with

$$W[i, j]_r = \begin{cases} \mathbb{Z}[\mathbb{Z}_2] & \text{if } i \leq r \leq j \\ 0 & \text{otherwise} \end{cases} , d = 1 + (-)^r T : W[i, j]_r \rightarrow W[i, j]_{r-1} .$$

(i) The $[i, j]$ -symmetric Q -groups of C are

$$Q_{[i, j]}^n(C) = H_n(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[i, j], C \otimes_A C)) .$$

An element $\phi \in Q_{[i, j]}^n(C)$ is an equivalence class of collections

$$\phi = \{\phi_s \in (C \otimes_A C)_{n+s} \mid i \leq s \leq j\}$$

satisfying

$$\begin{aligned} (d \otimes 1)\phi_s + (-)^r \phi_s(1 \otimes d) + (-)^{n+s-1}(\phi_{s-1} + (-)^s T \phi_{s-1}) &= 0 \\ \in (C \otimes_A C)_{n+s-1} &= \sum_r C_{n-r+s-1} \otimes_A C_r \quad (\phi_{i-1} = 0) . \end{aligned}$$

(ii) The $[i, j]$ -quadratic Q -groups of C are

$$Q_n^{[i, j]}(C) = H_n(W[i, j] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C)) .$$

An element $\psi \in Q_n^{[i, j]}(C)$ is an equivalence class of collections

$$\psi = \{\psi_s \in (C \otimes_A C)_{n-s} \mid i \leq s \leq j\}$$

satisfying

$$\begin{aligned} (d \otimes 1)\psi_s + (-)^r \psi_s(1 \otimes d) + (-)^{n-s-1}(\psi_{s+1} + (-)^s T\psi_{s+1}) &= 0 \\ \in (C \otimes_A C)_{n-s-1} &= \sum_r C_{n-r-s-1} \otimes_A C_r \quad (\psi_{j+1} = 0) . \end{aligned}$$

□

Proposition 5.2. ([49, 1.1]) *Let $-\infty \leq i \leq j \leq k \leq \infty$.*

(i) *The functions*

$$\begin{aligned} Q_{[i,j]}^n(C) &\rightarrow Q_{n-1}^{[-1-j, -1-i]}(C) ; \phi \mapsto \psi , \quad \psi_s = \phi_{-1-s} , \\ Q_{[i,j]}^n(C) &\rightarrow Q_{[i+1,j+1]}^{n+1}(SC) ; \phi \mapsto S\phi , \quad S\phi_s = \phi_{s-1} \end{aligned}$$

are isomorphisms.

(ii) *There are defined long exact sequences of Q -groups*

$$\begin{aligned} \cdots \rightarrow Q_{[j+1,k+1]}^n(C) &\rightarrow Q_{[i,k+1]}^n(C) \rightarrow Q_{[i,j]}^n(C) \rightarrow Q_{[j+1,k+1]}^{n-1}(C) \rightarrow \cdots \\ \cdots \rightarrow Q_n^{[i,j]}(C) &\rightarrow Q_n^{[i,k+1]}(C) \rightarrow Q_n^{[j+1,k+1]}(C) \rightarrow Q_{n-1}^{[i,j]}(C) \rightarrow \cdots \end{aligned}$$

Proof. (i) The functions are already isomorphisms on the chain level.

(ii) Immediate from the short exact sequence of $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes

$$0 \rightarrow W[i, j] \rightarrow W[i, k+1] \rightarrow W[j+1, k+1] \rightarrow 0$$

□

For an inner product space V with $\dim(V) = k$ ($1 \leq k \leq \infty$) give $S(LV)$ the standard \mathbb{Z}_2 -equivariant CW structure with cells

$$e_0 , Te_0 , e_1 , Te_1 , \dots , e_{k-1} , Te_{k-1}$$

and $(k-1)$ -dimensional free $\mathbb{Z}[\mathbb{Z}_2]$ -module cellular chain complex

$$C^{cell}(S(LV)) = W[0, k-1] .$$

Definition 5.3. ([49, pp. 100-101]) (i) Given $\mathbb{Z}[\mathbb{Z}_2]$ -modules M, N define a $\mathbb{Z}[\mathbb{Z}_2]$ -module structure on $\text{Hom}_{\mathbb{Z}}(M, N)$ by

$$T : \text{Hom}_{\mathbb{Z}}(M, N) \rightarrow \text{Hom}_{\mathbb{Z}}(M, N) ; f \mapsto T_N f T_M .$$

(ii) Given an inner product space V and $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes C, D define a V -coefficient \mathbb{Z}_2 -isovariant chain map $f : C \rightarrow D$ to be a cycle

$$f = \{f_s \mid 0 \leq s \leq \dim(V) - 1\} \in \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)), \text{Hom}_{\mathbb{Z}}(C, D))_0 ,$$

with $f_s \in \text{Hom}_{\mathbb{Z}}(C_r, D_{r+s})$ such that

$$\begin{aligned} d_D f_s + (-)^{s-1} f_s d_C + (-)^{s-1} (f_{s-1} + (-)^s T_D f_{s-1} T_C) &= 0 : \\ C_r &\rightarrow D_{r+s-1} \quad (f_{-1} = 0) , \end{aligned}$$

which can be regarded as a $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map $f : C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C \rightarrow D$. Thus $f_0 : C \rightarrow D$ is a \mathbb{Z} -module chain map, $f_1 : f_0 \simeq T_D f_0 T_C$ is a chain homotopy, f_2 is a higher chain homotopy, etc.

(iii) There is a corresponding notion of V -coefficient \mathbb{Z}_2 -isovariant chain homotopy $h : f \simeq g : C \rightarrow D$, and

$$\begin{aligned} H_0(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)), \text{Hom}_{\mathbb{Z}}(C, D))) \\ = H_0(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C, D)) \end{aligned}$$

is the abelian group of V -coefficient \mathbb{Z}_2 -isovariant chain homotopy classes of V -coefficient \mathbb{Z}_2 -isovariant chain maps.

(iv) A V -coefficient \mathbb{Z}_2 -isovariant chain map $f : C \rightarrow D$ induces a \mathbb{Z} -module chain map

$$\begin{aligned} f^{\%} : \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)), C) &\rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)), D) ; \\ g = \{g_s\} &\mapsto (f \otimes g) \Delta \end{aligned}$$

with

$$\Delta : C^{cell}(S(LV)) \rightarrow C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C^{cell}(S(LV)) ; 1_r \mapsto \sum_{s=0}^r 1_s \otimes (T_{r-s})^s$$

the *cellular diagonal chain approximation* $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map.

(v) The *composite* of V -coefficient \mathbb{Z}_2 -isovariant chain maps $f : C \rightarrow D$, $g : D \rightarrow E$ is the V -coefficient \mathbb{Z}_2 -isovariant chain map $gf : C \rightarrow E$ with

$$\begin{aligned} gf &= f^{\%}(g) : C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C \xrightarrow{\Delta \otimes 1} \\ C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C &\xrightarrow{1 \otimes f} C^{cell}(S(LV)) \otimes_{\mathbb{Z}} D \xrightarrow{g} E . \end{aligned}$$

(vi) The *cup product* of V -coefficient \mathbb{Z}_2 -isovariant chain maps $f : C \rightarrow D$, $g : E \rightarrow F$ is the V -coefficient \mathbb{Z}_2 -isovariant chain map with

$$\begin{aligned} f \cup g &= (f \otimes g) \Delta : C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C \otimes_{\mathbb{Z}} E \xrightarrow{\Delta \otimes 1} \\ C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C \otimes_{\mathbb{Z}} E &\xrightarrow{f \otimes g} D \otimes_{\mathbb{Z}} F . \end{aligned}$$

□

Example 5.4. (i) For $V = \mathbb{R}$ a V -coefficient \mathbb{Z}_2 -isovariant chain map $f : C \rightarrow D$ is just a \mathbb{Z} -module chain map $f_0 : C \rightarrow D$.
(ii) For $V = \mathbb{R}^2$ a V -coefficient \mathbb{Z}_2 -isovariant chain map $f : C \rightarrow D$ is a \mathbb{Z} -module chain map $f_0 : C \rightarrow D$ together with a \mathbb{Z} -module chain homotopy $f_1 : f_0 \simeq T_D f_0 T_C : C \rightarrow D$.

□

Proposition 5.5. *Let $f : C \rightarrow D$ is a V -coefficient \mathbb{Z}_2 -isovariant chain map of $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes which are bounded below. If $f_0 : C \rightarrow D$ induces isomorphisms $(f_0)_* : H_*(C) \cong H_*(D)$ the \mathbb{Z} -module chain map $f^\%$ induces isomorphisms*

$$(f^\%)_* : H_*(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)), C)) \cong H_*(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)), D)) .$$

Proof. Standard homological algebra.

□

Let $\widehat{S}(LV)$ be the suspension \mathbb{Z}_2 -spectrum with

$$\widehat{S}(LV)_j = \Sigma^{j-k} S(LV \oplus L\mathbb{R}^j)^+ \quad (j \geq k)$$

and $\mathbb{Z}[\mathbb{Z}_2]$ -module cellular chain complex

$$\dot{C}^{cell}(\widehat{S}(LV)) = C^{cell}(S(LV \oplus LV))_{*+k} = W[-k, k-1] .$$

Definition 5.6. Let C be an A -module chain complex, and let V be an inner product space with $\dim(V) = k$ ($1 \leq k \leq \infty$).

(i) The V -coefficient symmetric Q -groups of C are

$$Q_V^n(C) = H_n(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(S(LV)), C \otimes_A C)) = Q_{[0, k-1]}^n(C) .$$

(ii) The V -coefficient quadratic Q -groups of C are

$$Q_n^V(C) = H_n(C^{cell}(S(LV)) \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C)) = Q_n^{[0, k-1]}(C) .$$

(iii) The V -coefficient hyperquadratic Q -groups of C are

$$\widehat{Q}_V^n(C) = H_n(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(C^{cell}(\widehat{S(LV)}), C \otimes_A C)) = Q_{[-k, k-1]}^n(C) .$$

□

Proposition 5.7. (i) An A -module chain map $f : C \rightarrow D$ determines a $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$f \otimes f : C \otimes_A C \rightarrow D \otimes_A D$$

which can be regarded as a V -coefficient \mathbb{Z}_2 -isovariant chain map with

$$(f \otimes f)_s = \begin{cases} f \otimes f & \text{if } s = 0 \\ 0 & \text{if } s \geq 1 \end{cases} : (C \otimes_A C)_r \rightarrow (D \otimes_A D)_{r+s} .$$

(ii) An A -module chain homotopy $g : f \simeq f' : C \rightarrow D$ determines a V -coefficient \mathbb{Z}_2 -isovariant chain homotopy

$$g \otimes g : f \otimes f \simeq f' \otimes f' : C \otimes_A C \rightarrow D \otimes_A D$$

with

$$(g \otimes g)_s = \begin{cases} f \otimes g + (-)^q g \otimes f' & \text{if } s = 0 \\ (-)^q g \otimes g & \text{if } s = 1 \\ 0 & \text{if } s \geq 2 \end{cases} : (C \otimes_A C)_r \rightarrow (D \otimes_A D)_{r+s+1} = \sum_q D_{-q+r+s+1} \otimes_A D_q .$$

(iii) An A -module chain map $f : C \rightarrow D$ induces morphisms in the Q -groups

$$\begin{aligned} f^\% & : Q_V^n(C) \rightarrow Q_V^n(D) , \\ f_\% & : Q_n^V(C) \rightarrow Q_n^V(D) , \\ \widehat{f}^\% & : \widehat{Q}_V^n(C) \rightarrow \widehat{Q}_V^n(D) \end{aligned}$$

which depend only on the chain homotopy class of f , and are isomorphisms if f is a chain equivalence.

(iv) The chain level cup products

$$\begin{aligned} & \text{Hom}_{\mathbb{Z}}(C^{\text{cell}}S(LV), C \otimes_A C) \otimes_{\mathbb{Z}} \text{Hom}_{\mathbb{Z}}(C^{\text{cell}}S(LV), D \otimes_B D) \\ & \rightarrow \text{Hom}_{\mathbb{Z}}(C^{\text{cell}}S(LV), (C \otimes_{\mathbb{Z}} D) \otimes_{A \otimes_{\mathbb{Z}} B} (C \otimes_{\mathbb{Z}} D)) ; \\ & \phi \otimes \theta \mapsto \phi \cup \theta = (\phi \otimes \theta) \Delta \end{aligned}$$

induce cup products in the V -coefficient symmetric Q -groups

$$\cup : Q_V^m(C) \otimes_{\mathbb{Z}} Q_V^n(D) \rightarrow Q_V^{m+n}(C \otimes_{\mathbb{Z}} D) ; \phi \otimes \theta \mapsto \phi \cup \theta .$$

Similarly for products involving the V -coefficient quadratic and hyperquadratic Q -groups

$$\begin{aligned} Q_V^m(C) \otimes_{\mathbb{Z}} Q_n^V(D) & \rightarrow Q_{m+n}^V(C \otimes_{\mathbb{Z}} D) , \\ \widehat{Q}_m^V(C) \otimes_{\mathbb{Z}} \widehat{Q}_n^V(D) & \rightarrow \widehat{Q}_V^{m+n}(C \otimes_{\mathbb{Z}} D) . \end{aligned}$$

Proof. See [49, pp. 100-101, §8].

□

Definition 5.8. Let $\sigma \in \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, 1], S\mathbb{Z} \otimes_A S\mathbb{Z})_1$ be the cycle defined by

$$\sigma_1 = 1 : (S\mathbb{Z})^1 = \mathbb{Z} \rightarrow (S\mathbb{Z})_1 = \mathbb{Z} .$$

The *suspension* chain map is defined for any A -module chain complex C to be the evaluation of the cup product on σ

$$S = \sigma \cup - :$$

$$\begin{aligned} \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, k], C \otimes_A C) &\rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, k+1], SC \otimes_A SC)_{*+1} ; \phi \mapsto S\phi , \\ S\phi_s = \phi_{s-1} &\in (SC \otimes_A SC)_{n+s+1} = (C \otimes_A C)_{n+s-1} \quad (0 \leq s \leq k+1, \phi_{-1} = 0) \end{aligned}$$

inducing suspension maps in the symmetric Q -groups.

$$S : Q_V^n(C) \rightarrow Q_{V \oplus \mathbb{R}}^{n+1}(SC) .$$

Similarly, product with σ induces suspension maps in the quadratic and hyperquadratic Q -groups

$$\begin{aligned} S : Q_n^V(C) &\rightarrow Q_{n+1}^{V \oplus \mathbb{R}}(SC) , \\ S : \hat{Q}_V^n(C) &\rightarrow \hat{Q}_{V \oplus \mathbb{R}}^{n+1}(SC) . \end{aligned}$$

□

Proposition 5.9. ([49, 1.3])

(i) For any inner product spaces U, V there is defined a long exact sequence of Q -groups

$$\dots \rightarrow Q_n^V(C) \xrightarrow{1+T} Q_U^n(C) \xrightarrow{S^{\dim(V)}} Q_{U \oplus V}^{n+\dim(V)}(S^{\dim(V)}C) \rightarrow Q_{n-1}^V(C) \rightarrow \dots$$

with $S^{\dim(V)}C = C_{*-\dim(V)}$ and

$$\begin{aligned} 1+T : Q_n^U(C) &\rightarrow Q_V^n(C) ; \psi \mapsto (1+T)\psi , \\ (1+T)\psi_s &= \begin{cases} \psi_0 + T\psi_0 & \text{if } s = 0 \\ 0 & \text{if } s \geq 1 , \end{cases} \\ Q_{U \oplus V}^{n+\dim(V)}(S^{\dim(V)}C) &\rightarrow Q_{n-1}^V(C) ; \\ \phi \mapsto \psi , \psi_s &= \phi_{\dim(V)-s-1} \quad (0 \leq s \leq \dim(V)-1) \end{aligned}$$

(ii) If $U = \{0\}$ then $Q_U^*(C) = 0$ and the sequence in (i) gives the isomorphisms

$$Q_V^{n+\dim(V)}(S^{\dim(V)}C) \rightarrow Q_{n-1}^V(C) ; \phi \mapsto \psi , \psi_s = \phi_{\dim(V)-s-1}$$

of 5.2 (i), which are induced by the chain level \mathbb{Z}_2 -equivariant S -duality isomorphism of Example 4.89 (ii)

$$\begin{aligned} \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(SC^{\text{cell}}(S(LV)), \dot{C}^{\text{cell}}(LV^\infty) \otimes_{\mathbb{Z}} (C \otimes_A C)) \\ \xrightarrow{\cong} C^{\text{cell}}(S(LV)) \otimes_{\mathbb{Z}} (C \otimes_A C) . \end{aligned}$$

(iii) If $U = V$ there are defined isomorphisms

$$\begin{aligned} Q_{V \oplus V}^{n+\dim(V)}(S^{\dim(V)}C) &\rightarrow \widehat{Q}_V^n(C) ; \\ \phi &\mapsto \widehat{\phi} , \widehat{\phi}_s = \phi_{s+\dim(V)} \quad (-\dim(V) \leq s \leq \dim(V) - 1) \end{aligned}$$

and the sequence in (i) is

$$\dots \longrightarrow Q_n^V(C) \xrightarrow{1+T} Q_V^n(C) \xrightarrow{J} \widehat{Q}_V^n(C) \xrightarrow{H} Q_{n-1}^V(C) \longrightarrow \dots$$

with

$$\begin{aligned} J : Q_V^n(C) &\rightarrow \widehat{Q}_V^n(C) ; \phi \mapsto J\phi , J\phi_s = \begin{cases} \phi_s & \text{if } 0 \leq s \leq \dim(V) - 1 \\ 0 & \text{if } -\dim(V) \leq s \leq -1 \end{cases} , \\ H : \widehat{Q}_V^n(C) &\rightarrow Q_{n-1}^V(C) ; \widehat{\phi} \mapsto H\widehat{\phi} , H\widehat{\phi}_s = \widehat{\phi}_{-s-1} . \end{aligned}$$

(iv) The suspension maps in the hyperquadratic Q -groups

$$S : \widehat{Q}_V^n(C) \rightarrow \widehat{Q}_{V \oplus \mathbb{R}}^{n+1}(SC)$$

are isomorphisms, and there is defined a commutative braid of exact sequences of Q -groups

$$\begin{array}{ccccc} & & H & & \\ & \widehat{Q}_V^{n+1}(C) & \searrow & \widehat{Q}_n^{V \oplus \mathbb{R}}(SC) & \searrow & H_{n-1}(C \otimes_A C) \\ & & H & S & 1+T & \\ & Q_n^V(C) & & Q_{V \oplus \mathbb{R}}^n(SC) & & \\ & \nearrow & 1+T & S & \nearrow & \\ H_n(C \otimes_A C) & & Q_V^n(C) & & \widehat{Q}_V^n(C) & \\ & & J & & & \end{array}$$

Proof. (i) This is a special case of Proposition 5.2

$$\begin{aligned} \cdots \rightarrow Q_n^V(C) &= Q_{[-\dim(V), -1]}^{n+1}(C) \rightarrow Q_U^n(C) = Q_{[0, \dim(U)-1]}^n(C) \rightarrow \\ Q_{U \oplus V}^{n+\dim(V)}(C) &= Q_{[-\dim(V), \dim(U)-1]}^n(C) \rightarrow Q_{n-1}^V(C) = Q_{[-\dim(V), -1]}^n(C) \rightarrow \cdots \end{aligned}$$

Specifically, let $\dim(U) = j$, $\dim(V) = k$, and note that the cofibration sequence of pointed CW - \mathbb{Z}_2 -complexes

$$S(LV)^+ \rightarrow S(LU \oplus LV)^+ \rightarrow S(LU)^+ \wedge LV^\infty$$

induces the short exact sequence of cellular free $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes

$$\begin{aligned} 0 \rightarrow C^{cell}(S(LV)) &= W[0, k-1] \rightarrow C^{cell}(S(LU \oplus LV)) = W[0, j+k-1] \\ &\rightarrow \dot{C}^{cell}(S(LU)^+ \wedge LV^\infty) = W[k, j+k-1] \rightarrow 0 \end{aligned}$$

inducing a long exact sequence

$$\begin{aligned} \cdots \rightarrow Q_n^V(C) &= Q_n^{[0, k-1]}(C) = Q_{[0, k-1]}^{n+k+1}(S^k C) \\ &\rightarrow Q_U^n(C) = Q_{[0, j-1]}^n(C) = Q_{[k, j+k-1]}^{n+k}(S^k C) \\ &\rightarrow Q_{U \oplus V}^{n+\dim(V)}(S^{\dim(V)} C) = Q_{[0, j+k-1]}^{n+k}(S^k C) \\ &\rightarrow Q_{n-1}^V(C) = Q_{[0, k-1]}^{n+k}(S^k C) \rightarrow \cdots \end{aligned}$$

(ii) Immediate from (i).

(iii) Immediate from (i), noting that the long exact sequence is induced from the short exact sequence of $\mathbb{Z}[\mathbb{Z}_2]$ -module cellular chain complexes

$$0 \rightarrow C^{cell}(S(LV))^{-*-1} \rightarrow \dot{C}^{cell}(\hat{S}(LV)) \rightarrow C^{cell}(S(LV)) \rightarrow 0.$$

(iv) Immediate from (ii) and (iii).

□

Terminology 5.10 ([49, §1])

The infinite-dimensional inner product space

$$V = \mathbb{R}(\infty) = \varinjlim_k \mathbb{R}^k$$

has unit sphere

$$S(LV) = S(\infty) = E\mathbb{Z}_2$$

a contractible space with a free \mathbb{Z}_2 -action, and

$$C^{cell}(S(\infty)) = W[0, \infty] = W :$$

$$\cdots \rightarrow W_2 = \mathbb{Z}[\mathbb{Z}_2] \xrightarrow{1+T} W_1 = \mathbb{Z}[\mathbb{Z}_2] \xrightarrow{1-T} W_0 = \mathbb{Z}[\mathbb{Z}_2]$$

is the standard free $\mathbb{Z}[\mathbb{Z}_2]$ -module resolution of \mathbb{Z} .

(i) The *symmetric Q -groups* of an A -module chain complex C are

$$Q^n(C) = Q_{\mathbb{R}(\infty)}^n(C) = Q_{[0, \infty]}^n(C) .$$

(ii) The *quadratic Q -groups* of an A -module chain complex C are

$$Q_n(C) = Q_n^{\mathbb{R}(\infty)}(C) = Q_n^{[0, \infty]}(C) .$$

(ii) The *hyperquadratic Q -groups* of an A -module chain complex C are

$$\hat{Q}^n(C) = \hat{Q}_{\mathbb{R}(\infty)}^n(C) = Q_{[-\infty, \infty]}^n(C) .$$

□

Proposition 5.11. ([49, 1.2]) *Let C be an A -module chain complex.*

(i) *The symmetric and quadratic Q -groups are related by a long exact sequence*

$$\cdots \longrightarrow Q_n^{[0, k-1]}(C) \xrightarrow{1+T} Q^n(C) \xrightarrow{S^k} Q^{n+k}(S^k C) \xrightarrow{H} Q_{n-1}^{[0, k-1]}(C) \longrightarrow \cdots$$

for any $k \geq 0$.

(ii) *The symmetric, quadratic and hyperquadratic Q -groups are related by a long exact sequence*

$$\cdots \longrightarrow Q_n(C) \xrightarrow{1+T} Q^n(C) \xrightarrow{J} \hat{Q}^n(C) \xrightarrow{H} Q_{n-1}(C) \longrightarrow \cdots$$

and

$$\hat{Q}^n(C) = \varinjlim_k Q^{n+k}(S^k C)$$

is the direct limit of the suspension maps

$$Q^n(C) \xrightarrow{S} Q^{n+1}(SC) \xrightarrow{S} Q^{n+2}(S^2 C) \longrightarrow \cdots .$$

Proof. (i) This is the special case $U = \mathbb{R}(\infty)$, $V = \mathbb{R}^k$ of 5.9.

(ii) This is the special case $U = V = \mathbb{R}(\infty)$ of 5.9.

□

Definition 5.12. Let C be an A -module chain complex, and $1 \leq k \leq \infty$.

(i) A *symmetric k -class* is an element $\phi \in Q_{[0, k-1]}^n(C)$. A symmetric ∞ -class

is an element $\phi \in Q_{[0,\infty]}^n(C) = Q^n(C)$, and this is called a *symmetric class*. A symmetric class ϕ determines a symmetric k -class $\phi[0, k-1]$ for every k , via the morphism


$$Q^n(C) \rightarrow Q_{[0,k-1]}^n(C); \phi = \{\phi_s \mid 0 \leq s < \infty\} \mapsto \phi[0, k-1] = \{\phi_s \mid 0 \leq s < k\}.$$

(ii) A *quadratic k -class* is an element $\psi \in Q_n^{[0,k-1]}(C)$. For $k = \infty$ this is an element $\psi \in Q_n(C)$, and this is called a *quadratic class*.

(iii) A *quadratic refinement* of a symmetric k -class ϕ is a quadratic k -class ψ such that $(1+T)\psi = \phi$.

□

Proposition 5.13. (i) For any A -module chain complex C and $0 \leq k \leq \infty$ there is defined a commutative braid of exact sequences



with

$$\begin{aligned} Q_{[-k,k-1]}^n(C) &= Q_{[-k,k-1]}^{n+k}(S^k C), \\ Q_{[-k,\infty]}^n(C) &= Q^{n+k}(S^k C), \quad Q_{[k,\infty]}^n(C) = Q_{[0,k-1]}^{n+k}(S^k C). \end{aligned}$$

(ii) The following conditions on symmetric class $\phi \in Q^n(C)$ are equivalent:

- (a) there exists a quadratic refinement $\psi \in Q_n^{[0,k-1]}(C)$
- (b) $S^k \phi = 0 \in Q^{n+k}(S^k C)$,
- (c) $S^k \phi[0, k-1] = 0 \in Q_{[0,k-1]}^{n+k}(S^k C)$,

In particular, a chain level solution of $S^k \phi = 0$ gives a particular ψ , in which case

$$(1+T)\psi = \phi$$

$$\in \text{im}(1+T : Q_n^{[0,k-1]}(C) \rightarrow Q^n(C)) = \ker(S^k : Q^n(C) \rightarrow Q^{n+k}(S^k C)) .$$

This applies also for $k = \infty$, with $Q_n^{[0,k-1]}(C) = Q_n(C)$, $S^k = J$.

(iii) Let $A = \mathbb{Z}[\pi]$ be a group ring with an involution, so that for any A -modules M, N the diagonal π -action can be used to regard $M \otimes_{\mathbb{Z}} N$ as an A -module with

$$\mathbb{Z} \otimes_A (M \otimes_{\mathbb{Z}} N) = M \otimes_A N .$$

For any A -module chain complexes C, D and $k \geq 1$ there is a natural isomorphism of short exact sequences of \mathbb{Z} -module chain complexes

$$\begin{array}{ccc}
0 & & 0 \\
\downarrow & & \downarrow \\
\text{Hom}_{A[\mathbb{Z}_2]}(W[k, \infty] \otimes_{\mathbb{Z}} S^k C, S^k D \otimes_{\mathbb{Z}} S^k D) & \xrightarrow{\cong} & \text{Hom}_{A[\mathbb{Z}_2]}(W \otimes_{\mathbb{Z}} C, D \otimes_{\mathbb{Z}} D) \\
\downarrow & & \downarrow S^k \\
\text{Hom}_{A[\mathbb{Z}_2]}(W \otimes_{\mathbb{Z}} S^k C, S^k D \otimes_{\mathbb{Z}} S^k D) & \xrightarrow{\cong} & \text{Hom}_{A[\mathbb{Z}_2]}(W \otimes_{\mathbb{Z}} S^k C, S^k D \otimes_{\mathbb{Z}} S^k D) \\
\downarrow & & \downarrow \\
\text{Hom}_{A[\mathbb{Z}_2]}(W[0, k-1] \otimes_{\mathbb{Z}} S^k C, S^k D \otimes_{\mathbb{Z}} S^k D) & \xrightarrow{\cong} & \text{Hom}_{A[\mathbb{Z}_2]}(C, SW[0, k-1] \otimes_{\mathbb{Z}} (D \otimes_{\mathbb{Z}} D)) \\
\downarrow & & \downarrow \\
0 & & 0
\end{array}$$

as well as an isomorphism

$$\text{Hom}_{A[\mathbb{Z}_2]}(SW[0, k-1] \otimes_{\mathbb{Z}} S^k C, S^k D \otimes_{\mathbb{Z}} S^k D) \cong \text{Hom}_{A[\mathbb{Z}_2]}(C, W[0, k-1] \otimes_{\mathbb{Z}} (D \otimes_{\mathbb{Z}} D))$$

Suppose given A -module chain complexes C, C', D, D' with A -module chain maps

$$E_C : S^k C \rightarrow C' , E_D : S^k D \rightarrow D' , F : C \rightarrow D , F' : C' \rightarrow D' ,$$

such that E_C, E_D are chain equivalences with

$$F' E_C = E_D F : C \rightarrow D' .$$

Suppose given also $A[\mathbb{Z}_2]$ -module chain maps

$$\begin{aligned}
\phi_C : W \otimes_{\mathbb{Z}} C &\rightarrow C \otimes_{\mathbb{Z}} C , \phi_{C'} : W \otimes_{\mathbb{Z}} C' \rightarrow C' \otimes_{\mathbb{Z}} C' , \\
\phi_D : W \otimes_{\mathbb{Z}} D &\rightarrow D \otimes_{\mathbb{Z}} D , \phi_{D'} : W \otimes_{\mathbb{Z}} D' \rightarrow D' \otimes_{\mathbb{Z}} D'
\end{aligned}$$

such that

$$(F' \otimes F')\phi_{C'} = \phi_{D'}(1 \otimes F') : W \otimes_{\mathbb{Z}} C' \rightarrow D' \otimes_{\mathbb{Z}} D'$$

and a chain homotopy commutative diagram of $A[\mathbb{Z}_2]$ -module chain complexes and chain maps

$$\begin{array}{ccccc}
 W \otimes_{\mathbb{Z}} S^k C & \xrightarrow{S^k \phi_C} & S^k C \otimes_{\mathbb{Z}} S^k C & & \\
 \downarrow 1 \otimes F & \searrow \delta \phi_C & \swarrow E_C \otimes E_C & & \downarrow F \otimes F \\
 & W \otimes_{\mathbb{Z}} C' & \xrightarrow{\phi_{C'}} & C' \otimes_{\mathbb{Z}} C' & \\
 & \downarrow 1 \otimes F' & & \downarrow F' \otimes F' & \\
 & W \otimes_{\mathbb{Z}} D' & \xrightarrow{\phi_{D'}} & D' \otimes_{\mathbb{Z}} D' & \\
 \downarrow 1 \otimes F & \searrow \delta \phi_D & \swarrow E_D \otimes E_D & & \downarrow F \otimes F \\
 W \otimes_{\mathbb{Z}} S^k D & \xrightarrow{S^k \phi_D} & S^k D \otimes_{\mathbb{Z}} S^k D & &
 \end{array}$$

with chain homotopies

$$\delta \phi_C : (E_C \otimes E_C)(S^k \phi_C) \simeq \phi_{C'}(1 \otimes E_C) : W \otimes_{\mathbb{Z}} S^k C \rightarrow C' \otimes_{\mathbb{Z}} C' ,$$

$$\delta \phi_D : (E_D \otimes E_D)(S^k \phi_D) \simeq \phi_{D'}(1 \otimes E_D) : W \otimes_{\mathbb{Z}} S^k D \rightarrow D' \otimes_{\mathbb{Z}} D' .$$

The data determines an A -module chain map

$$\psi = \delta(F, \phi_C, \phi_D) : C \rightarrow W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (D \otimes_{\mathbb{Z}} D)$$

such that

$$(1 + T)\psi \simeq (F \otimes F)\phi_C - \phi_D F : C \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, D \otimes_{\mathbb{Z}} D) .$$

The induced \mathbb{Z} -module chain maps

$$\phi_C : \mathbb{Z} \otimes_A C \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, C \otimes_A C) ,$$

$$\phi_D : \mathbb{Z} \otimes_A D \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, D \otimes_A D) ,$$

$$\psi = \delta(f, \phi_C, \phi_D) : \mathbb{Z} \otimes_A C \rightarrow W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (D \otimes_A D)$$

are such that

$$(1 + T)\psi \simeq (F \otimes F)\phi_C - \phi_D F : \mathbb{Z} \otimes_A C \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, D \otimes_A D) .$$

Proof. (i) The short exact sequence of \mathbb{Z} -module chain complexes underlying the proof of Proposition 5.11

$$\begin{aligned}
0 &\rightarrow \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, j-1], C \otimes_A C) \\
&\xrightarrow{S^k} \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, j+k-1], S^k C \otimes_A S^k C)_{*+k} \\
&\rightarrow S(W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C)) \rightarrow 0
\end{aligned}$$

gives an identification

$$W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C) = \mathcal{C}(S^k)_{*+1}$$

and a chain equivalence

$$\begin{aligned}
\mathcal{C}(1+T : W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C) &\rightarrow \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, j-1], C \otimes_A C)) \\
&\simeq \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, j+k-1], S^k C \otimes_A S^k C)_{*+k} .
\end{aligned}$$

An n -cycle $\psi \in (W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C))_n$ is thus essentially the same (up to homology) as an n -cycle $\phi \in \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, j-1], C \otimes_A C)_n$ together with an $(n+k+1)$ -chain

$$\delta\phi \in \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, j+k-1], S^k C \otimes_A S^k C)_{n+k+1}$$

such that $S^k\phi = d(\delta\phi)$, so that up to signs

$$\begin{aligned}
\phi_r &= (d \otimes 1 + 1 \otimes d)\delta\phi_{r+k} + \delta\phi_{r+k-1} + T\delta\phi_{r+k-1} \\
&\in (C \otimes_A C)_{n+r} = (S^k C \otimes_A S^k C)_{n+2k+r} \\
&\quad (-k \leq r \leq j-1, \phi_q = 0 \text{ for } q < 0) , \\
\psi_s &= \delta\phi_{k-s-1} \in (C \otimes_A C)_{n-s} = (S^k C \otimes_A S^k C)_{n+2k-s} \quad (0 \leq s \leq k-1) .
\end{aligned}$$

and

$$\begin{aligned}
\phi &= (1+T)\psi \in \operatorname{im}(1+T : Q_n^{[0, k-1]}(C) \rightarrow Q_{[0, j-1]}^n(C)) \\
&= \ker(S^k : Q_{[0, j-1]}^n(C) \rightarrow Q_{[0, j+k-1]}^{n+k}(S^k C)) .
\end{aligned}$$

For $j = k = \infty$ this is just the short exact sequence of \mathbb{Z} -module chain complexes underlying the proof of Proposition 5.11

$$\begin{aligned}
0 &\rightarrow \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, C \otimes_A C) \\
&\xrightarrow{J} \varinjlim_k \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, S^k C \otimes_A S^k C)_{*+k} = \operatorname{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(\widehat{W}, C \otimes_A C) \\
&\xrightarrow{H} S(W \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C)) \rightarrow 0
\end{aligned}$$

which gives an identification

$$W \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C) = \mathcal{C}(J)_{*+1}$$

and a chain equivalence

$$\begin{aligned} \mathcal{C}(1+T : W \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C) &\rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, C \otimes_A C)) \\ &\simeq \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(\widehat{W}, C \otimes_A C) . \end{aligned}$$

An n -cycle $\psi \in (W \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (C \otimes_A C))_n$ is thus essentially the same (up to homology) as an n -cycle $\phi \in \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, C \otimes_A C)_n$ together with an $(n+k+1)$ -chain

$$\delta\phi \in \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, S^k C \otimes_A S^k C)_{n+k+1}$$

such that $S^k\phi = d(\delta\phi)$, and

$$\phi = (1+T)\psi \in \text{im}(1+T : Q_n^{[0, k-1]}(C) \rightarrow Q^n(C)) = \ker(S^k : Q^n(C) \rightarrow \widehat{Q}^n(C)) .$$

(ii) The A -module chain map

$$\begin{aligned} \theta &= (F'E_C \otimes F'E_C)(\phi_C) - (E_D \otimes E_D)(\phi_D)F : \\ &C \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, S^{-k}D' \otimes_{\mathbb{Z}} S^{-k}D') \end{aligned}$$

and the A -module chain null-homotopy

$$(F' \otimes F')\delta\phi_C - \delta\phi_D(1 \otimes F) : S^k\theta \simeq 0 : S^k C \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, D' \otimes_{\mathbb{Z}} D')$$

correspond by (i) to an A -module chain map

$$\psi' : C \rightarrow W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (S^{-k}D' \otimes_{\mathbb{Z}} S^{-k}D')$$

such that $(1+T)\psi' \simeq \theta$. Use any chain homotopy inverse $(E_D)^{-1} : S^{-k}D' \rightarrow D$ of E_D to define an A -module chain map

$$\psi = ((E_D)^{-1} \otimes (E_D)^{-1})\psi' : C \rightarrow W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (D \otimes_{\mathbb{Z}} D)$$

such that $(1+T)\psi \simeq (F \otimes F)\phi_C - \phi_D F$.

□

Remark 5.14. For a bounded f.g. projective A -module chain complex C the forgetful map

$$Q^n(C) \rightarrow H_n(C \otimes_A C) ; \phi \mapsto \phi_0$$

sends a symmetric class $\phi \in Q^n(C)$ to a chain homotopy class

$$\phi_0 \in H_n(C \otimes_A C) = H_0(\text{Hom}_A(C^{-*}, C))$$

of A -module chain maps $\phi_0 : C^{n-*} \rightarrow C$ with

$$d_{C^{n-*}} = (-)^r d_C^* : (C^{n-*})_r = C^{n-r} \rightarrow (C^{n-*})_{r-1} = C^{n-r+1} .$$

□

The symmetric L -groups $L^n(A)$ of Mishchenko [44] are the cobordism groups of n -dimensional symmetric Poincaré complexes (C, ϕ) over A , as defined by an n -dimensional f.g. free A -module chain complex C together with a symmetric class $\phi \in Q^n(C)$ such that $\phi_0 : C^{n-*} \rightarrow C$ is a chain homotopy class of chain equivalences. The quadratic L -groups $L_n(A)$ of Wall [68] were expressed in Ranicki [49] as the cobordism groups of n -dimensional quadratic Poincaré complexes (C, ψ) over A , as defined by an n -dimensional f.g. free A -module chain complex C together with a quadratic class $\psi \in Q_n(C)$ such that $(1 + T)\psi_0 : C^{n-*} \rightarrow C$ is a chain homotopy class of chain equivalences. The surgery obstruction of an n -dimensional normal map $(f, b) : M \rightarrow X$ was expressed in Ranicki [50] as the cobordism class

$$\sigma_*(f, b) = (C, \psi) \in L_n(\mathbb{Z}[\pi_1(X)])$$

with C a $\mathbb{Z}[\pi_1(X)]$ -module chain complex such that

$$H_*(C) = K_*(M) = \ker(\tilde{f}_* : H_*(\tilde{M}) \rightarrow H_*(\tilde{X})) ,$$

with \tilde{X} the universal cover of X and $\tilde{M} = f^*\tilde{X}$ the pullback cover of M . The above chain level mechanism was used in [50] to refine the symmetric structure $\phi \in Q^n(C)$ on the chain complex kernel C of a normal map (f, b) to a quadratic structure $\psi \in Q_n(C)$ using a stable $\pi_1(X)$ -equivariant map $F : \Sigma^k \tilde{X}^+ \rightarrow \Sigma^k \tilde{M}^+$ (k large) S -dual to the induced $\pi_1(X)$ -equivariant map of Thom spaces $T(\tilde{b}) : T(\tilde{\nu}_M) \rightarrow T(\tilde{\nu}_X)$ to provide a chain level solution of $S^k \phi = 0 \in Q^{n+k}(S^k C)$. In the following sections we shall use the geometric Hopf invariant to construct a homotopy theoretic mechanism inducing this chain level mechanism. In the first instance we ignore the fundamental group $\pi_1(X)$, returning to it in Chapter 7.

5.2 The symmetric construction $\phi_V(X)$

As before, let $C(X)$ denote the singular chain complex of a space X , with

$$H_*(C(X)) = H_*(X) , \quad H^*(C(X)) = H^*(X)$$

the singular homology and cohomology groups of X .

Proposition 5.15. (Eilenberg-Zilber)

(i) *For any spaces X, Y there exist inverse \mathbb{Z} -module chain equivalences*

$$\begin{aligned} E(X, Y) &: C(X) \otimes_{\mathbb{Z}} C(Y) \rightarrow C(X \times Y) , \\ F(X, Y) &: C(X \times Y) \rightarrow C(X) \otimes_{\mathbb{Z}} C(Y) \end{aligned}$$

which are natural in X, Y , and such that

$$E(X, Y)(a \otimes b) = a \times b , \quad F(X, Y)(a \times b) = a \otimes b$$

for 0-simplexes $a : \Delta^0 \rightarrow X$, $b : \Delta^0 \rightarrow Y$.

(ii) The diagrams

$$\begin{array}{ccc} C(X) \otimes_{\mathbb{Z}} C(Y) & \xrightarrow{E(X, Y)} & C(X \times Y) & C(X \times Y) & \xrightarrow{F(X, Y)} & C(X) \otimes_{\mathbb{Z}} C(Y) \\ \downarrow T & & \downarrow T & \downarrow T & & \downarrow T \\ C(Y) \otimes_{\mathbb{Z}} C(X) & \xrightarrow{E(Y, X)} & C(Y \times X) & C(Y \times X) & \xrightarrow{F(Y, X)} & C(Y) \otimes_{\mathbb{Z}} C(X) \end{array}$$

are naturally chain homotopy commutative, with

$$\begin{aligned} T &: X \times Y \rightarrow Y \times X ; (x, y) \mapsto (y, x) , \\ T &: C(X)_p \otimes_{\mathbb{Z}} C(Y)_q \rightarrow C(Y)_q \otimes_{\mathbb{Z}} C(X)_p ; x \otimes y \mapsto (-)^{pq} y \otimes x . \end{aligned}$$

(iii) There exist chain homotopy inverse $\mathbb{R}(\infty)$ -coefficient \mathbb{Z}_2 -isovariant chain equivalences

$$\begin{aligned} E(X) &= \{E(X)_s \mid s \geq 0\} : C(X) \otimes_{\mathbb{Z}} C(X) \rightarrow C(X \times X) , \\ F(X) &= \{F(X)_s \mid s \geq 0\} : C(X \times X) \rightarrow C(X) \otimes_{\mathbb{Z}} C(X) \end{aligned}$$

which are natural in X , with $E(X)_0 = E(X, X)$, $F(X)_0 = F(X, X)$.

Proof. Standard acyclic model theory. □

Example 5.16. (i) The diagonal chain approximation for X

$$\Delta_{C(X)} = F(X)_0 \Delta_X : C(X) \xrightarrow{\Delta_X} C(X \times X) \xrightarrow{F(X)_0} C(X) \otimes_{\mathbb{Z}} C(X)$$

is used to define the cup products

$$\cup : H^p(X) \times H^q(X) \rightarrow H^{p+q}(X) ; (x, y) \mapsto (z \mapsto \langle x \otimes y, \Delta_{C(X)}(z) \rangle) .$$

(ii) The diagonal chain approximation of (i) is the degree 0 component of an $\mathbb{R}(\infty)$ -coefficient \mathbb{Z}_2 -isovariant chain map

$$F(X) \Delta_X : C(X) \xrightarrow{\Delta_X} C(X \times X) \xrightarrow{F(X)} C(X) \otimes_{\mathbb{Z}} C(X)$$

which is used to define the Steenrod squares – see Example 5.20 below.

(iii) The restriction of the diagonal chain approximation $\Delta_{C(S(LV))}$ to the chain homotopy deformation retract $C^{cell}(S(LV)) \subset C(S(LV))$ is chain homotopic to the cellular diagonal chain approximation Δ of 5.3 (iv). \square

Definition 5.17. Let V be an inner product space, and X a space.

(i) The *space level V -coefficient symmetric construction* is the \mathbb{Z}_2 -equivariant map

$$\phi_V(X) : S(LV) \times X \rightarrow X \times X ; (v, x) \mapsto (x, x) .$$

(ii) The *chain level V -coefficient symmetric construction* is the V -coefficient $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$\phi_V(X) : C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C(X) \rightarrow C(X) \otimes_{\mathbb{Z}} C(X) ,$$

given by the composite of the \mathbb{Z}_2 -isovariant chain maps

$$\begin{aligned} C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C(X) &\subset C(S(LV)) \otimes_{\mathbb{Z}} C(X) \\ &\xrightarrow[\simeq]{E(S(LV), X)} C(S(LV) \times X) \xrightarrow{\phi_V(X)} C(X \times X) , \\ C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C(X \times X) &\subset C(S(\infty)) \otimes_{\mathbb{Z}} C(X \times X) \\ &\xrightarrow[\simeq]{F(X)} C(X) \otimes_{\mathbb{Z}} C(X) \end{aligned}$$

(using an embedding $V \subseteq \mathbb{R}(\infty)$), inducing morphisms

$$\phi_V(X) : H_n(X) \rightarrow Q_V^n(C(X)) .$$

\square

Proposition 5.18. For any inner product space V and spaces X, Y there exists a natural $\mathbb{Z}[\mathbb{Z}_2]$ -module chain homotopy

$$\begin{aligned} \delta\phi_V(X, Y) : \\ \phi_V(X \times Y)E(X, Y) &\simeq (E(X, Y) \otimes E(X, Y))(\phi_V(X) \cup \phi_V(Y)) : \\ C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C(X) \otimes_{\mathbb{Z}} C(Y) &\rightarrow C(X \times Y) \otimes_{\mathbb{Z}} C(X \times Y) \end{aligned}$$

in the diagram

$$\begin{array}{ccc}
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C(X) \otimes_{\mathbb{Z}} C(Y) & \xrightarrow{\phi_V(X) \cup \phi_V(Y)} & C(X) \otimes_{\mathbb{Z}} C(Y) \otimes_{\mathbb{Z}} C(X) \otimes_{\mathbb{Z}} C(Y) \\
\downarrow \simeq \downarrow 1 \otimes E(X, Y) & \searrow \delta \phi_V(X, Y) & \downarrow \simeq \downarrow E(X, Y) \otimes E(X, Y) \\
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} C(X \times Y) & \xrightarrow{\phi_V(X \times Y)} & C(X \times Y) \otimes_{\mathbb{Z}} C(X \times Y)
\end{array}$$

Proof. Standard acyclic model theory. □

Terminology 5.19 For $V = \mathbb{R}(\infty)$ write

$$\begin{aligned}
\phi(X) &= \phi_{\mathbb{R}(\infty)}(X) : S(\infty) \times X \rightarrow X \times X, \\
\phi(X) &= \phi_{\mathbb{R}(\infty)}(X) : W \otimes_{\mathbb{Z}} C(X) \rightarrow C(X) \otimes_{\mathbb{Z}} C(X), \\
\delta \phi(X, Y) &= \delta \phi_{\mathbb{R}(\infty)}(X, Y) : \\
\phi(X \times Y) E(X, Y) &\simeq (E(X, Y) \otimes E(X, Y))(\phi(X) \cup \phi(Y)) : \\
&C(X) \otimes_{\mathbb{Z}} C(Y) \rightarrow C(X \times Y) \otimes_{\mathbb{Z}} C(X \times Y).
\end{aligned}$$

□

For $V = \mathbb{R}^k$ ($1 \leq k \leq \infty$) the adjoint of the natural $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$\phi_V(X) : W[0, k-1] \otimes_{\mathbb{Z}} C(X) \rightarrow C(X) \otimes_{\mathbb{Z}} C(X)$$

is a natural \mathbb{Z} -module chain map

$$\phi_V(X) : C(X) \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W[0, k-1], C(X) \otimes C(X))$$

inducing a natural transformation of homology groups

$$\phi_V(X) : H_*(X) \rightarrow Q_V^*(C(X)) = Q_{[0, k-1]}^*(C(X)).$$

By Proposition 5.18 there is defined a commutative diagram

$$\begin{array}{ccc}
H_m(X) \otimes_{\mathbb{Z}} H_n(Y) & \xrightarrow{E(X, Y)} & H_{m+n}(X \times Y) \\
\phi_V(X) \cup \phi_V(Y) \downarrow & & \phi_V(X \times Y) \downarrow \\
Q_V^{m+n}(C(X) \otimes_{\mathbb{Z}} C(Y)) & \xrightarrow[\cong]{E(X, Y)^\%} & Q_V^{m+n}(C(X \times Y)).
\end{array}$$

The V -coefficient symmetric constructions $\phi_V(X)$ are the restrictions of $\phi(X)$

$$\begin{aligned}\phi_V(X) &: S(LV) \times X \subseteq S(\infty) \times X \xrightarrow{\phi(X)} X \times X, \\ \phi_V(X) &: W[0, k-1] \otimes_{\mathbb{Z}} C(X) \subseteq W \otimes_{\mathbb{Z}} C(X) \xrightarrow{\phi(X)} C(X) \otimes_{\mathbb{Z}} C(X)\end{aligned}$$

and similarly for $\delta\phi_V(X, Y)$.

Remark 5.20. (i) The chain level symmetric construction

$$\phi(X) : C(X) \rightarrow \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(W, C(X) \otimes C(X))$$

is the symmetric construction in Ranicki [50], with a natural transformation

$$\phi(X) : H_*(X) \rightarrow Q^*(C(X)) .$$

The components of $\phi(X)$ are natural transformations

$$\phi(X)_s = \phi(X)(1_s \otimes -) : C(X)_s \rightarrow (C(X) \otimes_{\mathbb{Z}} C(X))_{n+s} \quad (0 \leq s \leq k-1)$$

such that

$$\begin{aligned}d\phi(X)_s + (-)^r \phi(X)_s d + (-)^{n+s-1} (\phi(X)_{s-1} + (-)^s T\phi(X)_{s-1}) &= 0 : \\ C(X)_n \rightarrow (C(X) \otimes C(X))_{n+s-1} &= \sum_r C(X)_{n-r+s-1} \otimes C(X)_r \quad (\phi(X)_{-1} = 0) .\end{aligned}$$

(ii) The Steenrod squares of X are given by

$$\begin{aligned}Sq^i &: H^r(X; \mathbb{Z}_2) \rightarrow H^{r+i}(X; \mathbb{Z}_2) ; \\ x &\mapsto (y \mapsto \langle x \otimes x, \phi(X)_{r-i}(y) \rangle) \quad (= 0 \text{ for } r < i) .\end{aligned}$$

The symmetric Poincaré complex associated by Mishchenko [44] to an n -dimensional geometric Poincaré complex X is

$$\sigma^*(X) = (C(X), \phi(X)[X] \in Q^n(C(X))) ,$$

with $[X] \in H_n(X)$ the fundamental class.

(iii) The $\mathbb{R}(\infty)$ -coefficient \mathbb{Z}_2 -isovariant chain homotopy $\delta\phi(X, Y)$ is used in the chain level proof of the Cartan product formula for the Steenrod squares

$$\begin{aligned}Sq^k &= \sum_{i+j=k} Sq^i \otimes Sq^j : \\ H^r(X \times Y; \mathbb{Z}_2) &= \sum_{p+q=r} H^p(X; \mathbb{Z}_2) \otimes H^q(Y; \mathbb{Z}_2) \\ &\rightarrow H^{r+k}(X \times Y; \mathbb{Z}_2) = \sum_{p+q+i+j=r+k} H^{p+i}(X; \mathbb{Z}_2) \otimes H^{q+j}(Y; \mathbb{Z}_2) .\end{aligned}$$

(iv) As in Steenrod [62] the *Hopf invariant* $\text{Hopf}(F) \in \mathbb{Z}$ of a map $F : S^{2n-1} \rightarrow S^n$ is defined by

$$\text{Hopf}(F) : H^n(S^n \cup_F D^{2n}) = \mathbb{Z} \rightarrow H^{2n}(S^n \cup_F D^{2n}) = \mathbb{Z} ; 1 \mapsto 1 \cup 1$$

and the *mod 2 Hopf invariant* $\text{Hopf}_2(F) \in \mathbb{Z}_2$ of a map $F : S^{n+k} \rightarrow S^n$ can be defined by

$$\begin{aligned} Sq^{k+1} &= \text{Hopf}_2(F) : \\ H^n(S^n \cup_F D^{n+k+1}; \mathbb{Z}_2) &= \mathbb{Z}_2 \rightarrow H^{n+k+1}(S^n \cup_F D^{n+k+1}; \mathbb{Z}_2) = \mathbb{Z}_2 . \end{aligned}$$

For $n = k + 1$ $\text{Hopf}_2(F) \in \mathbb{Z}_2$ is the mod 2 reduction of $\text{Hopf}(F) \in \mathbb{Z}$. By Adams [1] $\text{Hopf}(F) \in 2\mathbb{Z} \subset \mathbb{Z}$ for $n - 1 \neq 1, 3, 7$, and $\text{Hopf}_2(F) = 0 \in \mathbb{Z}_2$ for $k \neq 1, 3, 7$. □

Definition 5.21. (i) The relative singular chain complex of a pair of spaces $(X, A \subseteq X)$ is defined by

$$C(X, A) = C(X)/C(A) ,$$

with $H_*(C(X, A)) = H_*(X, A)$ the relative singular homology of (X, A) .

(ii) For a pointed space X let

$$\dot{C}(X) = C(X, \{*\}) = C(X)/C(\{*\})$$

be the reduced singular chain complex, with $H_*(\dot{C}(X)) = \dot{H}_*(X)$ the reduced singular homology of X .

(iii) For a non-empty subspace $A \subseteq X$ let

$$p_{X/A} : C(X, A) = C(X)/C(A) \rightarrow \dot{C}(X/A)$$

be the projection. □

In our applications $p_{X/A}$ will be a chain equivalence, e.g. if X is a *CW* complex and $A \subset X$ is a subcomplex.

Given pointed spaces X, Y let

$$Z = X \times \{*\} \cup \{*\} \times Y \subseteq X \times Y$$

so that

$$(X \times Y)/Z = X \wedge Y ,$$

and there are defined \mathbb{Z} -module chain maps

$$\begin{aligned} E(X, Y) : \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(Y) &= C(X, \{*\}) \otimes_{\mathbb{Z}} C(Y, \{*\}) \rightarrow C(X \times Y, Z) , \\ \dot{E}(X, Y) &= p_{X \wedge Y} E(X, Y) : \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(Y) \rightarrow \dot{C}(X \wedge Y) \end{aligned}$$

with $E(X, Y)$ a chain equivalence, and $p_{X \wedge Y} : C(X \times Y, Z) \rightarrow \dot{C}(X \wedge Y)$ the projection. As noted above, in our applications $p_{X \wedge Y}$ will be a chain equivalence, e.g. for CW complexes X, Y , so that $\dot{E}(X, Y)$ will also be a chain equivalence.

Definition 5.17 and Proposition 5.18 have pointed versions:

Definition 5.22. Let V be an inner product space, and X a pointed space.
(i) The *reduced space level V -coefficient symmetric construction* is the \mathbb{Z}_2 -equivariant map

$$\dot{\phi}_V(X) : S(LV)^+ \wedge X \rightarrow X \wedge X ; (v, x) \mapsto (x, x) .$$

(ii) The *chain level V -coefficient symmetric construction* is the V -coefficient $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$\dot{\phi}_V(X) : C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X) \rightarrow \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(X) ,$$

given by the composite of the \mathbb{Z}_2 -isovariant chain maps

$$\begin{aligned} C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X) &\subset C(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X) \\ &\xrightarrow[\simeq]{E(S(LV), X)} \dot{C}(S(LV)^+ \wedge X) \xrightarrow{\dot{\phi}_V(X)} \dot{C}(X \wedge X) , \\ C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X \wedge X) &\subset \dot{C}(S(\infty)) \otimes_{\mathbb{Z}} \dot{C}(X \wedge X) \\ &\xrightarrow[\simeq]{F(X)} \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(X) \end{aligned}$$

(using an embedding $V \subseteq \mathbb{R}(\infty)$), inducing morphisms

$$\dot{\phi}_V(X) : \dot{H}_n(X) \rightarrow Q_V^n(\dot{C}(X)) .$$

□

Proposition 5.23. For any inner product space V and pointed spaces X, Y there exists a natural $\mathbb{Z}[\mathbb{Z}_2]$ -module chain homotopy

$$\begin{aligned} \delta \dot{\phi}_V(X, Y) : \\ \dot{\phi}_V(X \times Y) E(X, Y) &\simeq (E(X, Y) \otimes E(X, Y)) (\dot{\phi}_V(X) \cup \dot{\phi}_V(Y)) : \\ C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(Y) &\rightarrow \dot{C}(X \wedge Y) \otimes_{\mathbb{Z}} \dot{C}(X \wedge Y) \end{aligned}$$

in the diagram

$$\begin{array}{ccc}
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(Y) & \xrightarrow{\dot{\phi}_V(X) \cup \dot{\phi}_V(Y)} & \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(Y) \\
\downarrow \simeq \downarrow 1 \otimes E(X,Y) & \searrow \delta \dot{\phi}_V(X,Y) & \downarrow \simeq \downarrow E(X,Y) \otimes E(X,Y) \\
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X \wedge Y) & \xrightarrow{\dot{\phi}_V(X \wedge Y)} & \dot{C}(X \wedge Y) \otimes_{\mathbb{Z}} \dot{C}(X \wedge Y)
\end{array}$$

□

Example 5.24. The $\mathbb{Z}[\mathbb{Z}_2]$ -module chain homotopy

$$\begin{aligned}
& \delta \dot{\phi}_V(S^1, X) : \\
& \dot{\phi}_V(S^1 \wedge X) \dot{E}(S^1, X) \simeq (\dot{E}(S^1, X) \otimes \dot{E}(S^1, X))(\dot{\phi}_V(S^1) \cup \dot{\phi}_V(X)) : \\
& \dot{C}(S^1) \otimes_{\mathbb{Z}} \dot{C}(X) \rightarrow \dot{C}(S^1 \wedge X) \otimes_{\mathbb{Z}} \dot{C}(S^1 \wedge X)
\end{aligned}$$

for $V = \mathbb{R}(\infty)$ gives the stability of the Steenrod squares, i.e. that the diagrams

$$\begin{array}{ccc}
\dot{H}^r(X; \mathbb{Z}_2) & \xrightarrow[\cong]{S} & \dot{H}^{r+1}(\Sigma X; \mathbb{Z}_2) \\
\downarrow Sq^i & & \downarrow Sq^i \\
\dot{H}^{r+i}(X; \mathbb{Z}_2) & \xrightarrow[\cong]{S} & \dot{H}^{r+i+1}(\Sigma X; \mathbb{Z}_2)
\end{array}$$

commute, with S the suspension isomorphisms induced by the natural chain equivalence

$$S = E(S^1, X) : S\dot{C}(X) = \dot{C}^{cell}(S^1) \otimes_{\mathbb{Z}} \dot{C}(X) \rightarrow \dot{C}(\Sigma X) = \dot{C}(S^1 \wedge X).$$

For $i = r + 1$ this is just the vanishing of cup products in a suspension. □

Example 5.25. It follows from the stability of the Steenrod squares and from

$$Sq^i = 0 : H^r(X; \mathbb{Z}_2) \rightarrow H^{r+i}(X; \mathbb{Z}_2) \text{ for } i > r$$

that if $X = \Sigma^k Y$ is a k -fold suspension then

$$Sq^i = 0 : \dot{H}^r(X; \mathbb{Z}_2) \rightarrow \dot{H}^{r+i}(X; \mathbb{Z}_2) \text{ for } i > r - k.$$

□

Proposition 4.41 for $U = V \subseteq \mathbb{R}(\infty)$ and $X = Y$ compact gives a commutative braid of exact sequences of stable \mathbb{Z}_2 -equivariant homotopy groups:

$$\begin{array}{ccccc}
& \xrightarrow{\quad \dot{\phi}_V(X) \quad} & & & \\
\{X; S(LV)^+ \wedge X \wedge X\}_{\mathbb{Z}_2} & \xrightarrow{s_{LV}} & \{X; X \wedge X\}_{\mathbb{Z}_2} & \xrightarrow{s_{LV}^*} & \{S(LV)^+ \wedge X; X \wedge X\}_{\mathbb{Z}_2} \\
& \searrow & \nearrow & \searrow & \nearrow \\
& & \{X; X \wedge X\}_{\mathbb{Z}_2} & & \{S(LV \oplus LV)^+ \wedge X; LV^\infty \wedge X \wedge X\}_{\mathbb{Z}_2} \\
& \nearrow & \searrow & \nearrow & \searrow \\
\{LV^\infty \wedge X; X \wedge X\}_{\mathbb{Z}_2} & \xrightarrow{0_{LV}^*} & \{X; X \wedge X\}_{\mathbb{Z}_2} & \xrightarrow{0_{LV}} & \{X; LV^\infty \wedge X \wedge X\}_{\mathbb{Z}_2} \\
& \searrow & \nearrow & \searrow & \nearrow \\
& & \{X; LV^\infty \wedge X \wedge X\}_{\mathbb{Z}_2} & & A
\end{array}$$

with

$$\begin{aligned}
\Delta_X &= (0, 1) \in \{X; X \wedge X\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge (X \wedge X)\}_{\mathbb{Z}_2} \oplus \{X; X\} , \\
s_{LV}^*(\Delta_X) &= \dot{\phi}_V(X) \in \{S(LV)^+ \wedge X; X \wedge X\}_{\mathbb{Z}_2} , \\
[\dot{\phi}_V(X)] &= \dot{\phi}_{V \oplus V}(V^\infty \wedge X) \in \{S(LV \oplus LV)^+ \wedge X; LV^\infty \wedge X \wedge X\}_{\mathbb{Z}_2} \\
&= \{S(LV \oplus LV)^+ \wedge (V^\infty \wedge X); (V^\infty \wedge X) \wedge (V^\infty \wedge X)\}_{\mathbb{Z}_2} , \\
0_{LV}(\Delta_X) &= \Delta_{V^\infty \wedge X} = (0, 1) \\
&\in \{V^\infty \wedge X; (V^\infty \wedge X) \wedge (V^\infty \wedge X)\}_{\mathbb{Z}_2} = \{X; LV^\infty \wedge (X \wedge X)\}_{\mathbb{Z}_2} \\
&= \{X; S(\infty)/S(LV) \wedge (X \wedge X)\}_{\mathbb{Z}_2} \oplus \{X; X\} , \\
A &= \{S(LV)^+ \wedge X; LV^\infty \wedge X \wedge X\}_{\mathbb{Z}_2} = \{X; \Sigma S(LV)^+ \wedge (X \wedge X)\}_{\mathbb{Z}_2} .
\end{aligned}$$

If $X = V^\infty \wedge X_0$ for some pointed space X_0 then

$$\begin{aligned}
\Delta_{X_0} &\in \{X_0; X_0 \wedge X_0\}_{\mathbb{Z}_2} = \{LV^\infty \wedge X; X \wedge X\}_{\mathbb{Z}_2} , \\
\Delta_X &= 0_{LV}^*(\Delta_{X_0}) \in \text{im}(0_{LV}^* : \{LV^\infty \wedge X; X \wedge X\}_{\mathbb{Z}_2} \rightarrow \{X; X \wedge X\}_{\mathbb{Z}_2}) \\
&= \ker(s_{LV}^* : \{X; X \wedge X\}_{\mathbb{Z}_2} \rightarrow \{S(LV)^+ \wedge X; X \wedge X\}_{\mathbb{Z}_2}) , \\
\dot{\phi}_V(X) &= 0 \in \{S(LV)^+ \wedge X; X \wedge X\}_{\mathbb{Z}_2} ,
\end{aligned}$$

generalizing the vanishing of the cup product in a suspension (the special case $V = \mathbb{R}$, on the chain level). More precisely :

Proposition 5.26. (i) For any pointed spaces X, Y and inner product space V

$$\begin{aligned}
\dot{\phi}_V(X \wedge Y) &= \dot{\phi}_V(X) \wedge \Delta_Y = \Delta_X \wedge \dot{\phi}_V(Y) : \\
S(LV)^+ \wedge X \wedge Y &\rightarrow (X \wedge Y) \wedge (X \wedge Y) = (X \wedge X) \wedge (Y \wedge Y) .
\end{aligned}$$

(ii) The space level symmetric construction $\dot{\phi}_V(V^\infty)$ has a natural \mathbb{Z}_2 -equivariant null-homotopy

$$\delta \dot{\phi}_V(V^\infty) : \dot{\phi}_V(V^\infty) \simeq * : S(LV)^+ \wedge V^\infty \rightarrow V^\infty \wedge V^\infty .$$

(iii) *The space level V -coefficient symmetric construction $\dot{\phi}_V(V^\infty \wedge X)$ has a natural \mathbb{Z}_2 -equivariant null-homotopy*

$$\begin{aligned} \delta \dot{\phi}_V(V^\infty \wedge X) &= \delta \dot{\phi}_V(V^\infty) \wedge \Delta_X : \\ \dot{\phi}_V(V^\infty \wedge X) &\simeq * : S(LV)^+ \wedge V^\infty \wedge X \rightarrow V^\infty \wedge X \wedge V^\infty \wedge X \end{aligned}$$

inducing a natural $\mathbb{Z}[\mathbb{Z}_2]$ -module null-chain homotopy

$$\begin{aligned} \delta \dot{\phi}_V(V^\infty \wedge X) : \dot{\phi}_V(V^\infty \wedge X) &\simeq 0 : \\ C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(V^\infty \wedge X) &\rightarrow \dot{C}(V^\infty \wedge X) \otimes_{\mathbb{Z}} \dot{C}(V^\infty \wedge X) . \end{aligned}$$

Proof. (i) By construction.

(ii) Apply the construction of 2.10 to the \mathbb{Z}_2 -equivariant map

$$p = \kappa_V : LV^\infty \wedge V^\infty \rightarrow V^\infty \wedge V^\infty ; (r, s) \mapsto (r + s, -r + s)$$

to obtain a \mathbb{Z}_2 -equivariant null-homotopy

$$\delta \dot{\phi}_V(V^\infty) = \delta p : CS(LV)^+ \wedge V^\infty \rightarrow V^\infty \wedge V^\infty ; (t, u, v) \mapsto p([t, u], v)$$

of

$$[\delta \dot{\phi}_V(V^\infty)] = \dot{\phi}_V(V^\infty) : S(LV)^+ \wedge V^\infty \rightarrow V^\infty \wedge V^\infty ; (u, v) \mapsto (v, v) ,$$

using the projection

$$CS(LV)^+ \rightarrow LV^\infty ; (t, u) \mapsto [t, u] = \frac{tu}{1-t} .$$

(iii) Combine (i) and (ii).

□

More generally :

Proposition 5.27. *Let U, V be inner product spaces.*

(i) *There is defined a \mathbb{Z}_2 -equivariant homotopy*

$$\begin{aligned} \delta \dot{\phi}_{U,V} : \kappa_V \dot{\phi}_{U \oplus V}(V^\infty) &\simeq (\dot{\phi}_U(S^0) \wedge 1_{LV^\infty \wedge V^\infty})(j_{LU, LV} \wedge 1_{V^\infty}) \\ &: S(LU \oplus LV)^+ \wedge V^\infty \rightarrow LV^\infty \wedge V^\infty . \end{aligned}$$

in the diagram

$$\begin{array}{ccc}
S(LU \oplus LV)^+ \wedge V^\infty & \xrightarrow{j_{LU, LV} \wedge 1_{V^\infty}} & S(LU)^+ \wedge LV^\infty \wedge V^\infty \\
\downarrow \phi_{U \oplus V}(V^\infty) & & \downarrow \phi_U(S^0) \wedge 1_{LV^\infty \wedge V^\infty} \\
V^\infty \wedge V^\infty & \xrightarrow{\kappa_V} & LV^\infty \wedge V^\infty
\end{array}$$

with

$$\begin{aligned}
j_{LU, LV} &= \text{projection} : \\
S(LU \oplus LV)^+ &= (D(LU) \times S(LV) \cup S(LU) \times D(LV))^+ \\
&\rightarrow S(LU) \times D(LV) / S(LU) \times S(LV) = S(LU)^+ \wedge D(LV) / S(LV) \\
&= S(LU)^+ \wedge LV^\infty .
\end{aligned}$$

(ii) For any pointed space X there is defined a natural \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned}
\delta \phi_{U, V}(X) : \\
(\kappa_V \wedge 1_{X \wedge X}) \phi_{U \oplus V}(V^\infty \wedge X) &\simeq (\phi_U(X) \wedge 1_{LV^\infty \wedge V^\infty})(j_{LU, LV} \wedge 1_{V^\infty}) \\
&: S(LU \oplus LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge X \wedge X .
\end{aligned}$$

in the diagram

$$\begin{array}{ccc}
S(LU \oplus LV)^+ \wedge V^\infty \wedge X & \xrightarrow{j_{LU, LV} \wedge 1_{V^\infty \wedge X}} & S(LU)^+ \wedge LV^\infty \wedge V^\infty \wedge X \\
\downarrow \phi_{U \oplus V}(V^\infty \wedge X) & & \downarrow \phi_U(X) \wedge 1_{LV^\infty \wedge V^\infty} \\
V^\infty \wedge V^\infty \wedge X \wedge X & \xrightarrow{\kappa_V \wedge 1_{X \wedge X}} & LV^\infty \wedge V^\infty \wedge X \wedge X
\end{array}$$

(iii) The natural \mathbb{Z}_2 -equivariant homotopy in (ii) induces a natural $U \oplus V$ -coefficient \mathbb{Z}_2 -isovariant chain homotopy

$$\begin{aligned}
\delta \phi_{U, V} : \phi_{U \oplus V}(V^\infty \wedge X) &\simeq \phi_U(X) j_{LU, LV} \\
&: \dot{C}(V^\infty \wedge X) \rightarrow \dot{C}(V^\infty \wedge X) \otimes_{\mathbb{Z}} \dot{C}(V^\infty \wedge X) .
\end{aligned}$$

Proof. (i) Use the pushout square of \mathbb{Z}_2 -spaces given by Proposition 2.11 (ii)

$$\begin{array}{ccc}
S(LU \oplus LV)^+ & \xrightarrow{j_{LU, LV}} & S(LU)^+ \wedge LV^\infty \\
\downarrow & & \downarrow k_{LU, LV} \\
CS(LU \oplus LV)^+ & \xrightarrow{\delta j_{LU, LV}} & \Sigma S(LV)^+
\end{array}$$

to define

$$\delta \phi_{U, V} : I \times S(LU \oplus LV)^+ \wedge V^\infty \rightarrow LV^\infty \wedge V^\infty ; (t, (u, v, w)) \mapsto (tv, w) .$$

(For $U = \{0\}$ $\delta\dot{\phi}_{U,V} = \delta\dot{\phi}_V$ was already defined in the proof of Proposition 5.26 (ii)).

(ii)+(iii) Immediate from (i), with $\delta\dot{\phi}_{U,V}(X) = \delta\dot{\phi}_{U,V} \wedge \Delta_X$. \square

We shall also need a relative version of the symmetric construction:

Definition 5.28. Given a pair of pointed spaces $(A, B \subseteq A)$ there is defined a *relative space level reduced V -coefficient symmetric construction* \mathbb{Z}_2 -equivariant map

$$\dot{\phi}_V(A, B) : S(LV)^+ \wedge A/B \rightarrow (A \wedge A)/(B \wedge B)$$

which fits into a natural transformation of cofibration sequences of \mathbb{Z}_2 -spaces

$$\begin{array}{ccccccc} S(LV)^+ \wedge B & \longrightarrow & S(LV)^+ \wedge A & \longrightarrow & S(LV)^+ \wedge A/B & \longrightarrow & S(LV)^+ \wedge \Sigma B \\ \downarrow \phi_V(B) & & \downarrow \phi_V(A) & & \downarrow \phi_V(A, B) & & \downarrow \Sigma \phi_V(B) \\ B \wedge B & \longrightarrow & A \wedge A & \longrightarrow & (A \wedge A)/(B \wedge B) & \longrightarrow & \Sigma(B \wedge B) \end{array}$$

\square

Proposition 5.29. Given a pointed map $F : X \rightarrow Y$ let

$$(A, B \subseteq A) = (\mathcal{M}(F), \{1\} \times X), \quad Z = A/B = \mathcal{C}(F)$$

so that there is defined a cofibration sequence of pointed maps

$$X \xrightarrow{F} Y \xrightarrow{G} Z \xrightarrow{H} \Sigma X \xrightarrow{\Sigma F} \Sigma Y \longrightarrow \dots$$

with G the inclusion, H the projection. The space level reduced V -coefficient symmetric construction of Z is given by

$$\begin{aligned} \dot{\phi}_V(Z) : S(LV)^+ \wedge Z &\xrightarrow{\dot{\phi}_V(A, B)} \\ A \wedge A/B \wedge B &\longrightarrow A \times A/(A \times B \cup B \times A) = Z \wedge Z \end{aligned}$$

Proof. By construction. \square

5.3 The geometric Hopf invariant $h_V(F)$

Let $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ be a stable \mathbb{Z}_2 -map, with X, Y pointed \mathbb{Z}_2 -spaces and V an inner product \mathbb{Z}_2 -space, with

$$V = V_- \oplus V_+, \quad V_\pm = \{v \in V \mid Tv = \pm v\}.$$

(We shall be mainly concerned with the case when F is the square of a stable nonequivariant map). The geometric Hopf invariant uses the difference construction to associate to F a \mathbb{Z}_2 -map $\delta(p, q)$ whose \mathbb{Z}_2 -homotopy class is the primary obstruction to the desuspension of F . The diagram

$$\begin{array}{ccc} V^\infty \wedge X^{\mathbb{Z}_2} & \xrightarrow{1 \wedge i_X} & V^\infty \wedge X \\ 1 \wedge \rho(F) \downarrow & & \downarrow F \\ V^\infty \wedge Y^{\mathbb{Z}_2} & \xrightarrow{1 \wedge i_Y} & V^\infty \wedge Y \end{array}$$

does not commute in general, with $i_X : X^{\mathbb{Z}_2} \rightarrow X$, $i_Y : Y^{\mathbb{Z}_2} \rightarrow Y$ the inclusions and $\rho(F) : X^{\mathbb{Z}_2} \rightarrow Y^{\mathbb{Z}_2}$ the restriction of F to the fixed point sets. However, the \mathbb{Z}_2 -maps

$$p = F(1 \wedge i_X), \quad q = (1 \wedge i_Y)(1 \wedge \rho(F)) : V^\infty \wedge X^{\mathbb{Z}_2} \rightarrow V^\infty \wedge Y$$

are such that

$$p(0, v, x) = F(0, v, x) = (0, v, \rho(F)(x)) = q(0, v, x) \quad (v \in V, x \in X^{\mathbb{Z}_2})$$

so the relative difference (2.18 (ii)) \mathbb{Z}_2 -map

$$\delta(p, q) : \Sigma S(V_-)^+ \wedge V_+^\infty \wedge X^{\mathbb{Z}_2} \rightarrow V^\infty \wedge Y$$

is defined, with

$$p - q : V^\infty \wedge X^{\mathbb{Z}_2} \xrightarrow{\alpha_{V_-} \wedge i_X} \Sigma S(V_-)^+ \wedge V_+^\infty \wedge X \xrightarrow{\delta(p, q)} V^\infty \wedge Y.$$

If F is \mathbb{Z}_2 -homotopic to $1 \wedge F_0 : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ for some \mathbb{Z}_2 -map $F_0 : X \rightarrow Y$ then

$$\delta(p, q) \simeq \delta(1 \wedge F_0, 1 \wedge F_0) \simeq *.$$

Now let X, Y be spaces, and V an inner product space. The square of a nonequivariant stable map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is a stable \mathbb{Z}_2 -map

$$F \wedge F : V^\infty \wedge V^\infty \wedge X \wedge X \rightarrow V^\infty \wedge V^\infty \wedge Y \wedge Y .$$

The geometric Hopf invariant of F is the above difference construction

$$h_V(F) = \delta(p, q) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow V^\infty \wedge V^\infty \wedge Y \wedge Y$$

applied to the stable \mathbb{Z}_2 -map

$$\begin{aligned} G &= (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge 1) : \\ &LV^\infty \wedge V^\infty \wedge X \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y ; \\ &(u, v, x_1, x_2) \mapsto ((w_1 - w_2)/2, (w_1 + w_2)/2, y_1, y_2) \\ &(F(u + v, x_1) = (w_1, y_1), F(-u + v, x_2) = (w_2, y_2)) \end{aligned}$$

with

$$\rho(G) = F : V^\infty \wedge X \rightarrow V^\infty \wedge Y .$$

In essence, the geometric Hopf invariant measures the difference $(F \wedge F)\Delta_X - \Delta_Y F$, given that $(F \wedge F)\Delta_{V^\infty \wedge X} = \Delta_{V^\infty \wedge Y} F$. The diagram of \mathbb{Z}_2 -equivariant maps

$$\begin{array}{ccc} LV^\infty \wedge V^\infty \wedge X & \xrightarrow{1 \wedge \Delta_X} & LV^\infty \wedge V^\infty \wedge X \wedge X \\ \downarrow 1 \wedge F & & \downarrow G \\ LV^\infty \wedge V^\infty \wedge Y & \xrightarrow{1 \wedge \Delta_Y} & LV^\infty \wedge V^\infty \wedge Y \wedge Y \end{array}$$

does not commute in general. However, the \mathbb{Z}_2 -equivariant maps defined by

$$\begin{aligned} p &= G(1 \wedge \Delta_X) : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y ; \\ &(u, v, x) \mapsto ((w_1 - w_2)/2, (w_1 + w_2)/2, y_1, y_2) \\ &(F(u + v, x) = (w_1, y_1), F(-u + v, x) = (w_2, y_2)) , \\ q &= (1 \wedge \Delta_Y)(1 \wedge F) : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y ; \\ &(u, v, x) \mapsto (u, w, y, y) \quad (F(v, x) = (w, y)) \end{aligned}$$

agree on $0^+ \wedge V^\infty \wedge X = V^\infty \wedge X \subset LV^\infty \wedge V^\infty \wedge X$, with

$$\begin{aligned} p| &= q| = (\kappa_V^{-1} \wedge 1)\Delta_{V^\infty \wedge Y} F = (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge 1)\Delta_{V^\infty \wedge X} : \\ &V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y \end{aligned}$$

on account of the commutative diagram

$$\begin{array}{ccccc}
V^\infty \wedge X & \xrightarrow{\Delta_{V^\infty \wedge X}} & (V^\infty \wedge X) \wedge (V^\infty \wedge X) & \xrightarrow{\kappa_V^{-1} \wedge 1} & LV^\infty \wedge V^\infty \wedge X \wedge X \\
\downarrow F & & \downarrow F \wedge F & & \downarrow (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge 1) \\
V^\infty \wedge Y & \xrightarrow{\Delta_{V^\infty \wedge Y}} & (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) & \xrightarrow{\kappa_V^{-1} \wedge 1} & LV^\infty \wedge V^\infty \wedge Y \wedge Y .
\end{array}$$

Definition 5.30. The *geometric Hopf invariant* of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is the \mathbb{Z}_2 -equivariant map given by the relative difference of the \mathbb{Z}_2 -equivariant maps $p = (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge \Delta_X)$, $q = (1 \wedge \Delta_Y)(1 \wedge F)$

$$\begin{aligned}
h_V(F) &= \delta(p, q) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y ; \\
(t, u, v, x) &\mapsto \begin{cases} q([1 - 2t, u], v, x) & \text{if } 0 \leq t \leq 1/2 \\ p([2t - 1, u], v, x) & \text{if } 1/2 \leq t \leq 1 \end{cases} \\
&(t \in I, u \in S(LV), v \in V, x \in X) .
\end{aligned}$$

□

Remark 5.31. $h_V(F)$ is the geometric Hopf invariant of Crabb [12, p. 61], Crabb and James [13, p. 306]. □

Example 5.32. Suppose that $V = \{0\}$, so that

$$V^\infty = S^0, \quad S(LV) = \emptyset, \quad S(LV)^+ = \{*\} .$$

The geometric Hopf invariant of a map

$$F : V^\infty \wedge X = X \rightarrow V^\infty \wedge Y = Y$$

is

$$h_V(F) = * : \Sigma S(LV)^+ \wedge V^\infty \wedge X = \{*\} \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y = Y \wedge Y .$$

□

The geometric Hopf invariant $h_V(F)$ of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ has the following properties.

Proposition 5.33. (i) (*Naturality*) If $f : X \rightarrow X'$, $g : Y \rightarrow Y'$, $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$, $F' : V^\infty \wedge X' \rightarrow V^\infty \wedge Y'$ are maps such that there is defined a commutative diagram

$$\begin{array}{ccc}
V^\infty \wedge X & \xrightarrow{F} & V^\infty \wedge Y \\
1 \wedge f \downarrow & & \downarrow 1 \wedge g \\
V^\infty \wedge X' & \xrightarrow{F'} & V^\infty \wedge Y'
\end{array}$$

then there is defined a commutative diagram

$$\begin{array}{ccc}
\Sigma S(LV)^+ \wedge V^\infty \wedge X & \xrightarrow{h_V(F)} & LV^\infty \wedge V^\infty \wedge Y \wedge Y \\
1 \wedge f \downarrow & & \downarrow 1 \wedge g \wedge g \\
\Sigma S(LV)^+ \wedge V^\infty \wedge X' & \xrightarrow{h_V(F')} & LV^\infty \wedge V^\infty \wedge Y' \wedge Y'
\end{array}$$

(ii) (*Homotopy invariance*) The \mathbb{Z}_2 -equivariant homotopy class of $h_V(F)$ depends only on the homotopy class of F .

(iii) (*Suspension formula*) Let U be another inner product space. The geometric Hopf invariant $h_{U \oplus V}(1_{U^\infty} \wedge F)$ of the stabilization of $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$

$$1_{U^\infty} \wedge F : (U \oplus V)^\infty \wedge X = U^\infty \wedge V^\infty \wedge X \rightarrow (U \oplus V)^\infty \wedge Y = U^\infty \wedge V^\infty \wedge Y$$

is determined by $h_V(F)$, with a commutative diagram

$$\begin{array}{ccc}
U^\infty \wedge V^\infty \wedge \Sigma S(LU \oplus LV)^+ \wedge X & & \\
\downarrow h_{U \oplus V}(1_{U^\infty} \wedge F) & \searrow 1 \wedge \Sigma j_{LV, LU} & \\
& (U \oplus LU)^\infty \wedge V^\infty \wedge \Sigma S(LV)^+ \wedge X & \\
& \swarrow 1_{(U \oplus LU)^\infty} \wedge h_V(F) & \\
(U \oplus LU)^\infty \wedge (V \oplus LV)^\infty \wedge Y \wedge Y & &
\end{array}$$

with $j_{LV, LU}$ the \mathbb{Z}_2 -equivariant adjunction Umkehr map of the \mathbb{Z}_2 -equivariant embedding

$$S(LU) \times D(LV) \subset D(LU) \times S(LV) \cup S(LU) \times D(LV) = S(LU \oplus LV),$$

that is

$$\begin{aligned}
j_{LV, LU} &= \text{projection} : S(LU \oplus LV)^+ \rightarrow \\
&S(LU \oplus LV)/(S(LU) \times D(LV)) = LU^\infty \wedge S(LV)^+.
\end{aligned}$$

(iv) (*Symmetrization*) Write

$$A = V^\infty \wedge X, \quad B = LV^\infty \wedge V^\infty \wedge Y \wedge Y.$$

The images of the \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant $h_V(F) \in [\Sigma S(LV)^+ \wedge A; B]_{\mathbb{Z}_2}$ under the maps in the commutative braid of exact sequences of \mathbb{Z}_2 -equivariant homotopy groups

$$\begin{array}{ccccc}
 & & k_{LV, LV}^* & & \\
 & \nearrow & & \searrow & \\
 [\Sigma S(LV)^+ \wedge A; B]_{\mathbb{Z}_2} & & [S(LV)^+ \wedge LV^\infty \wedge A; B]_{\mathbb{Z}_2} & & \\
 \downarrow \alpha_{LV}^* & & \downarrow s_{LV}^* & & \downarrow j_{LV, LV}^* \\
 [LV^\infty \wedge A; B]_{\mathbb{Z}_2} & & [S(LV \oplus LV)^+ \wedge A; B]_{\mathbb{Z}_2} & & \\
 \uparrow 0_{LV}^* & & \downarrow s_{LV \oplus LV}^* & & \\
 [LV^\infty \wedge LV^\infty \wedge A; B]_{\mathbb{Z}_2} & & [A; B]_{\mathbb{Z}_2} & & \\
 \downarrow 0_{LV \oplus LV}^* & & \uparrow 0_{LV}^* & & \\
 & & & &
 \end{array}$$

are given by

$$\begin{aligned}
 \alpha_{LV}^* h_V(F) &= (F \wedge F) \Delta_X - \Delta_Y F \in [LV^\infty \wedge A; B]_{\mathbb{Z}_2}, \\
 k_{LV, LV}^* h_V(F) &= (F \wedge F) \dot{\phi}_V(X) - \dot{\phi}_V(Y) F \in [S(LV)^+ \wedge LV^\infty \wedge A; B]_{\mathbb{Z}_2}.
 \end{aligned}$$

In particular, the diagram of \mathbb{Z}_2 -equivariant maps

$$\begin{array}{ccc}
 LV^\infty \wedge V^\infty \wedge X & \xrightarrow{\alpha_{LV} \wedge 1} & \Sigma S(LV)^+ \wedge V^\infty \wedge X \\
 & \searrow (F \wedge F) \Delta_X - \Delta_Y (1 \wedge F) & \downarrow h_V(F) \\
 & & LV^\infty \wedge V^\infty \wedge Y \wedge Y
 \end{array}$$

commutes up to \mathbb{Z}_2 -equivariant homotopy.

(v) (Composition formula) The geometric Hopf invariant of the composite

$$GF : V^\infty \wedge X \rightarrow V^\infty \wedge Z$$

of maps $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$, $G : V^\infty \wedge Y \rightarrow V^\infty \wedge Z$ is given up to \mathbb{Z}_2 -equivariant homotopy by

$$\begin{aligned}
 h_V(GF) &= h_V(G)(1 \wedge F) + (\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge 1)h_V(F) : \\
 &\quad \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Z \wedge Z.
 \end{aligned}$$

(vi) (Product formula) Let $F_i : V_i^\infty \wedge X_i \rightarrow V_i^\infty \wedge Y_i$, $i = 1, 2$, be two maps. The geometric Hopf invariant $h_{V_1 \oplus V_2}(F_1 \wedge F_2)$ of the smash product

$$F_1 \wedge F_2 : (V_1 \oplus V_2)^\infty \wedge (X_1 \wedge X_2) \rightarrow (V_1 \oplus V_2)^\infty \wedge (Y_1 \wedge Y_2)$$

is the composite

$$\begin{aligned} & ((\Delta_{Y_1} F_1 \wedge h_{V_2}(F_2)) \vee (h_{V_1}(F_1) \wedge h_{V_2}(F_2)) \vee (h_{V_1}(F_1) \wedge \Delta_{Y_2} F_2)) \circ \Sigma c : \\ & \Sigma S(LV_1 \oplus LV_2)^+ \wedge (V_1 \oplus V_2)^\infty \wedge X_1 \wedge X_2 \\ & \rightarrow ((LV_1 \oplus LV_2)^\infty \wedge (V_1 \oplus V_2)^\infty \wedge (Y_1 \wedge Y_2) \wedge (Y_1 \wedge Y_2)) \end{aligned}$$

where

$$c : A = S(LV_1 \oplus LV_2)^+ \rightarrow B_1 \vee B_2 \vee B_3$$

is the Umkehr map for the inclusion of the submanifold

$$S(LV_1) \sqcup S(LV_2) \sqcup (S(LV_1) \times S(LV_2)) \hookrightarrow S(LV_1 \oplus LV_2) .$$

The \mathbb{Z}_2 -equivariant stable homotopy class

$$c = (c_1, c_2, c_3) \in \{A; B_1 \vee B_2 \vee B_3\}_{\mathbb{Z}_2} = \{A; B_1\}_{\mathbb{Z}_2} \oplus \{A; B_2\}_{\mathbb{Z}_2} \oplus \{A; B_3\}_{\mathbb{Z}_2}$$

has components

$$\begin{aligned} c_1 &= j_{LV_2, LV_1} : A = S(LV_1 \oplus LV_2) \\ &\rightarrow B_1 = S(LV_1 \oplus LV_2)/(S(LV_1) \times D(LV_2)) = LV_1^\infty \wedge S(LV_2)^+ , \\ c_2 &= j_{LV_1, LV_2} : A = S(LV_1 \oplus LV_2) \\ &\rightarrow B_2 = S(LV_1 \oplus LV_2)/(D(LV_1) \times S(LV_2)) = S(LV_1)^+ \wedge LV_2^\infty , \\ c_3 &= \text{projection} : A = S(LV_1 \oplus LV_2) \\ &\rightarrow B_3 = S(LV_1 \oplus LV_2)/(S(LV_1) \times D(LV_2) \sqcup D(LV_1) \times S(LV_2)) \\ &= \Sigma(S(LV_1)^+ \wedge S(LV_2)^+) . \end{aligned}$$

(vii) (One-point union formula) The geometric Hopf invariant of a map from a one-point union

$$F_1 \vee F_2 : V^\infty \wedge (X_1 \vee X_2) = (V^\infty \wedge X_1) \vee (V^\infty \wedge X_2) \rightarrow V^\infty \wedge Y$$

is given by

$$\begin{aligned} h_V(F_1 \vee F_2) &= h_V(F_1) \vee h_V(F_2) : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge (X_1 \vee X_2) &\rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y . \end{aligned}$$

(viii) (Sum map) For $V \neq \{0\}$ the geometric Hopf invariant of the sum map

$$\nabla_V : V^\infty \rightarrow V^\infty \vee V^\infty$$

is given up to \mathbb{Z}_2 -equivariant homotopy by the composite

$$h_V(\nabla_V) = i_V j_V : \Sigma S(LV)^+ \wedge V^\infty \rightarrow (V^\infty \vee V^\infty) \wedge (V^\infty \vee V^\infty)$$

with

$$\begin{aligned} i_V : (V^\infty \wedge V^\infty) \vee (V^\infty \wedge V^\infty) &\rightarrow (V^\infty \vee V^\infty) \wedge (V^\infty \vee V^\infty) ; \\ (u, v)_1 \mapsto (u_1, v_2) \quad , \quad (u, v)_2 &\mapsto (u_2, v_1) \end{aligned}$$

and

$$\begin{aligned} j_V : \Sigma S(LV)^+ \wedge V^\infty &\rightarrow (V^\infty \wedge V^\infty) \vee (V^\infty \wedge V^\infty) ; \\ (t, u, v) \mapsto \begin{cases} (\alpha_V(2t, u), v)_1 & \text{if } 0 \leq t \leq 1/2 \\ (v, \alpha_V(2t - 1, u))_2 & \text{if } 1/2 \leq t \leq 1 . \end{cases} \end{aligned}$$

(ix) (*Sum formula*) For $V \neq \{0\}$ the geometric Hopf invariant of the sum of maps $F, G : V^\infty \wedge X \rightarrow V^\infty \wedge Y$

$$F + G : V^\infty \wedge X \rightarrow V^\infty \wedge Y$$

is given up to \mathbb{Z}_2 -equivariant homotopy by

$$\begin{aligned} h_V(F + G) &= (h_V(F) \vee h_V(G) \vee ((F \wedge G) \vee (G \wedge F))(1 \wedge \Delta_X)) \circ (d \wedge 1) : \\ &\Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y \end{aligned}$$

with

$$\begin{aligned} d : A &= \Sigma S(LV)^+ \wedge V^\infty \rightarrow \\ B_1 \vee B_2 \vee B_3 &= (\Sigma S(LV)^+ \wedge V^\infty) \vee (\Sigma S(LV)^+ \wedge V^\infty) \\ &\quad \vee ((V^\infty \wedge V^\infty) \vee (V^\infty \wedge V^\infty)) \end{aligned}$$

a \mathbb{Z}_2 -equivariant map with stable components

$$d = (d_1, d_2, d_3) \in \{A; B_1 \vee B_2 \vee B_3\}_{\mathbb{Z}_2} = \{A; B_1\}_{\mathbb{Z}_2} \oplus \{A; B_2\}_{\mathbb{Z}_2} \oplus \{A; B_3\}_{\mathbb{Z}_2}$$

given by

$$\begin{aligned} d_1 = d_2 = 1 : A &= \Sigma S(LV)^+ \wedge V^\infty \rightarrow B_1 = B_2 = \Sigma S(LV)^+ \wedge V^\infty , \\ d_3 = j_V : A &= \Sigma S(LV)^+ \wedge V^\infty \rightarrow B_3 = (V^\infty \wedge V^\infty) \vee (V^\infty \wedge V^\infty) \end{aligned}$$

Thus up to stable \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned} h_V(F + G) &= h_V(F) + h_V(G) + [(F \wedge G) \vee (G \wedge F)](j_V \wedge \Delta_X) : \\ &\Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y . \end{aligned}$$

Proof. (i) By construction.

(ii) A homotopy $G : F \simeq F' : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ determines \mathbb{Z}_2 -equivariant homotopies

$$\begin{aligned}
1 \wedge G : 1 \wedge F &\simeq 1 \wedge F' : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y, \\
G \wedge G : F \wedge F &\simeq F' \wedge F' : (V^\infty \wedge X) \wedge (V^\infty \wedge X) \rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y)
\end{aligned}$$

and hence by 2.23 (i) a \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned}
&\delta((\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge \Delta_X), (1 \wedge \Delta_Y)(1 \wedge G)) : \\
h_V(F) &= \delta((\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge \Delta_X), (1 \wedge \Delta_Y)(1 \wedge F)) \\
&\simeq h_V(F') = \delta((\kappa_V^{-1} \wedge 1)(F' \wedge F')(\kappa_V \wedge \Delta_X), (1 \wedge \Delta_Y)(1 \wedge F')) \\
&: \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y.
\end{aligned}$$

(iii) The identity $h_{U \oplus V}(1_{U^\infty} \wedge F) = (1_{(U \oplus LV)^\infty} \wedge h_V(F))(1 \wedge \Sigma j_{LU, LV})$ follows from the commutative diagram

$$\begin{array}{ccc}
(LU \oplus LV)^\infty = LU^\infty \wedge LV^\infty & \xrightarrow{\alpha_{L(U \oplus V)}} & \Sigma S(L(U \oplus V))^\infty \\
& \searrow 1_{LU^\infty} \wedge \alpha_{LV} & \downarrow \Sigma j_{LU, LV} \\
& & LU^\infty \wedge \Sigma S(LV)^+.
\end{array}$$

(iv) This is a special case of 2.23 (i).

(v) By construction $h_V(F) = \delta(p, q)$, $h_V(G) = \delta(r, s)$ with

$$\begin{aligned}
p &= (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge \Delta_X), \quad q = (1 \wedge \Delta_Y)(1 \wedge F) : \\
&\quad LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y, \\
r &= (\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge \Delta_Y), \quad s = (1 \wedge \Delta_Z)(1 \wedge G) : \\
&\quad LV^\infty \wedge V^\infty \wedge Y \rightarrow LV^\infty \wedge V^\infty \wedge Z \wedge Z.
\end{aligned}$$

Now

$$\begin{aligned}
(\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge 1)q &= (\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge 1)\Delta_Y(1 \wedge F) \\
&= r(1 \wedge F) : \\
&\quad LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Z \wedge Z,
\end{aligned}$$

so Proposition 2.23 (v) can be applied to give a $(\mathbb{Z}_2$ -equivariant) homotopy

$$\begin{aligned}
h_V(GF) &= \delta((\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge 1)p, s(1 \wedge F)) \\
&\simeq \delta(r(1 \wedge F), s(1 \wedge F),) \\
&\quad + \delta((\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge 1)p, (\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge 1)q) \\
&= h_V(G)(1 \wedge F) + (\kappa_V^{-1} \wedge 1)(G \wedge G)(\kappa_V \wedge 1)h_V(F) : \\
&\quad \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Z \wedge Z.
\end{aligned}$$

(vi) Consider first the special case of a product map of the type

$$F \wedge 1_Z : V^\infty \wedge X \wedge Z \rightarrow V^\infty \wedge Y \wedge Z$$

when the product formula to be obtained is

$$\begin{aligned} h_V(F \wedge 1_Z) &= h_V(F) \wedge \Delta_Z : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge X \wedge Z &\rightarrow LV^\infty \wedge V^\infty \wedge ((Y \wedge Z) \wedge (Y \wedge Z)) . \end{aligned}$$

By construction $h_V(F) = \delta(p_F, q_F)$ with

$$\begin{aligned} p_F &= (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge \Delta_X) , \quad q_F = (1 \wedge \Delta_Y)(1 \wedge F) : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge X &\rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y . \end{aligned}$$

It now follows from

$$\Delta_{X \wedge Z} = \Delta_X \wedge \Delta_Z , \quad \Delta_{Y \wedge Z} = \Delta_Y \wedge \Delta_Z$$

that

$$\begin{aligned} h_V(F \wedge 1) &= \delta(p_{F \wedge 1}, q_{F \wedge 1}) = \delta(p_F \wedge \Delta_Z, q_F \wedge \Delta_Z) \\ &= \delta(p_F, q_F) \wedge \Delta_Z = h_V(F) \wedge \Delta_Z : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge X \wedge Z &\rightarrow LV^\infty \wedge V^\infty \wedge ((Y \wedge Z) \wedge (Y \wedge Z)) . \end{aligned}$$

The general case follows from the suspension formula (iii), the composition formula (v) and the special case, because

$$F_1 \wedge F_2 = (1 \wedge F_2) \circ (F_1 \wedge 1) : (V_1^\infty \wedge X_1) \wedge (V_2^\infty \wedge X_2) \rightarrow (V_1^\infty \wedge Y_1) \wedge (V_2^\infty \wedge Y_2) .$$

(vii) By construction.

(viii) The Hurewicz map

$$\begin{aligned} &[\Sigma S(LV)^+ \wedge V^\infty, (V^\infty \vee V^\infty) \wedge (V^\infty \vee V^\infty)]_{\mathbb{Z}_2} \rightarrow \\ &\rightarrow H_0(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(\dot{C}(\Sigma S(LV)^+ \wedge V^\infty), \dot{C}((V^\infty \vee V^\infty) \wedge (V^\infty \vee V^\infty)))) \\ &= Q_0^V(\mathbb{Z} \oplus \mathbb{Z}) = Q_0^V(\mathbb{Z}) \oplus Q_0^V(\mathbb{Z}) \oplus H_0((\mathbb{Z} \oplus \mathbb{Z}) \otimes (\mathbb{Z} \oplus \mathbb{Z})) = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}^4 \end{aligned}$$

is an isomorphism and the symmetrization map

$$\begin{aligned} 1 + T &= 2 \oplus 2 \oplus 1 : Q_0^V(\mathbb{Z} \oplus \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}^4 \\ &\rightarrow Q_V^0(\mathbb{Z} \oplus \mathbb{Z}) = Q_V^0(\mathbb{Z}) \oplus Q_V^0(\mathbb{Z}) \oplus H_0((\mathbb{Z} \oplus \mathbb{Z}) \otimes (\mathbb{Z} \oplus \mathbb{Z})) = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}^4 \end{aligned}$$

is an injection with

$$\begin{aligned}
(1+T)h_V(\nabla) &= \nabla^{\%}\dot{\phi}_V(S^0) - \dot{\phi}_V(S^0 \vee S^0)\nabla \\
&= (0, 0, (1, 0) \otimes (0, 1) + (0, 1) \otimes (1, 0)) \\
&= (1+T)i_V j_V \in Q_V^0(\mathbb{Z} \oplus \mathbb{Z}) .
\end{aligned}$$

(In Example 6.33 we shall give a more geometric proof of $h_V(\nabla) = i_V j_V$ using the fact that ∇ is an Umkehr map of an immersion $\{*_1, *_2\} \rightarrow \{*\}$ of 0-dimensional manifolds with a single double point.)

(ix) By definition, $F + G$ is the composite

$$\begin{aligned}
F + G &= (F \vee G)(\nabla_V \wedge 1) : \\
V^\infty \wedge X &\xrightarrow{\nabla_V \wedge 1} (V^\infty \vee V^\infty) \wedge X = V^\infty \wedge (X \vee X) \xrightarrow{F \vee G} V^\infty \wedge Y
\end{aligned}$$

with $\nabla_V : V^\infty \rightarrow V^\infty \vee V^\infty$ a sum map as in Definition 2.12 (iii). By the composition formula (v), the product formula (vi), the one-point union formula (vii) and the sum map formula (viii)

$$\begin{aligned}
&h_V(F + G) \\
&= h_V(F \vee G)(\nabla_V \wedge 1) + ((F \vee G) \wedge (F \vee G))h_V(\nabla_V \wedge 1) \\
&= (h_V(F) \vee h_V(G))(\nabla_V \wedge 1) + ((F \vee G) \wedge (F \vee G))(h_V(\nabla_V) \wedge \Delta_X) \\
&= h_V(F) + h_V(G) + ((F \vee G) \wedge (F \vee G))(h_V(\nabla_V) \wedge \Delta_X) \\
&= h_V(F) + h_V(G) + ((F \wedge G) \vee (G \vee F))(j_V \wedge \Delta_X) : \\
&\quad \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge (Y \wedge Y) .
\end{aligned}$$

□

Remark 5.34. As a special case of the composition formula 5.33 (v), we have the naturality of the geometric Hopf invariant for X and Y in the homotopy category, that is

$$h_V((1 \wedge g) \circ F \circ (1 \wedge f)) = (1 \wedge (g \wedge g)) \circ h_V(F) \circ (1 \wedge f)$$

for maps $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$, $f : X' \rightarrow X$ and $g : Y \rightarrow Y'$. □

In the case $X = Y = S^0$ we have:

Proposition 5.35. *The stable \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant of a stable map $F : V^\infty \rightarrow V^\infty$ of degree $d \in \mathbb{Z}$ is*

$$h_V(F) = \frac{d(d-1)}{2} \in \{\Sigma S(LV)^+ \wedge V^\infty; LV^\infty \wedge V^\infty\}_{\mathbb{Z}_2} = \{S^0; P(V)^+\} = \mathbb{Z} .$$

Proof. We offer three proofs:

(I) The maps $p, q : LV^\infty \wedge V^\infty \rightarrow LV^\infty \wedge V^\infty$ used to define $h_V(F) = \delta(p, q)$ have

$$\text{degree}(p) = d^2, \text{degree}(q) = d \in \mathbb{Z}$$

so that by Proposition 4.21

$$\begin{aligned} h_V(F) &= \text{semidegree}(h_V(F)) \\ &= \frac{\text{degree}(p) - \text{degree}(q)}{2} \\ &= \frac{d^2 - d}{2} \in [\Sigma S(LV)^+ \wedge V^\infty, LV^\infty \wedge V^\infty]_{\mathbb{Z}_2} = \mathbb{Z}. \end{aligned}$$

(II) As in the proof of Proposition 5.33 (ix) note that the Hurewicz map

$$\begin{aligned} [\Sigma S(LV)^+ \wedge V^\infty, V^\infty \wedge V^\infty]_{\mathbb{Z}_2} \rightarrow \\ H_0(\text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(\dot{C}(\Sigma S(LV)^+ \wedge V^\infty), \dot{C}(V^\infty \wedge V^\infty))) = Q_0^V(\mathbb{Z}) = \mathbb{Z} \end{aligned}$$

is an isomorphism and the symmetrization map

$$1 + T = 2 : Q_0^V(\mathbb{Z}) = \mathbb{Z} \rightarrow Q_V^0(\mathbb{Z}) = \mathbb{Z}$$

is an injection with

$$(1 + T)h_V(d) = d^2 \dot{\phi}_V(S^0) - \dot{\phi}_V(S^0)d = d^2 - d \in Q_V^0(\mathbb{Z}) = \mathbb{Z}.$$

(III) Proposition 4.41 (i) and Example 4.80 (v) give a split short exact sequence

$$\begin{aligned} \{S^1; LV^\infty\}_{\mathbb{Z}_2} = 0 \rightarrow \{\Sigma S(LV)^+; LV^\infty\}_{\mathbb{Z}_2} &= \{S^0; P(V)^+\}_{\mathbb{Z}_2} = \mathbb{Z} \\ \rightarrow \{LV^\infty; LV^\infty\}_{\mathbb{Z}_2} &= \{S^0; S^0\}_{\mathbb{Z}_2} = A(\mathbb{Z}_2) \\ &\xrightarrow{\rho} \{S^0; LV^\infty\}_{\mathbb{Z}_2} = \{S^0; S^0\} = \mathbb{Z} \rightarrow 0 \end{aligned}$$

with $A(\mathbb{Z}_2)$ the Burnside ring of finite \mathbb{Z}_2 -sets and ρ the \mathbb{Z}_2 -fixed point map. Assume that F is transverse regular at $0 \in V^\infty$, so that $\{x \in V \mid F(x) = 0\}$ is a finite set with d points (counted algebraically), with

$$F = d \in [V^\infty, V^\infty] = \mathbb{Z}.$$

The image of $h_V(F)$ in $A(\mathbb{Z}_2)$ is the class of the finite free \mathbb{Z}_2 -set

$$S = \{(x, y) \in V \times V \mid F(x) = F(y) = 0, x \neq y\}$$

with $d^2 - d = d(d - 1)$ points, so that

$$h_V(F) = \frac{|S|}{2} = \frac{d(d - 1)}{2} \in \mathbb{Z}.$$

□

Example 5.36. (i) Given $c \in O(V)$ define

$$d = \text{degree}(c^\infty : V^\infty \rightarrow V^\infty) = \begin{cases} +1 & \text{if } \det(c) > 0 \\ -1 & \text{if } \det(c) < 0 . \end{cases}$$

By Proposition 5.35 the stable \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant of $c^\infty : V^\infty \rightarrow V^\infty$ is given by

$$\begin{aligned} h_V(c^\infty) &= \frac{d(d-1)}{2} = \begin{cases} 0 & \text{if } \det(c) > 0 \\ 1 & \text{if } \det(c) < 0 \end{cases} \\ &\in \omega_0(P(\mathbb{R}(\infty))) = \ker(\rho : \omega_{0,0} \rightarrow \omega_0) = \mathbb{Z} \end{aligned}$$

(using the terminology of Example 4.80).

(ii) In the special case of (i)

$$\begin{aligned} V &= \mathbb{R}, c = -1 \in GL(V) = \mathbb{R} \setminus \{0\}, \\ c^\infty &= -1 : V^\infty = S^1 \rightarrow V^\infty = S^1 ; t \mapsto 1 - t \end{aligned}$$

with degree $d = -1$ the formula in (i) gives that

$$h_{\mathbb{R}}(-1) = 1 \in \mathbb{Z} .$$

□

Example 5.37. Let $V = \mathbb{R}$, $X = S^{2n}$, $Y = S^n$, for any $n \geq 0$. The \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant of a map

$$F : V^\infty \wedge X = S^{2n+1} \rightarrow V^\infty \wedge Y = S^{n+1}$$

is an integer

$$\begin{aligned} h_{\mathbb{R}}(F) &\in [\Sigma S(LV)^+ \wedge V^\infty \wedge X, LV^\infty \wedge V^\infty \wedge Y \wedge Y]_{\mathbb{Z}_2} \\ &= [V^\infty \wedge V^\infty \wedge X, LV^\infty \wedge V^\infty \wedge Y \wedge Y] = \pi_{2n+2}(S^{2n+2}) = \mathbb{Z} \end{aligned}$$

(using $\Sigma S(LV)^+ = V^\infty \vee LV^\infty$), namely the degree of the map

$$h_{\mathbb{R}}(F)| : V^\infty \wedge V^\infty \wedge X = S^{2n+2} \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y = S^{2n+2} .$$

(i) For $n = 0$ and any $d \in \mathbb{Z}$ the stable \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant of the map

$$d : \Sigma X = S^1 \rightarrow \Sigma Y = S^1 ; t \mapsto dt$$

is given by Proposition 5.35 to be

$$h_{\mathbb{R}}(d) = \frac{d(d-1)}{2} \in \mathbb{Z} ,$$

See 6.35 below for a geometric interpretation in the case $d \geq 1$.

(ii) See 6.68 below for the identification of the \mathbb{Z}_2 -equivariant homotopy class of the geometric Hopf invariant of a map $F : \Sigma X = S^{2n+1} \rightarrow \Sigma Y = S^{n+1}$ in the case $n \geq 1$ with the classical Hopf invariant of F . \square

The geometric Hopf invariant is closely related to the relative version of the symmetric construction, for maps:

Proposition 5.38. (i) *Given a map $F : X \rightarrow Y$ let*

$$Z = \mathcal{C}(F) = Y \cup_F CX$$

be the mapping cone, so that there is defined a cofibration sequence

$$X \xrightarrow{F} Y \xrightarrow{G} Z \xrightarrow{H} \Sigma X$$

with G the inclusion, H the projection. Let $E : CX \rightarrow Z$ be the inclusion, a canonical null-homotopy of $GF : X \rightarrow Z$. For any inner product space V the V -coefficient symmetric construction $\dot{\phi}_V(Z)$ is the composite \mathbb{Z}_2 -equivariant map

$$\dot{\phi}_V(Z) : S(LV)^+ \wedge Z \xrightarrow{1 \wedge H} \Sigma S(LV)^+ \wedge X \xrightarrow{\delta(r, Tr)} Z \wedge Z$$

with

$$r = (E \wedge GF)\phi_V(X) , \quad Tr = (GF \wedge E)\phi_V(X) : CS(LV)^+ \wedge X \rightarrow Z \wedge Z$$

two null-homotopies of

$$(GF \wedge GF)\phi_V(X) : S(LV)^+ \wedge X \rightarrow Z \wedge Z .$$

(ii) *The geometric Hopf invariant of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is the relative difference (1.3) of the null-homotopies of*

$$\begin{aligned} (F \wedge F)\dot{\phi}_V(V^\infty \wedge X) &= \dot{\phi}_V(V^\infty \wedge Y)(1 \wedge F) : \\ S(LV)^+ \wedge V^\infty \wedge X &\rightarrow V^\infty \wedge V^\infty \wedge Y \wedge Y \end{aligned}$$

given by 5.26

$$\begin{aligned} & \delta \dot{\phi}_V(V^\infty \wedge Y)(1 \wedge F) , (F \wedge F) \delta \dot{\phi}_V(V^\infty \wedge X) : \\ & CS(LV)^+ \wedge V^\infty \wedge X \rightarrow V^\infty \wedge V^\infty \wedge Y \wedge Y \end{aligned}$$

that is

$$\begin{aligned} h_V(F) &= (\kappa_V^{-1} \wedge 1) \delta(\delta \dot{\phi}_V(V^\infty \wedge Y)(1 \wedge F), (F \wedge F) \delta \dot{\phi}_V(V^\infty \wedge X)) : \\ & \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y . \end{aligned}$$

(iii) Given a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ let

$$V^\infty \wedge X \xrightarrow{F} V^\infty \wedge Y \xrightarrow{G} Z \xrightarrow{H} \Sigma V^\infty \wedge X$$

be as in (i) with $Z = \mathcal{C}(F)$ the mapping cone, G the inclusion and H the projection. The V -coefficient symmetric construction $\dot{\phi}_V(Z)$ is given up to \mathbb{Z}_2 -equivariant homotopy by

$$\dot{\phi}_V(Z) = (G \wedge G) h_V(F) H : S(LV)^+ \wedge Z \rightarrow Z \wedge Z .$$

Proof. (i) By construction.

(ii) By Proposition 2.20 (i).

(iii) Combine (ii) and Proposition 1.11 (ii). □

The geometric Hopf invariant $h_V(F)$ has a particularly simple description in the special case $V = \mathbb{R}$.

Definition 5.39. For $V = \mathbb{R}$

$$\begin{aligned} V^\infty &= S^1 , S(LV) = S^0 , LV \setminus \{0\} = \mathbb{R}_+ \sqcup \mathbb{R}_- , \\ \Sigma S(LV)^+ &= (LV \setminus \{0\})^\infty = S^1 \vee S^1 \end{aligned}$$

with \mathbb{Z}_2 acting on S^0 and $S^1 \vee S^1$ by transposition, and

$$\mathbb{R}_+ = \{x \in \mathbb{R} \mid x > 0\} , \mathbb{R}_- = \{x \in \mathbb{R} \mid x < 0\} .$$

The geometric Hopf invariant of a map

$$F : \mathbb{R}^\infty \wedge X \rightarrow \Sigma X \rightarrow \mathbb{R}^\infty \wedge Y \rightarrow \Sigma Y$$

is the \mathbb{Z}_2 -equivariant map

$$h_{\mathbb{R}}(F) = h_{\mathbb{R}}(F)_+ \vee h_{\mathbb{R}}(F)_- : (S^1 \vee S^1) \wedge \Sigma X \rightarrow L\mathbb{R}^\infty \wedge \mathbb{R}^\infty \wedge Y \wedge Y = \Sigma Y \wedge \Sigma Y$$

which is entirely determined by the non-equivariant \mathbb{R} -coefficient geometric Hopf invariant

$$h_{\mathbb{R}}(F)_+ : (S^1 \vee \{*\}) \wedge \Sigma X = \Sigma^2 X \rightarrow \Sigma Y \wedge \Sigma Y .$$

□

Proposition 5.40. (i) *The geometric Hopf invariant of the sum of maps $F, G : \Sigma X \rightarrow \Sigma Y$*

$$F + G : \Sigma X \rightarrow \Sigma Y ; (s, x) \mapsto \begin{cases} F(2s, x) & \text{if } 0 \leq s \leq 1/2 \\ G(2s - 1, x) & \text{if } 1/2 \leq s \leq 1 \end{cases}$$

is given up to homotopy by

$$h_{\mathbb{R}}(F + G)_+ = h_{\mathbb{R}}(F)_+ + h_{\mathbb{R}}(G)_+ + F \wedge G : \Sigma^2 X \rightarrow \Sigma Y \wedge \Sigma Y .$$

(ii) *The geometric Hopf invariant of the map defined for any pointed spaces A_1, A_2 by*

$$F : \Sigma(A_1 \times A_2) \rightarrow \Sigma(A_1 \vee A_2) ; (s, a_1, a_2) \mapsto \begin{cases} (2s, a_1) & \text{if } 0 \leq s \leq 1/2 \\ (2s - 1, a_2) & \text{if } 1/2 \leq s \leq 1 \end{cases}$$

is given up to homotopy by

$$h_{\mathbb{R}}(F)_+ : \Sigma^2(A_1 \times A_2) \xrightarrow{\Sigma^2 p} \Sigma^2(A_1 \wedge A_2) \subset \Sigma(A_1 \vee A_2) \wedge \Sigma(A_1 \vee A_2)$$

with $p : A_1 \times A_2 \rightarrow A_1 \wedge A_2$ the projection.

Proof. (i) This is the special case $V = \mathbb{R}$ of the sum formula of Proposition 5.33 (ix).

(ii) Apply (i) to $F = \Sigma F_1 + \Sigma F_2$, with

$$F_i : A_1 \times A_2 \rightarrow A_1 \vee A_2 ; (a_1, a_2) \mapsto a_i .$$

□

Example 5.41. For any pointed space X define the map

$$F = \Sigma p_1 + \Sigma p_2 : \Sigma(X \times X) \rightarrow \Sigma X$$

with

$$p_i : X \times X \rightarrow X ; (x_1, x_2) \mapsto x_i \quad (i = 1, 2) .$$

By Proposition 5.40 the geometric Hopf invariant of F is given up to homotopy by

$$h_{\mathbb{R}}(F)_+ = \Sigma^2 p : \Sigma^2(X \times X) \rightarrow \Sigma^2(X \wedge X)$$

with $p : X \times X \rightarrow X \wedge X$ the projection. \square

5.4 The stable geometric Hopf invariant $h'_V(F)$

The stable relative difference $\delta'(p, q)$ defined in 3.6 above will now be used to construct a second form of the geometric Hopf invariant of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$, to be a \mathbb{Z}_2 -equivariant map

$$h'_V(F) : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge S(LV)^+ \wedge Y \wedge Y$$

with LV as defined in section 5.3. We shall regard $h'_V(F)$ as a stable \mathbb{Z}_2 -equivariant map

$$h'_V(F) : X \rightarrow S(LV)^+ \wedge Y \wedge Y .$$

The symmetric construction on the mapping cone of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ will be shown to factorize (up to \mathbb{Z}_2 -equivariant homotopy) through the stable geometric Hopf invariant $h'_V(F)$ of §5.4. Thus $h'_V(F)$ is a geometric version of the functional Steenrod squares (cf. Example 5.51 below).

As before, define the \mathbb{Z}_2 -equivariant maps

$$\begin{aligned} p &= (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge \Delta_X) , \quad q = (1 \wedge \Delta_Y)(1 \wedge F) : \\ &LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y \end{aligned}$$

which agree on $0^+ \wedge V^\infty \wedge X = V^\infty \wedge X \subset LV^\infty \wedge V^\infty \wedge X$. The geometric Hopf invariant (5.30) is the \mathbb{Z}_2 -equivariant map defined by

$$h_V(F) = \delta(p, q) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y .$$

Definition 5.42. The *stable geometric Hopf invariant* of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is the stable relative difference of p and q (3.6), the \mathbb{Z}_2 -equivariant map

$$h'_V(F) = \delta'(p, q) : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge S(LV)^+ \wedge Y \wedge Y$$

which we shall regard as a stable \mathbb{Z}_2 -equivariant map

$$h'_V(F) : X \rightarrow S(LV)^+ \wedge Y \wedge Y .$$

□

Proposition 5.43. (i) *The \mathbb{Z}_2 -equivariant S -duality isomorphism*

$$\{\Sigma S(LV)^+ \wedge V^\infty \wedge X; LV^\infty \wedge V^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2} \rightarrow \{X; S(LV)^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2} ;$$

$$f \mapsto (1 \wedge f)(\Delta_{\alpha_{LV}} \wedge 1)$$

sends the geometric Hopf invariant map

$$h_V(F) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y$$

to the stable geometric Hopf invariant map $h'_V(F)$.

(ii) *The stable geometric Hopf invariant is such that there is a commutative diagram of \mathbb{Z}_2 -equivariant maps*

$$\begin{array}{ccc} LV^\infty \wedge V^\infty \wedge X & \xrightarrow{h'_V(F)} & LV^\infty \wedge V^\infty \wedge S(LV)^+ \wedge Y \wedge Y \\ & \searrow 1 \wedge ((\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge \Delta_X) - \Delta_Y(1 \wedge F)) & \downarrow s_{LV} \wedge 1 \\ & & LV^\infty \wedge V^\infty \wedge Y \wedge Y \end{array}$$

with

$$s_{LV} : S(LV)^+ \rightarrow S(LV)^+/S(LV) = 0^+ ; u \mapsto 0$$

the \mathbb{Z}_2 -equivariant pinch map.

(iii) *The stable geometric Hopf invariant defines a function*

$$h : [V^\infty \wedge X; V^\infty \wedge Y] \rightarrow \{X; S(LV)^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2} ; F \mapsto h'_V(F)$$

such that for any $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$, $G : V^\infty \wedge Y \rightarrow V^\infty \wedge Z$

$$h'_V(GF) = h'_V(G)F + (G \wedge G)h'_V(F) : X \rightarrow S(LV)^+ \wedge Z \wedge Z .$$

Proof. (i) This is a special case of 3.44 (i).

(ii) This is a special case of 3.44 (ii).

(iii) By construction and the composition formula of 5.33 (v). □

Proposition 4.41 gives a commutative braid of exact sequences of stable \mathbb{Z}_2 -equivariant homotopy groups

$$\begin{array}{ccccc}
& & \xrightarrow{\quad} & & \\
\{X; S(LV)^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2} & & \{S(LV)^+ \wedge X; Y \wedge Y\}_{\mathbb{Z}_2} & & \\
\downarrow s_{LV} & & \uparrow s_{LV}^* & & \searrow \\
& \{X; Y \wedge Y\}_{\mathbb{Z}_2} & & \{S(LV \oplus LV)^+ \wedge X; LV^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2} & \\
\uparrow 0_{LV}^* & & \downarrow 0_{LV} & & \searrow \\
\{LV^\infty \wedge X; Y \wedge Y\}_{\mathbb{Z}_2} & & \{X; LV^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2} & & \\
& \xrightarrow{\quad} & & & \searrow \\
& & & & A
\end{array}$$

and for a pointed map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ with $V \subseteq \mathbb{R}(\infty)$

$$\begin{aligned}
p &= (F \wedge F) \Delta_X = (h'_{\mathbb{R}(\infty)}(F), F), \quad q = \Delta_Y F = (0, F) \\
&\in \{X; Y \wedge Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge (Y \wedge Y)\}_{\mathbb{Z}_2} \oplus \{X; Y\}, \\
s_{LV}^*(p) &= (F \wedge F) \dot{\phi}_V(X), \quad s_{LV}^*(q) = \dot{\phi}_V(Y) F \in \{S(LV)^+ \wedge X; Y \wedge Y\}_{\mathbb{Z}_2}, \\
0_{LV}(p) &= 0_{LV}(q) = (0, F) \\
&\in \{V^\infty \wedge X; (V^\infty \wedge Y) \wedge (V^\infty \wedge Y)\}_{\mathbb{Z}_2} = \{X; LV^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2} \\
&= \{X; S(\infty)/S(LV) \wedge (Y \wedge Y)\}_{\mathbb{Z}_2} \oplus \{X; Y\}, \\
p - q &= s_{LV} h'_V(F) \in \ker(0_{LV} : \{X; Y \wedge Y\}_{\mathbb{Z}_2} \rightarrow \{X; LV^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2}) \\
&= \text{im}(s_{LV} : \{X; S(LV)^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2} \rightarrow \{X; Y \wedge Y\}_{\mathbb{Z}_2}), \\
\{LV^\infty \wedge X; Y \wedge Y\}_{\mathbb{Z}_2} &= \{S(\infty)/S(LV) \wedge X; Y \wedge Y\}_{\mathbb{Z}_2} \oplus \{X; Y\}, \\
A &= \{S(LV)^+ \wedge X; LV^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2} = \{X; \Sigma S(LV)^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2}.
\end{aligned}$$

For any pointed space X there is defined a split short exact sequence

$$0 \longrightarrow \{X; P(\mathbb{R}(\infty))^+\} \longrightarrow \tilde{\omega}_{\mathbb{Z}_2}^0(X) \xrightarrow{\rho} \tilde{\omega}^0(X) \longrightarrow 0$$

with ρ the \mathbb{Z}_2 -fixed point map and

$$\begin{aligned}
\sigma &: \tilde{\omega}^0(X) \rightarrow \tilde{\omega}_{\mathbb{Z}_2}^0(X); \\
(F : V^\infty \wedge X \rightarrow V^\infty) &\mapsto (1 \wedge F : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty)
\end{aligned}$$

such that $\rho\sigma = 1 : \tilde{\omega}^0(X) \rightarrow \tilde{\omega}^0(X)$ (Example 4.44 (iv)).

Definition 5.44. (Crabb [12, pp.32-33])

(i) The *squaring operation* from stable cohomotopy to stable \mathbb{Z}_2 -cohomotopy

$$\begin{aligned}
P^2 &: \omega^0(X) \rightarrow \omega_{\mathbb{Z}_2}^0(X); \quad (F : V^\infty \wedge X^+ \rightarrow V^\infty) \mapsto \\
&((\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge 1)(1 \wedge \Delta_X) : LV^\infty \wedge V^\infty \wedge X^+ \rightarrow LV^\infty \wedge V^\infty)
\end{aligned}$$

is a non-additive function such that $\rho P^2 = 1 : \omega^0(X) \rightarrow \omega^0(X)$.

(ii) The *reduced squaring operation* is defined by

$$\begin{aligned} \overline{P}^2 : \omega^0(X) &\rightarrow \ker(\rho : \omega_{\mathbb{Z}_2}^0(X \times X) \rightarrow \omega^0(X)) = \{X; P(\mathbb{R}(\infty))^+\} ; \\ (F : V^\infty \wedge X^+ \rightarrow V^\infty) &\mapsto P^2(F) - \sigma(F) = h'_V(F) . \end{aligned}$$

□

Example 5.45. (i) \overline{P}^2 agrees with the Hopf invariant function θ^2 of Segal [57] (cf. Crabb [12, p.35]).

(ii) For $j \equiv 1 \pmod{2}$ the composite of \overline{P}^2 for $X = S^j$ and the Hurewicz map

$$\omega_j = \tilde{\omega}^0(S^j) \xrightarrow{\overline{P}^2} \{S^j; P(\mathbb{R}(\infty))^+\} = \omega_j(P(\mathbb{R}(\infty))) \longrightarrow H_j(P(\mathbb{R}(\infty))) = \mathbb{Z}_2$$

is the classical mod 2 Hopf invariant ([12, 4.16] and Example 5.20 (iv)).

(iii) For $j \equiv 3 \pmod{4}$ the composite of \overline{P}^2 for $X = S^j$ and the real KO_* -theory Hurewicz map

$$\begin{aligned} \omega_j = \tilde{\omega}^0(S^j) &\xrightarrow{\overline{P}^2} \{S^j; P(\mathbb{R}(\infty))^+\} = \omega_j(P(\mathbb{R}(\infty))) \\ &\longrightarrow KO_j(P(\mathbb{R}(\infty))) = \mathbb{Z}[1/2]/\mathbb{Z} \end{aligned}$$

is the 2-primary e -invariant ([12, 4.17]).

□

5.5 The quadratic construction $\psi_V(F)$

The V -coefficient ‘quadratic construction’ on a pointed space X is the pointed space $S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X)$.

Definition 5.46. Let V be an inner product space, and let $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ be a pointed map.

(i) The *space level V -coefficient quadratic construction* on F is the stable map

$$\psi_V(F) : X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)$$

given by the image of the \mathbb{Z}_2 -equivariant stable geometric Hopf invariant (5.42)

$$h'_V(F) = \delta'(p, q) : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge S(LV)^+ \wedge Y \wedge Y$$

under the isomorphism given by Proposition 4.32

$$\{X; S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\} \cong \{X; S(LV)^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2}.$$

(ii) The *chain level V -coefficient quadratic construction* on F is the \mathbb{Z} -module chain map induced by the space level $\psi_V(F)$

$$\psi_V(F) : \dot{C}(X) \rightarrow C^{cell}(S(LV)) \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (\dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(Y)),$$

inducing morphisms

$$\psi_V(F) : \dot{H}_n(X) \rightarrow Q_n^V(\dot{C}(Y)).$$

More directly, note that the chain level quadratic construction $\psi_V(F)$ is the adjoint of the $\mathbb{Z}[\mathbb{Z}_2]$ -module chain map

$$h_V(F) : SC^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(X) \rightarrow C^{cell}(LV^\infty) \otimes_{\mathbb{Z}} (\dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(Y))$$

induced by the geometric Hopf invariant \mathbb{Z}_2 -equivariant map

$$h_V(F) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y,$$

with respect to the chain level \mathbb{Z}_2 -equivariant S -duality isomorphism of 4.81 (ii) and 5.9 (ii)

$$\begin{aligned} & C^{cell}(S(LV)) \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (\dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(Y)) \\ & \cong \text{Hom}_{\mathbb{Z}[\mathbb{Z}_2]}(SC^{cell}(S(LV)), \dot{C}(LV^\infty) \otimes_{\mathbb{Z}} (\dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(Y))). \end{aligned}$$

□

Remark 5.47. We refer to Crabb and James [13] for an extended treatment of fibrewise homotopy theory - there is a brief account in Appendix A below. Here is an explicit formula for the space level quadratic construction $\psi_V(F)$ in terms of fibrewise \mathbb{Z}_2 -equivariant homotopy theory. Think of the Hopf invariant map

$$h_V(F) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y$$

as a fibrewise \mathbb{Z}_2 -equivariant map

$$S(LV) \times (\Sigma V^\infty \wedge X) \rightarrow S(LV) \times (LV^\infty \wedge V^\infty \wedge Y \wedge Y)$$

over $S(LV)$. By dividing out the free \mathbb{Z}_2 -action, we get a fibrewise map

$$P(V) \times (\Sigma V^\infty \wedge X) \rightarrow S(LV) \times_{\mathbb{Z}_2} (LV^\infty \wedge V^\infty \wedge Y \wedge Y) \quad (*)$$

over the real projective space $P(V)$.

We now use fibrewise duality theory as in Crabb and James [13, Part II, (14.43)]. This requires an embedding of $P(V)$ in some Euclidean space. To be definite, we use the embedding

$$P(V) \hookrightarrow M = \text{End}(V) ; [x] \mapsto (v \mapsto \langle x, v \rangle x) \quad (x \in S(LV))$$

mapping a line to the orthogonal projection onto that subspace. Use the inner product to construct a tubular neighbourhood $\nu \hookrightarrow M$, where ν is the (total space of) the normal bundle. Notice that the direct sum $\mathbb{R} \oplus \tau$ of a trivial line and the tangent bundle τ of $P(V)$ is just $S(LV) \times_{\mathbb{Z}_2} LV$. And $\tau \oplus \nu$ is the trivial bundle $P(V) \times M$.

So the fibrewise smash product of $(*)$ with the identity on the fibrewise one-point compactification $\nu_{P(V)}^\infty$ of ν gives a fibrewise map

$$\begin{aligned} \nu_{P(V)}^\infty \wedge_{P(V)} (P(V) \times (\Sigma V^\infty \wedge X)) &\rightarrow \\ (P(V) \times M^+) \wedge_{P(V)} (S(LV) \times_{\mathbb{Z}_2} (\Sigma V^\infty \wedge Y \wedge Y)) & \end{aligned}$$

over $P(V)$. Collapsing the basepoint section $P(V)$, we get a map

$$P(V)^\nu \wedge (\Sigma V^\infty \wedge X) \rightarrow (M^+ \wedge \Sigma V^\infty) \wedge (S(LV)^+ \wedge_{\mathbb{Z}_2} Y \wedge Y)$$

Compose this with the Pontryagin-Thom map $M^\infty \rightarrow P(V)^\nu$, and we have produced an explicit map

$$\psi_V(F) : (M^+ \wedge \Sigma V^\infty) \wedge X \rightarrow (M^+ \wedge \Sigma V^\infty) \wedge (S(LV)^+ \wedge_{\mathbb{Z}_2} Y \wedge Y) .$$

□

Proposition 5.48. *Let $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ be a stable map, with $V = \mathbb{R}^k$. The quadratic construction $\psi_V(F) : X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)$ induces the chain level quadratic construction of Ranicki [50, §1], [51, p.29]*

$$\begin{aligned} \psi_V(F) : \dot{C}(X) &\rightarrow \dot{C}^{cell}(S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)) \\ &= W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (\dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(Y)) \end{aligned}$$

and hence also the quadratic construction on the level of homology groups

$$\psi_V(F) : \dot{H}_*(X) \rightarrow \dot{H}_*(S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)) = Q_*^{[0, k-1]}(\dot{C}(Y)) ,$$

identifying $C^{cell}(S(LV)) = W[0, k-1]$.

Proof. From Proposition 5.27 we have a diagram of $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes, chain maps and chain homotopies

$$\begin{array}{ccc}
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(V^\infty) \otimes_{\mathbb{Z}} \dot{C}(X) & \xrightarrow{\dot{\phi}_V(V^\infty) \cup \dot{\phi}_V(X)} & \dot{C}(V^\infty) \otimes_{\mathbb{Z}} \dot{C}(X) \otimes_{\mathbb{Z}} \dot{C}(V^\infty) \otimes_{\mathbb{Z}} \dot{C}(X) \\
1 \otimes \dot{E}(V^\infty, X) \downarrow \simeq & \nearrow \delta \dot{\phi}_V(V^\infty, X) \quad \dot{E}(V^\infty, X) \otimes \dot{E}(V^\infty, X) \downarrow \simeq & \\
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(V^\infty \wedge X) & \xrightarrow{\dot{\phi}_V(V^\infty \wedge X)} & \dot{C}(V^\infty \wedge X) \otimes_{\mathbb{Z}} \dot{C}(V^\infty \wedge X) \\
1 \otimes F \downarrow & & \downarrow F \otimes F \\
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(V^\infty \wedge Y) & \xrightarrow{\dot{\phi}_V(V^\infty \wedge Y)} & \dot{C}(V^\infty \wedge Y) \otimes_{\mathbb{Z}} \dot{C}(V^\infty \wedge Y) \\
1 \otimes \dot{E}(V^\infty, Y) \uparrow \simeq & \nwarrow \delta \dot{\phi}_V(V^\infty, Y) \quad \dot{E}(V^\infty, Y) \otimes \dot{E}(V^\infty, Y) \uparrow \simeq & \\
C^{cell}(S(LV)) \otimes_{\mathbb{Z}} \dot{C}(V^\infty) \otimes_{\mathbb{Z}} \dot{C}(Y) & \xrightarrow{\dot{\phi}_V(V^\infty) \cup \dot{\phi}_V(Y)} & \dot{C}(V^\infty) \otimes_{\mathbb{Z}} \dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(V^\infty) \otimes_{\mathbb{Z}} \dot{C}(Y)
\end{array}$$

Evaluation on a generator of $\dot{C}(V^\infty) = S^k \mathbb{Z}$ now gives a diagram of $\mathbb{Z}[\mathbb{Z}_2]$ -module chain complexes, chain maps and chain homotopies

$$\begin{array}{ccc}
W[0, k-1] \otimes_{\mathbb{Z}} S^k \dot{C}(X) & \xrightarrow{S^k \dot{\phi}_{\mathbb{R}^k}(X)} & S^k \dot{C}(X) \otimes_{\mathbb{Z}} S^k \dot{C}(X) \\
1 \otimes \dot{E}(S^k, X) \downarrow \simeq & \nearrow \delta \dot{\phi}_{\mathbb{R}^k}(S^k, X) \quad \dot{E}(S^k, X) \otimes \dot{E}(S^k, X) \downarrow \simeq & \\
W[0, k-1] \otimes_{\mathbb{Z}} \dot{C}(\Sigma^k X) & \xrightarrow{\dot{\phi}_{\mathbb{R}^k}(\Sigma^k X)} & \dot{C}(\Sigma^k X) \otimes_{\mathbb{Z}} \dot{C}(\Sigma^k X) \\
1 \otimes F \downarrow & & \downarrow F \otimes F \\
W[0, k-1] \otimes_{\mathbb{Z}} \dot{C}(\Sigma^k Y) & \xrightarrow{\dot{\phi}_{\mathbb{R}^k}(\Sigma^k Y)} & \dot{C}(\Sigma^k Y) \otimes_{\mathbb{Z}} \dot{C}(\Sigma^k Y) \\
1 \otimes \dot{E}(S^k, Y) \uparrow \simeq & \nwarrow \delta \dot{\phi}_{\mathbb{R}^k}(S^k, Y) \quad \dot{E}(S^k, Y) \otimes \dot{E}(S^k, Y) \uparrow \simeq & \\
W[0, k-1] \otimes_{\mathbb{Z}} S^k \dot{C}(Y) & \xrightarrow{S^k \dot{\phi}_{\mathbb{R}^k}(Y)} & S^k \dot{C}(Y) \otimes_{\mathbb{Z}} S^k \dot{C}(Y)
\end{array}$$

The stable geometric Hopf invariant $h'_{\mathbb{R}^k}(F)$ of Proposition 5.42 (ii) induces the same chain map as the quadratic construction $\psi_{\mathbb{R}^k}(F)$, which is thus the chain map given by Proposition 5.13 (ii)

$$\psi_{\mathbb{R}^k}(F) = \delta(F, \dot{\phi}_{\mathbb{R}^k}(X), \dot{\phi}_{\mathbb{R}^k}(Y)) : \dot{C}(X) \rightarrow W[0, k-1] \otimes_{\mathbb{Z}} (\dot{C}(Y) \otimes_{\mathbb{Z}} \dot{C}(X)).$$

But this is exactly the chain level quadratic construction of [50, 51].

□

The quadratic construction $\psi_V(F)$ has the following properties :

Proposition 5.49. (i) *The stable \mathbb{Z}_2 -equivariant homotopy class of $\psi_V(F) : X \rightarrow S(LV)^+ \wedge (Y \wedge Y)$ depends only on the homotopy class of $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$. The function*

$$\psi_V : [V^\infty \wedge X, V^\infty \wedge Y] \rightarrow \{X; S(LV)^+ \wedge (Y \wedge Y)\}_{\mathbb{Z}_2} ; F \mapsto \psi_V(F)$$

is such that

$$\psi_V(F + G) = \psi_V(F) + \psi_V(G) + i((F \wedge G) \circ \Delta_X) ,$$

where

$$i : \{X; Y \wedge Y\} = \{X; (S^0)^+ \wedge (Y \wedge Y)\}_{\mathbb{Z}_2} \rightarrow \{X; S(LV)^+ \wedge (Y \wedge Y)\}_{\mathbb{Z}_2}$$

is induced by the map

$$Y \wedge Y = (S^0)^+ \wedge (Y \wedge Y) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y) ; (\pm 1, y_1, y_2) \mapsto (\pm v, y_1, y_2)$$

for any $v \in S(LV)$.

(ii) The symmetrization of the quadratic construction is the difference of the symmetric constructions, in the sense that $\psi_V(F) \in \{X; S(LV)^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2}$ has images

$$s_{LV}(\psi_V(F)) = (F \wedge F)\Delta_X - \Delta_Y F \in \{X; Y \wedge Y\}_{\mathbb{Z}_2} ,$$

$$s_{LV}^* s_{LV}(\psi_V(F)) = (F \wedge F)\dot{\phi}_V(X) - \dot{\phi}_V(Y)F \in \{S(LV)^+ \wedge X; Y \wedge Y\}_{\mathbb{Z}_2} .$$

(iii) The quadratic construction on the suspension

$$\Sigma F : (V \oplus \mathbb{R})^\infty \wedge X = \Sigma(V^\infty \wedge X) \rightarrow (V \oplus \mathbb{R})^\infty \wedge Y = \Sigma(V^\infty \wedge Y)$$

is the composite

$$\psi_{V \oplus \mathbb{R}}(\Sigma F) : X \xrightarrow{\psi_V(F)} S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y) \xrightarrow{e \wedge 1} S(L(V \oplus \mathbb{R}))^\infty \wedge_{\mathbb{Z}_2} Y \wedge Y$$

with $e : S(LV) \rightarrow S(L(V \oplus \mathbb{R}))$ the inclusion induced by

$$e : V \rightarrow V \oplus \mathbb{R} ; x \mapsto (x, 0) .$$

(iv) If $V = \{0\}$ then $\psi_V(F) = 0$, i.e. the quadratic construction is 0 for an unstable map $F : X \rightarrow Y$.

(v) (Sum formula) The sum of maps $F_1, F_2 : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ (with $\dim(V) > 0$) is a map

$$F_1 + F_2 : V^\infty \wedge X \rightarrow V^\infty \wedge Y$$

with quadratic construction

$$\begin{aligned} \psi_V(F_1 + F_2) &= \psi_V(F_1) + \psi_V(F_2) + (F_1 \wedge F_2 + F_2 \wedge F_1)\Delta_X \\ &: X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y) \end{aligned}$$

with

$$F_1 \wedge F_2 + F_2 \wedge F_1 : X \wedge X \xrightarrow{\begin{pmatrix} F_1 \wedge F_2 \\ F_2 \wedge F_1 \end{pmatrix}} \\ (Y \wedge Y) \vee (Y \wedge Y) = (S^0)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y) \hookrightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)$$

where $S^0 \subset S(LV)$; $\pm \mapsto \pm v$ for any $v \in S(LV)$.

(vi) (Product formula) Let $F_i : V_i^\infty \wedge X_i \rightarrow V_i^\infty \wedge Y_i$, $i = 1, 2$, be two maps. The quadratic construction $\psi_{V_1 \oplus V_2}(F_1 \wedge F_2)$ of the smash product

$$F_1 \wedge F_2 : (V_1 \oplus V_2)^\infty \wedge (X_1 \wedge X_2) \rightarrow (V_1 \oplus V_2)^\infty \wedge (Y_1 \wedge Y_2)$$

is the sum

$$\begin{aligned} & \psi_{V_1 \oplus V_2}(F_1 \wedge F_2) \\ &= (a_1 \wedge 1)(\psi_{V_1}(F_1) \wedge \Delta_{Y_2} F_2) + (a_2 \wedge 1)(\Delta_{Y_1} F_1 \wedge \psi_{V_2}(F_2)) \\ & \quad + (a_3 \wedge 1)(\psi_{V_1}(F_1) \wedge \psi_{V_2}(F_2)) : \\ & X_1 \wedge X_2 \rightarrow S(LV_1 \oplus LV_2)^+ \wedge (Y_1 \wedge Y_1) \wedge (Y_2 \wedge Y_2) \end{aligned}$$

with

$$\begin{aligned} a_1 & : S(LV_1) \rightarrow S(LV_1 \oplus LV_2) ; x_1 \mapsto (x_1, 0) , \\ a_2 & : S(LV_2) \rightarrow S(LV_1 \oplus LV_2) ; x_2 \mapsto (0, x_2) , \\ a_3 & : S(LV_1) \times S(LV_2) \rightarrow S(LV_1 \oplus LV_2) ; (x_1, x_2) \mapsto (x_1/\sqrt{2}, x_2/\sqrt{2}) . \end{aligned}$$

(vii) (Composition formula) The composite of maps

$$F : V^\infty \wedge X \rightarrow V^\infty \wedge Y , \quad G : V^\infty \wedge Y \rightarrow V^\infty \wedge Z$$

is a map $GF : V^\infty \wedge X \rightarrow V^\infty \wedge Z$ with quadratic construction

$$\psi_V(GF) = (G \wedge G)\psi_V(F) + \psi_V(G)F : X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Z \wedge Z) .$$

Proof. Immediate from the corresponding properties of the geometric Hopf invariant $h_V(F)$ (5.33). □

For any pointed space X write

$$\Sigma^\infty X = \mathbb{R}(\infty)^\infty \wedge X$$

with $\mathbb{R}(\infty) = \varinjlim_k \mathbb{R}^k$. The quadratic construction $\psi_V(F)$ also has a stable version, for $V = \mathbb{R}(\infty)$:

Definition 5.50. The *quadratic construction* on a stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ is the stable homotopy class

$$\psi(F) = \psi_{\mathbb{R}(\infty)}(F) \in \{X; S(\infty)^+ \wedge (Y \wedge Y)\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\}$$

with $S(\infty)$ a contractible space with a free \mathbb{Z}_2 -action. \square

Remark 5.51. (i) The stable homotopy group $\pi_{2i}^S(K(\mathbb{Z}_2, i)) = \mathbb{Z}_2$ is generated by the Hopf construction $S^{2i+1} \rightarrow \Sigma K(\mathbb{Z}_2, i)$ on the map

$$S^i \times S^i \rightarrow K(\mathbb{Z}_2, i) \times K(\mathbb{Z}_2, i) \rightarrow K(\mathbb{Z}_2, i) ,$$

with the mod 2 coefficient quadratic construction defining an isomorphism

$$\begin{aligned} \pi_{2i}^S(K(\mathbb{Z}_2, i)) &\xrightarrow{\cong} \pi_{2i}^S(S(\infty)^+ \wedge_{\mathbb{Z}_2} (K(\mathbb{Z}_2, i) \wedge K(\mathbb{Z}_2, i))) = \mathbb{Z}_2 ; \\ (F : V^\infty \wedge S^{2i} \rightarrow V^\infty \wedge K(\mathbb{Z}_2, i)) &\mapsto \psi(F) . \end{aligned}$$

(ii) The mod 2 coefficient quadratic construction of a stable map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ can be expressed in terms of the functional Steenrod squares: for any $y \in H^i(Y; \mathbb{Z}_2) = [Y, K(\mathbb{Z}_2, i)]$

$$\begin{aligned} y\% \psi_V(F) : H_j(X; \mathbb{Z}_2) &\xrightarrow{\psi_V(F)} Q_j(C(Y; \mathbb{Z}_2)) \xrightarrow{y\%} Q_j(S^i \mathbb{Z}_2) = \mathbb{Z}_2 ; \\ x &\mapsto \langle Sq_{yF}^{j-i+1}(\iota), x \rangle \end{aligned}$$

with $\iota \in H^i(K(\mathbb{Z}_2, i) = \mathbb{Z}_2$ the generator (cf. [50, Proposition 1.6]). If $j = 2i$ and $x \in \text{im}(\pi_{2i}^S(X) \rightarrow H_{2i}(X; \mathbb{Z}_2))$ then

$$y\% \psi_V(x) = yFx \in \pi_{2i}^S(K(\mathbb{Z}_2, i)) = \mathbb{Z}_2 ,$$

with (i) the special case $Y = K(\mathbb{Z}_2, i)$.

(iii) The π -equivariant version of the quadratic construction was used in [50] to express the non-simply-connected surgery obstruction of Wall [68] as the cobordism class of a quadratic Poincaré complex: see 6.9 below for a more detailed discussion. \square

5.6 The ultraquadratic construction $\widehat{\psi}(F)$

The ultraquadratic construction is the quadratic construction $\psi_V(F)$ on a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ in the special case $V = \mathbb{R}$. A knot determines

such a map F , and the ultraquadratic construction on F is a homotopy theoretic version of the Seifert form on the homology of a Seifert surface. See Ranicki [51, pp. 814–842] for an earlier account of the ultraquadratic theory.

Definition 5.52. The *ultraquadratic construction* on a map $F : \Sigma X \rightarrow \Sigma Y$ is the quadratic construction for the special case $V = \mathbb{R}$

$$\widehat{\psi}(F) = \psi_{\mathbb{R}}(F) : X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y) = Y \wedge Y ,$$

identifying $S(LV) = S^0 = \{1, -1\}$ with \mathbb{Z}_2 acting by permutation. \square

Proposition 5.53. The ultraquadratic construction defines a function

$$\begin{aligned} \widehat{\psi} : [\Sigma X, \Sigma Y] &= [\mathbb{R}^\infty \wedge X, \mathbb{R}^\infty \wedge Y] \rightarrow \\ &\{X; S(L\mathbb{R})^+ \wedge Y \wedge Y\}_{\mathbb{Z}_2} = \{X; Y \wedge Y\} ; F \mapsto \psi_{\mathbb{R}}(F) , \end{aligned}$$

with a commutative braid of exact sequences of abelian groups

$$\begin{array}{ccccc} & & 1+T & & \\ & \curvearrowright & & \curvearrowleft & \\ \{X; Y \wedge Y\} & & & & \{X; Y \wedge Y\} \\ & \searrow s_{L\mathbb{R}} & & \nearrow s_{L\mathbb{R}}^* & \\ & & \{X; Y \wedge Y\}_{\mathbb{Z}_2} & & \\ & \nearrow 0_{L\mathbb{R}}^* & & \searrow 0_{L\mathbb{R}} & \\ \{L\mathbb{R}^\infty \wedge X; Y \wedge Y\}_{\mathbb{Z}_2} & & & & \{X; L\mathbb{R}^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2} \\ & \curvearrowright & & \curvearrowleft & \\ & & & & A \end{array}$$

where $A = \{S(L\mathbb{R} \oplus L\mathbb{R})^+ \wedge X; L\mathbb{R}^\infty \wedge Y \wedge Y\}_{\mathbb{Z}_2}$, $s_{L\mathbb{R}}^* : \{X; Y \wedge Y\}_{\mathbb{Z}_2} \rightarrow \{X; Y \wedge Y\}$ is the forgetful map, and

$$s_{L\mathbb{R}} \widehat{\psi}(F) = (F \wedge F) \Delta_X - \Delta_Y F \in \{X; Y \wedge Y\}_{\mathbb{Z}_2} .$$

Proof. This is the special case $V = \mathbb{R}$ of Proposition 5.49 (i)+(ii). \square

Example 5.54. Let $N^n \subset S^{n+1}$ be a codimension 1 framed submanifold with boundary ∂N , with Pontryagin-Thom Umkehr map

$$F : S^{n+1} = \Sigma S^n \rightarrow S^{n+1}/\text{cl.}(S^{n+1} - N \times [0, 1]) = \Sigma(N/\partial N)$$

(interpreting $N/\partial N$ as N^+ if $\partial N = \emptyset$). The Hurewicz image

$$[\widehat{\psi}(F)] \in H_n(\dot{C}(N/\partial N) \otimes_{\mathbb{Z}} \dot{C}(N/\partial N))$$

of the ultraquadratic construction $\widehat{\psi}(F) \in \{S^n; (N/\partial N) \wedge (N/\partial N)\}$ is a \mathbb{Z} -module chain map (or rather a chain homotopy class)

$$[\widehat{\psi}(F)] : \dot{C}(N/\partial N)^{n-*} \rightarrow \dot{C}(N/\partial N) ,$$

defining an ‘ n -dimensional ultraquadratic complex over \mathbb{Z} ’ in the sense of Ranicki [51, p.814]. If $\partial N = S^{n-1}$ then N is a Seifert surface for the $(n-1)$ -knot $\partial N = S^{n-1} \subset S^{n+1}$ and $(\dot{C}(N/\partial N), [\widehat{\psi}(F)])$ is a chain complex generalization of the Seifert form on $\dot{H}_*(N/\partial N) = H_*(N, \partial N)$ – see Example 6.91 below for the connection with linking numbers and the Hopf invariant. \square

In dealing with the special case $V = \mathbb{R}$ we shall use the following terminology :

Definition 5.55. (i) The *positive and negative lines*

$$V^\infty = \{v \in V \mid v \geq 0\}^\infty , \quad V^- = \{v \in V \mid v \leq 0\}^\infty \subset V^\infty$$

are homeomorphic to $[0, 1]$, and such that

$$V^\infty = V^\infty \cup_{0,\infty} V^- .$$

(ii) The *positive and negative circles*

$$S^+(V) = \alpha_V(V^\infty) , \quad S^-(V) = \alpha_V(V^-) \subset \Sigma S(V)^+$$

(with $\alpha_V : V^\infty \rightarrow V^\infty/0^+ = \Sigma S(V)^+$ the canonical projection) are homeomorphic to S^1 , and such that

$$\Sigma S(V)^+ = S^+(V) \vee S^-(V) .$$

(iii) The *positive and negative differences* of maps $p, q : V^\infty \wedge X \rightarrow Y$ such that $p|_{0^+} = q|_{0^+} : 0^+ \wedge X \rightarrow Y$ are the restrictions of the relative difference $\delta(p, q) : \Sigma S(V)^+ \wedge X \rightarrow Y$ (2.18 (ii)) to the positive and negative circles

$$\begin{aligned} \delta^+(p, q) &= \delta(p, q)| : S^+(V) \wedge X \rightarrow Y , \\ \delta^-(p, q) &= \delta(p, q)| : S^-(V) \wedge X \rightarrow Y \end{aligned}$$

with

$$\delta(p, q) = \delta^+(p, q) \vee \delta^-(p, q) : (S^+(V) \wedge X) \vee (S^-(V) \wedge X) \rightarrow Y .$$

□

More directly, working with suspension coordinates the positive and negative differences of maps $p, q : \Sigma X \rightarrow Y$ such that

$$p(1/2, x) = q(1/2, x) \in Y \quad (x \in X)$$

are given (up to homotopy and rescaling) by

$$\begin{aligned} \delta^+(p, q) : \Sigma X \rightarrow Y ; (t, x) &\mapsto \begin{cases} q(1-t, x) & \text{if } 0 \leq t \leq 1/2 \\ pq(t, x) & \text{if } 1/2 \leq t \leq 1 \end{cases} , \\ \delta^-(p, q) : \Sigma X \rightarrow Y ; (t, x) &\mapsto \begin{cases} q(t, x) & \text{if } 0 \leq t \leq 1/2 \\ p(1-t, x) & \text{if } 1/2 \leq t \leq 1 \end{cases} . \end{aligned}$$

Lemma 5.56. *For any map $q : V^\infty \wedge X \rightarrow Y$ such that $q(0^+ \wedge X) = *$ the positive difference $\delta^+(*, q)$ is homotopic to the map*

$$[q]^+ : S^+(V) \wedge X \rightarrow Y ; (v, x) \mapsto q(v, x) .$$

Similarly for the negative difference.

Proof. The homotopies are the restrictions of the homotopy of 2.23 (iv)

$$\begin{aligned} \delta(*, q) &= \delta^+(*, q) \vee \delta^-(*, q) \simeq [q] = [q]^+ \vee [q]^- : \\ \Sigma S(V)^+ \wedge X &= (S^+(V) \wedge X) \vee (S^-(V) \wedge X) \rightarrow Y . \end{aligned}$$

□

Proposition 5.57. *Suppose that $V = \mathbb{R}$, so that (as in 5.55)*

$$\Sigma S(LV)^+ = S^+(LV) \vee S^-(LV) .$$

The \mathbb{Z}_2 -action $T : \Sigma S(LV)^+ \rightarrow \Sigma S(LV)^+$ interchanges the positive and negative circles

$$S^+(LV) = T(S^-(LV)) , \quad S^-(LV) = T(S^+(LV)) .$$

The Hopf invariant (5.30) of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is a \mathbb{Z}_2 -equivariant map

$$\begin{aligned}
h_V(F) &= \delta(p, q) = \delta^+(p, q) \vee T\delta^+(p, q) : \\
\Sigma S(LV)^+ \wedge V^\infty \wedge X &= (S^+(LV) \wedge V^\infty \wedge X) \vee T(S^+(LV) \wedge V^\infty \wedge X) \\
&\rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y)
\end{aligned}$$

which is determined by the positive difference

$$h_V(F) \mid = \delta^+(p, q) : S^+(LV) \wedge V^\infty \wedge X \rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) .$$

The positive difference is homotopic to the Hopf map of Boardman and Steer [4]

$$\delta^+(p, q) \simeq \lambda_2(F) : V^\infty \wedge V^\infty \wedge X \rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) .$$

Proof. In the first instance, we recall the maps μ_2, λ_2 of [4]. Express $V^\infty \wedge V^\infty$ as a union

$$V^\infty \wedge V^\infty = H^+(V) \cup_{\Delta(V)^\infty} H^-(V)$$

of two contractible subspaces

$$\begin{aligned}
H^+(V) &= \{(v, w) \in V^\infty \wedge V^\infty \mid v \geq w\} , \\
H^-(V) &= \{(v, w) \in V^\infty \wedge V^\infty \mid v \leq w\} ,
\end{aligned}$$

with

$$H^+(V) \cap H^-(V) = \Delta(V)^\infty = \{(v, w) \in V^\infty \wedge V^\infty \mid v = w\} .$$

(Up to homeomorphism this is just $S^2 = D^2 \cup_{S^1} D^2$). Given pointed spaces X, Y_1, Y_2 and a map

$$f : V^\infty \wedge X \rightarrow Y_1 \vee Y_2$$

let $\pi_i : Y_1 \vee Y_2 \rightarrow Y_i$ ($i = 1, 2$) be the projections, and define

$$f_i = \pi_i f : V^\infty \wedge X \rightarrow Y_i .$$

Note that for each $(v, x) \in V^\infty \wedge X$ either $f_1(v, x) = *$ or $f_2(v, x) = *$. The μ_2 -function of [4, 5.1]

$$\mu_2 : [V^\infty \wedge X, Y_1 \vee Y_2] \rightarrow [V^\infty \wedge V^\infty \wedge X, Y_1 \wedge Y_2]$$

is defined by

$$\begin{aligned}
\mu_2(f) &= (f_1 \wedge f_2)(1 \wedge \Delta_X) \cup * : \\
V^\infty \wedge V^\infty \wedge X &= (H^+(V) \wedge X) \cup_{\Delta(V)^\infty \wedge X} (H^-(V) \wedge X) \rightarrow Y_1 \wedge Y_2 .
\end{aligned}$$

The projection

$$\pi^+ : V^\infty \wedge V^\infty \rightarrow H^+(V)/\Delta(V)^\infty ; (v, w) \mapsto \begin{cases} (v, w) & \text{if } v \geq w \\ * & \text{if } v \leq w \end{cases}$$

is a homotopy equivalence such that there is defined a commutative diagram

$$\begin{array}{ccc} V^\infty \wedge V^\infty \wedge X & \xrightarrow[\cong]{\pi^+ \wedge 1} & (H^+(V)/\Delta(V)^\infty) \wedge X \\ & \searrow \mu_2(f) & \downarrow [(f_1 \wedge f_2)(1 \wedge \Delta_X)] \\ & & Y_1 \wedge Y_2 . \end{array}$$

Let $\nabla : V^\infty \rightarrow V^\infty \vee V^\infty$ be a sum map, with the components

$$\nabla_i = \pi_i \nabla : V^\infty \xrightarrow{\nabla} V^\infty \vee V^\infty \xrightarrow{\pi_i} V^\infty$$

both homotopic to the identity, and chosen such that

$$(\nabla_1 \wedge \nabla_2) | = * : H^+(V)/\Delta(V)^\infty \rightarrow V^\infty \wedge V^\infty .$$

(If $\dot{\phi} : (0, 1) \rightarrow \mathbb{R}$ is an order-preserving homeomorphism, then the sum map

$$\nabla : V^\infty \rightarrow V^\infty \vee V^\infty ; \dot{\phi}(t) \mapsto \begin{cases} \dot{\phi}(2t)_1 & \text{if } t \leq 1/2 \\ \dot{\phi}(2t-1)_2 & \text{if } t \geq 1/2 \end{cases}$$

has these properties, with $\dot{\phi}(0) = \dot{\phi}(1) = \infty$). The Hopf map of [4] is given by the composite

$$\begin{aligned} \lambda_2 &= \mu_2(\nabla \wedge 1_Y) : \\ [V^\infty \wedge X, V^\infty \wedge Y] &\xrightarrow{\nabla \wedge 1_Y} [V^\infty \wedge X, (V^\infty \wedge Y) \vee (V^\infty \wedge Y)] \\ &\xrightarrow{\mu_2} [V^\infty \wedge V^\infty \wedge X, (V^\infty \wedge Y) \wedge (V^\infty \wedge Y)] . \end{aligned}$$

The Hopf invariant of a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is thus given by

$$\begin{aligned} \lambda_2(F) &= \mu_2((\nabla \wedge 1_Y)F) \\ &= [((\nabla_1 \wedge 1_Y)F \wedge (\nabla_2 \wedge 1_Y)F)(1 \wedge \Delta_X)] \cup * : \\ V^\infty \wedge V^\infty \wedge X &= (H^+(V) \wedge X) \cup_{\Delta(V)^\infty \wedge X} (H^-(V) \wedge X) \\ &\rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) \end{aligned}$$

with a commutative diagram

$$\begin{array}{ccc}
V^\infty \wedge V^\infty \wedge X & \xrightarrow[\simeq]{\pi^+ \wedge 1} & (H^+(V)/\Delta(V)^\infty) \wedge X \\
& \searrow \lambda_2(F) & \downarrow [(\nabla_1 \wedge \nabla_2 \wedge 1)((F \wedge F)(1 \wedge \Delta_X))] \\
& & (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) .
\end{array}$$

Now return to the difference construction (5.30) of the Hopf invariant $h_V(F) = \delta(p, q)$, with

$$\begin{aligned}
p &= (\kappa_V \wedge \Delta_Y)(1 \wedge F) \quad , \quad q = (F \wedge F)(\kappa_V \wedge \Delta_X) : \\
LV^\infty \wedge V^\infty \wedge X &\rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) .
\end{aligned}$$

The composites

$$\begin{aligned}
LV^\infty \wedge V^\infty \wedge X &\longrightarrow LV^\infty \wedge V^\infty \wedge X \xrightarrow{p} (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) \\
&\xrightarrow{\nabla_1 \wedge \nabla_2 \wedge 1} (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) , \\
0^+ \wedge V^\infty \wedge X &\longrightarrow LV^\infty \wedge V^\infty \wedge X \xrightarrow{q} (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) \\
&\xrightarrow{\nabla_1 \wedge \nabla_2 \wedge 1} (V^\infty \wedge Y) \wedge (V^\infty \wedge Y)
\end{aligned}$$

are both $*$. There exist homotopies

$$\nabla_1 \simeq \nabla_2 \simeq 1 : V^\infty \rightarrow V^\infty$$

and hence also a homotopy

$$\nabla_1 \wedge \nabla_2 \simeq 1 : V^\infty \wedge V^\infty \rightarrow V^\infty \wedge V^\infty .$$

Thus there exists a homotopy

$$\delta^+(p, q) \simeq (\nabla_1 \wedge \nabla_2 \wedge 1)\delta^+(p, q) = \delta^+(*, (\nabla_1 \wedge \nabla_2 \wedge 1)q) .$$

Furthermore, 5.56 gives a homotopy

$$\delta^+(*, (\nabla_1 \wedge \nabla_2 \wedge 1)q) \simeq [(\nabla_1 \wedge \nabla_2 \wedge 1)q]^+ ,$$

so that there exists a homotopy

$$\delta^+(p, q) \simeq [(\nabla_1 \wedge \nabla_2 \wedge 1)q]^+ : S^+(LV) \wedge V^\infty \wedge X \rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) .$$

The homeomorphism

$$\kappa_V^+ : S^+(LV) \wedge V^\infty \rightarrow H^+(V)/\Delta(V)^\infty ; (v, w) \mapsto (v + w, -v + w)$$

is such that there is defined a commutative diagram

$$\begin{array}{ccc}
 S^+(LV) \wedge V^\infty \wedge X & & \\
 \downarrow & \searrow \kappa_V^\infty \wedge 1 & \\
 [(\nabla_1 \wedge \nabla_2 \wedge 1)q]^+ & & (H^+(V)/\Delta(V)^\infty) \wedge X \\
 & \swarrow & \\
 (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) & & [(\nabla_1 \wedge \nabla_2 \wedge 1)((F \wedge F)(1 \wedge \Delta_X))]
 \end{array}$$

Putting this together, there are obtained a homotopy equivalence

$$\iota = (\kappa_V^\infty)^{-1} \pi^+ : V^\infty \wedge V^\infty \rightarrow S^+(LV) \wedge V^\infty$$

and a homotopy

$$\lambda_2(F) \simeq \delta^+(p, q)(\iota \wedge 1) : V^\infty \wedge V^\infty \wedge X \rightarrow (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) .$$

□

Terminology 5.58 In view of 5.57 we shall regard the Hopf invariant of a map $F : \Sigma X \rightarrow \Sigma Y$ as a non-equivariant map

$$h_{\mathbb{R}}(F) : \Sigma^2 X \rightarrow \Sigma Y \wedge \Sigma Y .$$

□

Definition 5.59. (i) The *James map* ([28]) is

$$\begin{aligned}
 J : X \times X &\rightarrow Q_{\mathbb{R}}(X) = \Omega \Sigma X ; \\
 (x_1, x_2) &\mapsto \left(s \mapsto \begin{cases} (2s, x_1) & \text{if } 0 \leq s \leq 1/2 \\ (2s-1, x_2) & \text{if } 1/2 \leq s \leq 1 \end{cases} \right) .
 \end{aligned}$$

(ii) Let $E : Q_{\mathbb{R}}(X) \rightarrow X$ be the stable map defined by evaluation

$$E : \Sigma Q_{\mathbb{R}}(X) \rightarrow \Sigma X ; (s, \omega) \mapsto \omega(s) ,$$

so that

$$EJ = \pi_1 + \pi_2 : X \times X \rightarrow X$$

with $\pi_i : X \times X \rightarrow X ; (x_1, x_2) \mapsto x_i$. Note that the composite of the map

$$i : X \rightarrow X \times X ; x \mapsto (x, *)$$

and J is just the inclusion

$$Ji : X \rightarrow Q_{\mathbb{R}}(X) ; x \mapsto (v \mapsto (v, x)) .$$

□

Proposition 5.60. (i) *The ultraquadratic construction*

$$\widehat{\psi}(E) : Q_{\mathbb{R}}(X) \rightarrow X \wedge X$$

is such that up to stable homotopy

$$\begin{aligned} \widehat{\psi}(E)J &= \widehat{\psi}(EJ) = \text{proj.} : X \times X \rightarrow X \wedge X , \\ \widehat{\psi}(E)Ji &= \{*\} : X \rightarrow X \wedge X . \end{aligned}$$

(ii) *The adjoint of a stable map $F : \Sigma W \rightarrow \Sigma X$*

$$\text{adj}(F) : W \rightarrow Q_{\mathbb{R}}(X) ; x \mapsto (s \mapsto F(s, x))$$

is such that $F = E(\text{adj}(F))$ with

$$\widehat{\psi}(F) = \widehat{\psi}(E)\text{adj}(F) : W \rightarrow (X \wedge X) .$$

$$\begin{array}{ccc} W & \xrightarrow{\text{adj}(F)} & Q_{\mathbb{R}}(X) \\ & \searrow \widehat{\psi}(F) & \downarrow \widehat{\psi}(E) \\ & & X \wedge X . \end{array}$$

Proof. By construction. (This is just the special case $V = \mathbb{R}$ of 6.46 (below).)

□

5.7 The spectral quadratic construction $s\psi_V(F)$

The chain level ‘spectral quadratic construction’ ([50, §9.3], [51, §7.3]) of a ‘semistable’ map $F : X \rightarrow V^\infty \wedge Y$ is induced by the following space level geometric Hopf invariant map :

Definition 5.61. The *spectral Hopf invariant map* of a map $F : X \rightarrow V^\infty \wedge Y$ is the \mathbb{Z}_2 -equivariant map given by the relative difference (1.3)

$$sh_V(F) = \delta((G \wedge G)\delta\dot{\phi}_V(V^\infty \wedge Y)(1 \wedge F), \dot{\phi}_V(\mathcal{C}(F))(1 \wedge GF)) : \\ \Sigma S(LV)^+ \wedge X \rightarrow \mathcal{C}(F) \wedge \mathcal{C}(F)$$

with $G : V^\infty \wedge Y \rightarrow \mathcal{C}(F)$ the inclusion in the mapping cone and

$$1 \wedge GF : CS(LV)^+ \wedge X = S(LV)^+ \wedge CX \rightarrow S(LV)^+ \wedge \mathcal{C}(F)$$

the null-homotopy of $1 \wedge GF : S(LV)^+ \wedge X \rightarrow S(LV)^+ \wedge \mathcal{C}(F)$ determined by the inclusion $GF : CX \rightarrow \mathcal{C}(F) = (V^\infty \wedge Y) \cup_F CX$. \square

The spectral Hopf invariant has the same properties as the Hopf invariant obtained in 5.33. Here are two particularly important special cases:

Proposition 5.62. (i) *The \mathbb{Z}_2 -equivariant homotopy class of $sh_V(F)$ depends only on the homotopy class of F .*
(ii) *The spectral Hopf invariant of the composite $EF : X \rightarrow V^\infty \wedge Z$ of maps $F : X \rightarrow V^\infty \wedge Y$, $E : V^\infty \wedge Y \rightarrow V^\infty \wedge Z$ is given up to \mathbb{Z}_2 -equivariant homotopy by*

$$sh_V(EF) = (k \wedge \ell)sh_V(F) + (k \wedge \ell)h_V(E)(1 \wedge F) : \\ \Sigma S(LV)^+ \wedge X \rightarrow \mathcal{C}(EF) \wedge \mathcal{C}(EF)$$

with

$$k = E \cup 1 : \mathcal{C}(F) = V^\infty \wedge Y \cup_F CX \rightarrow \mathcal{C}(EF) = V^\infty \wedge Z \cup_{GF} CX$$

and $\ell : V^\infty \wedge Z \rightarrow \mathcal{C}(GF)$ the inclusion.

Proof. (i) As for the homotopy invariance 5.33 (ii) of the Hopf invariant h_V .
(ii) As for the composition formula 5.33 (v) for the Hopf invariant h_V . \square

The spectral Hopf invariant is closely related to the symmetric construction on the mapping cone:

Proposition 5.63. *Given a map $F : X \rightarrow V^\infty \wedge Y$ let*

$$Z = \mathcal{C}(F) = V^\infty \wedge Y \cup_F CX$$

be the mapping cone, so that there is defined a cofibration sequence

$$X \xrightarrow{F} V^\infty \wedge Y \xrightarrow{G} Z \xrightarrow{H} \Sigma X$$

with G the inclusion and H the projection.

(i) The symmetric construction $\dot{\phi}_V(Z)$ is determined by the spectral Hopf invariant $sh_V(F)$, with a \mathbb{Z}_2 -equivariant homotopy commutative diagram

$$\begin{array}{ccc} S(LV)^+ \wedge Z & \xrightarrow{1 \wedge H} & S(LV)^+ \wedge \Sigma X = \Sigma S(LV)^+ \wedge X \\ & \searrow \dot{\phi}_V(Z) & \downarrow sh_V(F) \\ & & Z \wedge Z . \end{array}$$

(ii) If $X = V^\infty \wedge X_0$ the spectral Hopf invariant $sh_V(F)$ is determined by the Hopf invariant $h_V(F)$, with a commutative diagram

$$\begin{array}{ccc} \Sigma S(LV)^+ \wedge X & \xrightarrow{h_V(F)} & (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) \\ & \searrow sh_V(F) & \downarrow G \wedge G \\ & & Z \wedge Z . \end{array}$$

(iii) Suppose given a space W and a homotopy equivalence $Z \simeq V^\infty \wedge W$, so that the cofibration sequence can be written as

$$X \xrightarrow{F} V^\infty \wedge Y \xrightarrow{G} V^\infty \wedge W \xrightarrow{H} \Sigma X .$$

The spectral Hopf invariant $sh_V(F)$ is determined by the Hopf invariant $h_V(G)$, with a stable \mathbb{Z}_2 -equivariant homotopy commutative diagram

$$\begin{array}{ccc} \Sigma S(LV)^+ \wedge X & \xrightarrow{1 \wedge F} & \Sigma S(LV)^+ \wedge V^\infty \wedge Y \\ & \searrow sh_V(F) & \downarrow -h_V(G) \\ & & Z \wedge Z . \end{array}$$

Proof. (i) As in 5.38 there is defined a natural transformation of cofibration sequences

$$\begin{array}{ccccccc}
S(LV)^+ \wedge X & \xrightarrow{1 \wedge F} & S(LV)^+ \wedge V^\infty \wedge Y & \xrightarrow{1 \wedge G} & S(LV)^+ \wedge Z & \xrightarrow{1 \wedge H} & S(LV)^+ \wedge \Sigma X \\
\dot{\phi}_V(X) \downarrow & & \dot{\phi}_V(V^\infty \wedge Y) \downarrow & & \dot{\phi}_V(F) \downarrow & & \Sigma \dot{\phi}_V(X) \downarrow \\
X \wedge X & \xrightarrow{F \wedge F} & (V^\infty \wedge Y) \wedge (V^\infty \wedge Y) & \longrightarrow & \mathcal{C}(F \wedge F) & \longrightarrow & \Sigma(X \wedge X)
\end{array}$$

and 5.26 gives a \mathbb{Z}_2 -equivariant null-homotopy $\delta \dot{\phi}_V(V^\infty \wedge Y) : \dot{\phi}_V(V^\infty \wedge Y) \simeq *$. Now apply 1.11 (iii) to get a \mathbb{Z}_2 -equivariant homotopy $\dot{\phi}_V(F) \simeq \delta(i, j)$ (for the appropriate i, j) and compose with the projection $\mathcal{C}(F \wedge F) \rightarrow Z \wedge Z$ to get a \mathbb{Z}_2 -equivariant homotopy $\dot{\phi}_V(Z) \simeq sh_V(F)(1 \wedge H)$.

(ii) By construction.

(iii) The application of 5.62 to $GF \simeq *$ shows that up to stable \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned}
sh_V(GF) &= sh_V(F) + h_V(G)(1 \wedge F) \\
&= sh_V(*) = 0 : \Sigma S(LV)^+ \wedge X \rightarrow Z \wedge Z .
\end{aligned}$$

□

Remark 5.64. The expression in 5.63 (i) of the symmetric construction $\dot{\phi}_V(\mathcal{C}(F))$ on the mapping cone $\mathcal{C}(F)$ of a map $F : X \rightarrow V^\infty \wedge Y$ in terms of the spectral Hopf invariant $sh_V(F)$ is a generalization of the relationship between the functional Steenrod squares of a stable map F and the Steenrod squares of $\mathcal{C}(F)$. □

Definition 5.65. The *spectral quadratic construction* on a map $F : X \rightarrow V^\infty \wedge Y$ is the stable \mathbb{Z}_2 -equivariant map

$$s\psi_V(F) : LV^\infty \wedge X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (\mathcal{C}(F) \wedge \mathcal{C}(F))$$

given by the image of the \mathbb{Z}_2 -equivariant spectral Hopf invariant (5.42)

$$sh_V(F) : \Sigma S(LV)^+ \wedge X \rightarrow \mathcal{C}(F) \wedge \mathcal{C}(F)$$

under the composite of the \mathbb{Z}_2 -equivariant S -duality isomorphism of Proposition 4.85

$$\{\Sigma S(LV)^+ \wedge X; \mathcal{C}(F) \wedge \mathcal{C}(F)\}_{\mathbb{Z}_2} \cong \{LV^\infty \wedge X; S(LV)^+ \wedge \mathcal{C}(F) \wedge \mathcal{C}(F)\}_{\mathbb{Z}_2} .$$

□

Example 5.66. The spectral quadratic construction on a map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is the composite

$$\begin{aligned}
s\psi_V(F) : V^\infty \wedge LV^\infty \wedge X &\xrightarrow{1 \wedge \psi_V(F)} S(LV)^+ \wedge LV^\infty \wedge V^\infty \wedge Y \wedge Y \\
&\cong S(LV)^+ \wedge ((V^\infty \wedge Y) \wedge (V^\infty \wedge Y)) \\
&\xrightarrow{1 \wedge G \wedge G} S(LV)^+ \wedge (\mathcal{C}(F) \wedge \mathcal{C}(F))
\end{aligned}$$

with $G : V^\infty \wedge Y \rightarrow \mathcal{C}(F)$ the inclusion.

□

Remark 5.67. (i) The spectral quadratic construction $s\psi_V(F)$ is a space level version of the chain level spectral quadratic construction of Ranicki [50, §9.3], [51, p.589]. A triangulated n -dimensional normal space

$$(X, \nu_X : X \rightarrow BG(k), \rho_X : S^{n+k} \rightarrow T(\nu_X))$$

in the sense of Quinn [47] determines an n -dimensional normal complex $(C(X), \phi, \gamma, \chi)$ in the X -local additive category with chain duality $\mathbb{A}(\mathbb{Z}, X)$ ([50, Example 9.12]). The boundary

$$\partial(C(X), \phi, \gamma, \chi) = (C, \psi)$$

is an $(n-1)$ -dimensional quadratic Poincaré complex in $\mathbb{A}(\mathbb{Z}, X)$ with $C = \mathcal{C}([X] \cap -)_{*+1}$ the algebraic mapping cone with a dimension shift of the cap product $[X] \cap - : C(X)^{n-*} \rightarrow C(X)$ with the fundamental class $[X] = [\rho_X(S^{n+k})] \in H_n(X)$. The quadratic structure ψ is determined by the spectral quadratic construction on the semistable map

$$F : T(\nu_X)^* \rightarrow V^\infty \wedge X^+ = \Sigma^k X^+ \quad (V = \mathbb{R}^k)$$

from an S -dual $T(\nu_X)^*$ of the Thom space $T(\nu_X)$ inducing

$$F = [X] \cap - : \dot{C}(T(\nu_X)^*)_{*+k} = C(X)^{n-*} \rightarrow \dot{C}(V^\infty \wedge X^+)_{*+k} = C(X)$$

([51, Proposition 7.3.4]). The cobordism class

$$\hat{\sigma}^*(X) = (C(X), \phi, \gamma, \chi) \in H_n(X; \hat{\mathbb{L}}(\mathbb{Z}))$$

is the normal (= hyperquadratic) signature of [48], [51, Proposition 7.4.1], [52, §2].

(ii) An n -dimensional geometric Poincaré complex X has a Spivak normal structure (ν_X, ρ_X) . The total surgery obstruction (Ranicki [48], [51, §7.4], [52, §16], Kühl, Macko, Mole [39]) of a triangulated X is the cobordism class

$$s(X) = \partial \hat{\sigma}^*(X) = (C, \psi) \in \mathbb{S}_n(X)$$

in the cobordism group of $(n-1)$ -dimensional quadratic Poincaré complexes in $\mathbb{A}(\mathbb{Z}, X)$ with $\mathbb{Z}[\pi_1(X)]$ -coefficient Poincaré duality, with the property that

$s(X) = 0$ if (and for $n > 5$ only if) X is homotopy equivalent to an n -dimensional topological manifold.

□

5.8 Stably trivialized vector bundles

We now apply the geometric Hopf invariant to the classification of pairs $(\delta\xi, \xi)$ consisting of a U -vector bundle $\xi : X \rightarrow BO(U)$ and a V -stable trivialization

$$\delta\xi : \xi \oplus \epsilon_V \cong \epsilon_{U \oplus V}$$

with U, V finite-dimensional inner product spaces. We relate the classifying map

$$c = (\delta\xi, \xi) : X \rightarrow O(V, U \oplus V)$$

to the geometric Hopf invariant $h_V(F)$ of the V -stable map

$$\begin{aligned} F : V^\infty \wedge T(\xi) &= T(\xi \oplus \epsilon_V) \xrightarrow[\cong]{T(\delta\xi)} T(\epsilon_{U \oplus V}) = (U \oplus V)^\infty \wedge X^+ \\ &\xrightarrow{\text{proj.}} V^\infty \wedge U^\infty, \end{aligned}$$

with stable homotopy class

$$\begin{aligned} h_V(F) &= h'_V(F) = \psi_V(F) \in \{\Sigma S(LV)^+ \wedge T(\xi); LV^\infty \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2} \\ &= \{T(\xi); S(LV)^+ \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2} \\ &= \{T(\xi); S(LV)^+ \wedge_{\mathbb{Z}_2} (U^\infty \wedge U^\infty)\} \\ &= \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} LU^\infty\}. \end{aligned}$$

The stable homotopy relationship between the Stiefel space $O(V, U \oplus V)$ and the stunted projective space

$$S(LV)^\infty \wedge_{\mathbb{Z}_2} LU^\infty = P(U \oplus V)/P(U)$$

studied by James [30] and Crabb [12] is used in Proposition 5.75 (v) to identify the quadratic construction $\psi_V(F)$ with the ‘local obstruction’ $\theta(c)$ of [12] (cf. Definition 5.72 below)

$$\psi_V(F) = \theta(c) \in \{X^+; S(LV)^\infty \wedge_{\mathbb{Z}_2} LU^\infty\}.$$

In particular, $c^{univ} = 1 : O(V, U \oplus V) \rightarrow O(V, U \oplus V)$ classifies the universal U -bundle with a V -stable trivialization

$$(\eta(U) : O(V, U \oplus V) \rightarrow BO(U) , \delta\eta(U) : \eta(U) \oplus \epsilon_V \cong \epsilon_{U \oplus V})$$

with a corresponding V -stable map

$$F^{univ} : V^\infty \wedge T(\eta(U)) \cong (V \oplus U)^\infty \wedge O(V, U \oplus V)^+ \rightarrow V^\infty \wedge U^\infty .$$

The quadratic construction/local obstruction

$$\begin{aligned} \psi_V(F^{univ}) &= \theta(c^{univ}) \\ &\in \{T(\eta(U)); S(LV)^+ \wedge_{\mathbb{Z}_2} (U^\infty \wedge U^\infty)\} = \{O(V, U \oplus V)^+; S(LV)^\infty \wedge_{\mathbb{Z}_2} LU^\infty\} \end{aligned}$$

defines a stable map $\theta : O(V, U \oplus V)^+ \rightarrow S(LV)^\infty \wedge_{\mathbb{Z}_2} LU^\infty$ which is $2 \dim(U)$ -connected. It follows that the function

$$[X, O(V, U \oplus V)] \rightarrow \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} LU^\infty\} ; c \mapsto \psi_V(F) = \theta(c)$$

is a bijection if X is a CW complex with $\dim(X) < 2 \dim(U)$.

Definition 5.68. (i) Given an inner product space V and a 1-dimensional subspace $L = \mathbb{R}x \subset V$ ($x \in S(V)$) let

$$L^\perp = \{v \in V \mid \langle v, x \rangle = 0\} ,$$

and define the linear isometry reflecting V in $L^\perp \subset V$

$$\begin{aligned} R_L &= -1_L \oplus 1_{L^\perp} : V = L \oplus L^\perp \rightarrow V = L \oplus L^\perp ; \\ v &= (\lambda, \mu) \mapsto v - 2\langle v, x \rangle x = (-\lambda, \mu) . \end{aligned}$$

(ii) The *reflection* map

$$\begin{aligned} R &: P(U \oplus V)/P(U) \rightarrow O(V, U \oplus V) = O(U \oplus V)/O(U) ; \\ L &= \mathbb{R}x \mapsto (R_L| : v \mapsto (0, v) - 2\langle (0, v), x \rangle x) \quad (x \in S(U \oplus V), v \in V) \end{aligned}$$

sends $L \subset U \oplus V$ to $R_L| : V \rightarrow U \oplus V$.

□

Example 5.69. (i) For $U = \{0\}$, $V = \mathbb{R}$ the reflection map is a homeomorphism

$$R : P(\mathbb{R})^+ = S^0 \rightarrow O(\mathbb{R}) ; \pm 1 \mapsto \pm 1 .$$

(ii) For $U = \{0\}$, $V = \mathbb{R}^2$ the reflection map is the embedding

$$R : P(\mathbb{R}^2)^+ = (S^1)^+ \hookrightarrow O(\mathbb{R}^2) ; \begin{cases} [\cos t, \sin t] \mapsto \begin{pmatrix} -\cos 2t & -\sin 2t \\ -\sin 2t & \cos 2t \end{pmatrix} \\ * \mapsto \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \end{cases}$$

with image consisting of all the reflections $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ in lines through the origin and the identity map. \square

Example 5.70. (i) The reflection map

$$R : P(U \oplus \mathbb{R})/P(U) = U^\infty \rightarrow O(\mathbb{R}, U \oplus \mathbb{R}) = S(U \oplus \mathbb{R}) ; \\ u \mapsto \left(\frac{-2u}{|u|^2 + 1}, \frac{|u|^2 - 1}{|u|^2 + 1} \right), \quad \infty \mapsto (0, 1)$$

is a homeomorphism, with inverse

$$S(U \oplus \mathbb{R}) \rightarrow U^\infty ; (x, y) \mapsto \frac{x}{y - 1} .$$

(ii) The composite

$$\tilde{R} : S(V) \longrightarrow P(V) \subset P(V)^+ = P(0 \oplus V)/P(0) \xrightarrow{R} O(V) ; \\ v \mapsto (w \mapsto w - 2\langle v, w \rangle v)$$

is the clutching function for the tangent V -bundle of $S(V \oplus \mathbb{R})$

$$\tau_{S(V \oplus \mathbb{R})} : S(V \oplus \mathbb{R}) = D(V) \cup_{S(V)} D(V) \rightarrow BO(V) ,$$

with

$$E(\tau_{S(V \oplus \mathbb{R})}) = D(V) \times V \cup_{(x, v) \sim (x, \tilde{R}(x)(v))} D(V) \times V \quad (x \in S(V), v \in V) .$$

The classifying map for $\tau_{S(V \oplus \mathbb{R})}$ fits into a fibration

$$S(V \oplus \mathbb{R}) \xrightarrow{\tau_{S(V \oplus \mathbb{R})}} BO(V) \longrightarrow BO(V \oplus \mathbb{R})$$

corresponding to the stable isomorphism

$$\tau_{S(V \oplus \mathbb{R})} \oplus \epsilon_{\mathbb{R}} \cong \epsilon_{V \oplus \mathbb{R}}$$

determined by the framed embedding $S(V \oplus \mathbb{R}) \subset V \oplus \mathbb{R}$. \square

The reflection map $R : P(U \oplus V)/P(U) \rightarrow O(V, U \oplus V)$ is injective, which we use to regard $P(U \oplus V)/P(U)$ as a subspace of $O(V, U \oplus V)$.

Proposition 5.71. (James [30, Prop. 1.3 and Thm. 3.4])
The pair $(O(V, U \oplus V), P(U \oplus V)/P(U))$ is $2 \dim(U)$ -connected, so that

$$R_* : \pi_i(P(U \oplus V)/P(U)) \rightarrow \pi_i(O(V, U \oplus V))$$

is an isomorphism for $i < 2 \dim(U)$ and a surjection for $i = 2 \dim(U)$.

Proof. (Sketch) By induction on $\dim(V)$, using the Blakers-Massey theorem to compare the morphism induced by R from the homotopy exact sequence (in a certain range of dimensions) of the cofibration sequence

$$P(U \oplus V)/P(U) \rightarrow P(U \oplus V \oplus \mathbb{R})/P(U) \rightarrow (U \oplus V)^\infty$$

to the homotopy exact sequence of the fibration sequence

$$O(V, U \oplus V) \rightarrow O(V \oplus \mathbb{R}, U \oplus V \oplus \mathbb{R}) \rightarrow (U \oplus V)^\infty .$$

□

For any finite-dimensional inner product space V the \mathbb{Z}_2 -action on LV restricts to the antipodal involution on the unit sphere $S(LV)$, with

$$S(LV)/\mathbb{Z}_2 = P(V) , \quad LV^\infty/\mathbb{Z}_2 = sP(V) .$$

The cofibration sequence of Proposition 2.17 (iii)

$$\begin{aligned} S(LU) &\rightarrow S(LU \oplus LV) \rightarrow LU^\infty \wedge S(LV)^+ \\ &\rightarrow LU^\infty \rightarrow (LU \oplus LV)^\infty \rightarrow LU^\infty \wedge \Sigma S(LV)^+ \rightarrow \dots \end{aligned}$$

is \mathbb{Z}_2 -equivariant, so that passing to the \mathbb{Z}_2 -quotients there is defined a cofibration sequence

$$\begin{aligned} P(U) &\rightarrow P(U \oplus V) \rightarrow P(U \oplus V)/P(U) = LU^\infty \wedge_{\mathbb{Z}_2} S(LV)^+ \\ &\rightarrow sP(U) \rightarrow sP(U \oplus V) \rightarrow LU^\infty \wedge_{\mathbb{Z}_2} \Sigma S(LV)^+ \rightarrow \dots \end{aligned}$$

If $U = \{0\}$ interpret $P(U \oplus V)/P(U)$ as $P(V)^+$.

Let $j = \dim(U)$, $k = \dim(V)$, so that

$$O(V, U \oplus V) = V_{j+k, k} .$$

The reduced homology groups of the stunted projective space

$$P(U \oplus V)/P(U) = LU^\infty \wedge_{\mathbb{Z}_2} S(LV)^+$$

are the Q -groups

$$\begin{aligned} \dot{H}_i(P(U \oplus V)/P(U)) &= Q_{i+j}^{[0,k-1]}(S^j\mathbb{Z}) \\ &= \begin{cases} 0 & \text{if } i < j \\ \mathbb{Z} & \text{if } i = j \text{ is even, or if } i = j, k = 1 \\ \mathbb{Z}_2 & \text{if } i = j \text{ is odd and } k \geq 2. \end{cases} \end{aligned}$$

The induced morphisms in homology

$$\begin{aligned} \theta_* : H_i(O(V, U \oplus V)) &= H_i(V_{j+k,k}) \\ &\rightarrow H_i(P(U \oplus V)/P(U)) = Q_{i+j}^{[0,k-1]}(S^j\mathbb{Z}) \quad (i > 0) \end{aligned}$$

are split surjections, which are isomorphisms for $i < 2j$ by 5.71 (cf. 3.25 for $i \leq j$).

The Stiefel space

$$O(V, U \oplus V) = O(U \oplus V)/O(U) = V_{j+k,k}$$

fits into a fibration

$$O(V, U \oplus V) \rightarrow BO(U) \rightarrow BO(U \oplus V) .$$

The canonical U -bundle $\eta(U)$ over $O(V, U \oplus V)$ is such that

$$\begin{aligned} E(\eta(U)) &= \{(f, x) \mid f \in O(V, U \oplus V), x \in f(V)^\perp \subset U \oplus V\} \\ &= O(U \oplus V) \times_{O(U)} U \end{aligned}$$

with the canonical $U \oplus V$ -bundle isomorphism

$$\delta\eta(U) : \eta(U) \oplus \epsilon_V \cong \epsilon_{U \oplus V}$$

defined by

$$\begin{aligned} \delta\eta(U) : E(\eta(U) \oplus \epsilon_V) &= E(\eta(U)) \times V \\ &\rightarrow E(\epsilon_{U \oplus V}) = O(V, U \oplus V) \times U \oplus V ; (f, x, v) \mapsto (f, x + f(v)) . \end{aligned}$$

A map $c : X \rightarrow O(V, U \oplus V)$ classifies a U -bundle $\xi : X \rightarrow BO(U)$ with a stable isomorphism $\delta\xi : \xi \oplus \epsilon_V \cong \epsilon_{U \oplus V}$, where

$$E(\xi) = \{(x \in X, y \in c(x)^\perp \subset U \oplus V)\} \subset E(\epsilon_{U \oplus V}) = X \times U \oplus V .$$

The adjoint \mathbb{Z}_2 -equivariant map

$$F_c : LV^\infty \wedge X^+ \rightarrow (LU \oplus LV)^\infty ; (v, x) \mapsto c(x)(v)$$

represents the \mathbb{Z}_2 -equivariant Euler class of $L\xi : X \rightarrow BO^{\mathbb{Z}_2}(U)$

$$\gamma^{\mathbb{Z}_2}(L\xi) = F_c \in \omega_{\mathbb{Z}_2}^0(X; -L\xi) = \{X; LU^\infty\}_{\mathbb{Z}_2} .$$

The standard pair $(1, \epsilon_U)$ is classified by the constant map

$$0 : X \rightarrow O(V, U \oplus V) ; x \mapsto (v \mapsto (0, v))$$

with adjoint \mathbb{Z}_2 -equivariant map

$$F_0 : LV^\infty \wedge X^+ \rightarrow (LU \oplus LV)^\infty ; (v, x) \mapsto (0, v) .$$

The adjoint \mathbb{Z}_2 -equivariant maps F_c, F_0 are such that

$$F_c(0, x) = F_0(0, x) = (0, 0) \in (LU \oplus LV)^\infty .$$

Definition 5.72. (Crabb [12, 2.6])

(i) The *local obstruction* of $c : X \rightarrow O(V, U \oplus V)$ is the relative difference \mathbb{Z}_2 -equivariant map

$$\begin{aligned} \theta(c) &= \delta(F_c, F_0) : \Sigma S(LV)^+ \wedge X^+ \rightarrow (LU \oplus LV)^\infty ; \\ ([t, v], x) &\mapsto \begin{cases} (0, [1 - 2t, v]) & \text{if } 0 \leq t \leq 1/2 \\ c(x)[2t - 1, v] & \text{if } 1/2 \leq t \leq 1 . \end{cases} \end{aligned}$$

The stable \mathbb{Z}_2 -equivariant homotopy class

$$\theta(c) \in \{\Sigma S(LV)^+ \wedge X^+; (LU \oplus LV)^\infty\}_{\mathbb{Z}_2}$$

has image

$$[\theta(c)] = \gamma^{\mathbb{Z}_2}(L\xi) - \gamma^{\mathbb{Z}_2}(L\epsilon_U) \in \{X^+; LU^\infty\}_{\mathbb{Z}_2} .$$

(ii) Let

$$\theta'(c) \in \{X^+; S(LV)^+ \wedge LU^\infty\}_{\mathbb{Z}_2} = \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} LU^\infty\}$$

be the \mathbb{Z}_2 -equivariant S -dual of $\theta(c) \in \{\Sigma S(LV)^+ \wedge X^+; (LU \oplus LV)^\infty\}_{\mathbb{Z}_2}$. \square

Example 5.73. For any inner product space U the homeomorphism

$$c : S(U \oplus \mathbb{R}) \rightarrow O(\mathbb{R}, U \oplus \mathbb{R}) ; (u, v) \mapsto (t \mapsto (tu, tv)) \\ (t \in V = \mathbb{R}, (u, v) \in S(U \oplus \mathbb{R}))$$

classifies the tangent U -bundle

$$\xi = \tau_{S(U \oplus \mathbb{R})} : S(U \oplus \mathbb{R}) \rightarrow BO(U)$$

with the $U \oplus \mathbb{R}$ -bundle isomorphism

$$\delta\xi : \epsilon_{\mathbb{R}} \oplus \xi \cong \epsilon_{U \oplus \mathbb{R}}$$

determined by the embedding $S(U \oplus \mathbb{R}) \subset U \oplus \mathbb{R}$. The local obstruction of c is

$$\theta(c) = 1 \in \{S(U \oplus \mathbb{R})^+; S(L\mathbb{R})^+ \wedge LU^\infty\}_{\mathbb{Z}_2} = \{S(U \oplus \mathbb{R})^+; U^\infty\} = \mathbb{Z}.$$

The map

$$F_c : \mathbb{R}^\infty \wedge S(U \oplus \mathbb{R})^+ \rightarrow \mathbb{R}^\infty \wedge U^\infty \wedge S(U \oplus \mathbb{R})^+ ; \\ (t, u, v) \mapsto (c(u, v)(t), (u, v)) = (tv, tu, u, v) \ (t \in \mathbb{R}, (u, v) \in S(U \oplus \mathbb{R})) .$$

has stable homotopy class

$$F_c = 1 \in \{S(U \oplus \mathbb{R})^+; U^\infty \wedge S(U \oplus \mathbb{R})^+\} = \{S(U \oplus \mathbb{R})^+; U^\infty\} = \mathbb{Z} ,$$

and \mathbb{R} -coefficient quadratic construction

$$\psi_{\mathbb{R}}(F_c) = \theta(c)F_c = 1 \\ \in \{S(U \oplus \mathbb{R})^+; S(L\mathbb{R})^+ \wedge U^\infty \wedge LU^\infty \wedge S(U \oplus \mathbb{R})^+ \wedge S(U \oplus \mathbb{R})^+\}_{\mathbb{Z}_2} \\ = \{S(U \oplus \mathbb{R})^+; U^\infty \wedge U^\infty \wedge S(U \oplus \mathbb{R})^+ \wedge S(U \oplus \mathbb{R})^+\} = \mathbb{Z} .$$

□

Example 5.74. Suppose $U = \{0\}$, and $c \in O(V)$. The \mathbb{Z}_2 -equivariant homotopy class of the local obstruction $\theta(c) : \Sigma S(LV)^+ \rightarrow LV^\infty$ is given by Proposition 4.21 to be

$$\theta(c) = \text{bi-degree}(\delta(F_c, F_0)) \\ = \left(\frac{\text{degree}(F_c) - \text{degree}(F_0)}{2}, \text{degree}(G_c) - \text{degree}(G_0) \right) \\ = \left(\frac{\text{degree}(c) - 1}{2}, 1 - 1 \right) \\ = \begin{cases} (0, 0) & \text{if } \det(c) = 1 \\ (-1, 0) & \text{if } \det(c) = -1 \end{cases} \in [\Sigma S(LV)^+, LV^\infty]_{\mathbb{Z}_2} = \mathbb{Z} \oplus \mathbb{Z}$$

with

$$G_c = G_0 = 1 : (LV^\infty)^{\mathbb{Z}_2} = \{0\}^+ \rightarrow \{0\}^+$$

the fixed point maps.

□

Proposition 5.75. *Let $\xi : X \rightarrow BO(U)$ be a U -vector bundle with a V -stable trivialization $\delta\xi : \xi \oplus \epsilon_V \cong \epsilon_{U \oplus V}$.*

(i) *The map*

$$T(\delta\xi) : T(\xi \oplus \epsilon_V) = V^\infty \wedge T(\xi) \rightarrow T(\epsilon_{U \oplus V}) = V^\infty \wedge U^\infty \wedge X^+$$

is a homeomorphism inducing an isomorphism

$$T(\delta\xi)^* : \{X; S^0\} \xrightarrow{\cong} \{T(\xi); U^\infty\}.$$

The stable cohomotopy Thom class (3.47) of ξ

$$u(\xi) = T(\delta\xi)^*(1) \in \{T(\xi); U^\infty\} = \omega^0(X; \xi - \epsilon_U)$$

is represented by the composite

$$\begin{aligned} F &= (1 \wedge p)T(\delta\xi) : T(\epsilon_V \oplus \xi) = V^\infty \wedge T(\xi) \xrightarrow{T(\delta\xi)} \\ &T(\epsilon_{U \oplus V}) = V^\infty \wedge U^\infty \wedge X^+ \xrightarrow{1 \wedge p} V^\infty \wedge U^\infty \end{aligned}$$

with $p : X^+ \rightarrow S^0$ the projection sending X to the non-base-point of S^0 . The adjoint map

$$F_c : V^\infty \wedge X^+ \rightarrow V^\infty \wedge U^\infty ; (v, x) \mapsto c(x)(v)$$

is the composite

$$F_c : V^\infty \wedge X^+ \xrightarrow{1 \wedge z_\xi} V^\infty \wedge T(\xi) \xrightarrow{F} V^\infty \wedge U^\infty$$

with $z_\xi : X^+ \rightarrow T(\xi)$ the zero section. The images of $u(\xi) \in \{T(\xi); U^\infty\}$ in the commutative square

$$\begin{array}{ccc} \{T(\xi); U^\infty\} & \longrightarrow & \dot{H}^j(T(\xi)) \\ \downarrow z_\xi^* & & \downarrow z_\xi^* \\ \{X^+; U^\infty\} & \longrightarrow & H^j(X) \end{array}$$

are the Thom class $[u(\xi)] \in \dot{H}^j(T(\xi))$, the stable cohomotopy Euler class (3.47)

$$z_\xi^* u(\xi) = \gamma(\xi) \in \{X^+; U^\infty\}$$

and the Euler class $[\gamma(\xi)] \in H^j(X)$. The adjoint \mathbb{Z}_2 -equivariant maps

$$F_c, F_0 : LV^\infty \wedge X^+ \rightarrow (LV \oplus LU)^\infty$$

with 0 the constant map

$$0 : X \rightarrow O(V, U \oplus V) ; x \mapsto (v \mapsto (0, v))$$

represent the stable \mathbb{Z}_2 -equivariant cohomotopy Euler classes

$$\gamma^{\mathbb{Z}_2}(L\xi) = F_c, \gamma^{\mathbb{Z}_2}(L\epsilon_U) = F_0 \in \{X^+; LU^\infty\}_{\mathbb{Z}_2}.$$

(ii) The geometric Hopf invariants of F and F_c

$$\begin{aligned} h_V(F) &\in \{\Sigma S(LV)^+ \wedge T(\xi); LV^\infty \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2}, \\ h_V(F_c) &\in \{\Sigma S(LV)^+ \wedge X^+; LV^\infty \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2} \end{aligned}$$

are related by

$$h_V(F_c) = h_V(F)(1 \wedge z_\xi).$$

(iii) The \mathbb{Z}_2 -equivariant stable homotopy class of the geometric Hopf invariant of F_c is given by

$$\begin{aligned} h_V(F_c) &= (\theta(c) \wedge F_c)(1 \wedge \Delta_X) \\ &\in \{\Sigma S(LV)^+ \wedge X^+; LV^\infty \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2} \\ &= \{\Sigma S(LV)^+ \wedge V^\infty \wedge X^+; (LU \oplus LV)^\infty \wedge (U \oplus V)^\infty\}_{\mathbb{Z}_2} \end{aligned}$$

and is represented by the \mathbb{Z}_2 -equivariant stable map

$$\begin{aligned} &(\theta(c) \wedge F_c)(1 \wedge \Delta_X) : \\ &\Sigma S(LV)^+ \wedge V^\infty \wedge X^+ \rightarrow (LU \oplus LV)^\infty \wedge (U \oplus V)^\infty ; \\ &([t, u], v, x) \mapsto (\theta(c)([t, u], x), c(x)(v)) . \end{aligned}$$

(iv) The isomorphism of exact sequences

$$\begin{array}{ccccc} \{X^+; S(LV)^+ \wedge LU^\infty\}_{\mathbb{Z}_2} & \longrightarrow & \{X^+; LU^\infty\}_{\mathbb{Z}_2} & \longrightarrow & \{X^+; (LU \oplus LV)^\infty\}_{\mathbb{Z}_2} \\ \cong \downarrow T(\delta\xi)^* \wedge \kappa_U & & \cong \downarrow T(\delta\xi)^* \wedge \kappa_U & & \cong \downarrow T(\delta\xi)^* \wedge \kappa_U \\ \{T(\xi); S(LV)^+ \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2} & \simeq & \{T(\xi); U^\infty \wedge U^\infty\}_{\mathbb{Z}_2} & \simeq & \{T(\xi); LV^\infty \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2} \end{array}$$

sends $\theta'(c) \in \{X; S(LV)^+ \wedge LU^\infty\}_{\mathbb{Z}_2}$ to the stable geometric Hopf invariant $h'_V(F) \in \{T(\xi); S(LV)^+ \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2}$. The stable geometric Hopf invariant of F_c is given by

$$h'_V(F_c) = h'_V(F)z_\xi \in \{X^+; S(LV)^+ \wedge (U^\infty \wedge U^\infty)\}_{\mathbb{Z}_2}.$$

(v) *The isomorphism*

$$T(\delta\xi)^* \wedge \kappa_U : \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} LU^\infty\} \xrightarrow{\cong} \{T(\xi); S(LV)^+ \wedge_{\mathbb{Z}_2} (U^\infty \wedge U^\infty)\}$$

sends the local obstruction

$$\theta(c) \in \{X^+; P(U \oplus V)/P(U)\} = \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} LU^\infty\}$$

to the V -coefficient quadratic construction $\psi_V(F)$, so we can identify

$$\theta(c) = \psi_V(F) \in \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} LU^\infty\} = \{T(\xi); S(LV)^+ \wedge_{\mathbb{Z}_2} (U^\infty \wedge U^\infty)\}.$$

(vi) *The V -coefficient quadratic construction on F_c is the product of $F_c = \gamma(\xi) \in \{X^+; U^\infty\}$ and $\theta(c)$, that is*

$$\psi_V(F_c) = (\theta(c) \times \gamma(\xi))\Delta_X \in \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} (U^\infty \wedge U^\infty)\}$$

inducing

$$\psi_V(F_c) : H_i(X) \rightarrow Q_i^{[0, k-1]}(S^j \mathbb{Z}).$$

The local obstruction stable map

$$\theta : O(V, U \oplus V) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} LU^\infty = P(U \oplus V)/P(U)$$

is $2j$ -connected (5.71), so that for $i < 2j$

$$\theta(c)_* = c_* : H_i(X) \rightarrow H_i(O(V, U \oplus V)) = Q_{i+j}^{[0, k-1]}(S^j \mathbb{Z}).$$

Now F_c induces the \mathbb{Z} -module chain map

$$F_c = [\gamma(\xi)] \cap - : C(X) \rightarrow S^j \mathbb{Z}$$

and $\theta(c)$ induces a \mathbb{Z} -module chain map

$$\theta(c) : C(X) \rightarrow W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} S^j(\mathbb{Z}, -1),$$

so that $\psi_V(F_c)$ induces the \mathbb{Z} -module chain map

$$\begin{aligned} \psi_V(F_c) : C(X) &\xrightarrow{\Delta_X} C(X) \otimes_{\mathbb{Z}} C(X) \\ &\xrightarrow{\theta(c) \otimes \gamma(\xi)} W[0, k-1] \otimes_{\mathbb{Z}[\mathbb{Z}_2]} (S^j \mathbb{Z} \otimes_{\mathbb{Z}} S^j \mathbb{Z}). \end{aligned}$$

(vii) *Let $U = U' \oplus \mathbb{R}$. If $\xi = \xi' \oplus \epsilon_{\mathbb{R}}$ for some U' -bundle $\xi' : X \rightarrow BO(U')$ then*

$$\begin{aligned}
F_c &\simeq * : V^\infty \wedge X^+ \rightarrow (U \oplus V)^\infty \text{ (non-equivariantly) ,} \\
\gamma(\xi) &= 0 \in \{X^+; U^\infty\} , \\
h_V(F_c) &= 0 \in \{\Sigma S(LV)^+ \wedge X^+; U^\infty \wedge (LU \oplus LV)^\infty\}_{\mathbb{Z}_2} , \\
\psi_V(F_c) &= 0 \in \{X^+; S(LV)^+ \wedge_{\mathbb{Z}_2} (U^\infty \wedge U^\infty)\} .
\end{aligned}$$

If X is a j -dimensional CW complex then $\xi = \xi' \oplus \epsilon_{\mathbb{R}}$ if and only if

$$\gamma(\xi) = 0 \in \{X^+; U^\infty\} = H^j(X) .$$

Proof. (i)+(ii) By construction.

(iii) The geometric Hopf invariant of F_c is the relative difference $h_V(F_c) = \delta(p, q)$ of the \mathbb{Z}_2 -equivariant maps

$$p, q : LV^\infty \wedge V^\infty \wedge X \rightarrow (LU \oplus LV)^\infty \wedge (U \oplus V)^\infty$$

defined by

$$\begin{aligned}
p(v_1, v_2, x) &= \left(\frac{c(x)(v_1 + v_2) - c(x)(-v_1 + v_2)}{2}, \frac{c(x)(v_1 + v_2) + c(x)(-v_1 + v_2)}{2} \right) \\
&= (c(x)(v_1), c(x)(v_2)) , \\
q(v_1, v_2, x) &= (v_1, c(x)(v_2))
\end{aligned}$$

with

$$p(0, v, x) = q(0, v, x) = (0, c(x)(v)) \quad (v \in V, x \in X) .$$

By Example 2.24

$$\begin{aligned}
h_V(F_c) &= \delta(p, q) = (\theta(c) \wedge F_c)(1 \wedge \Delta_X) : \\
\Sigma S(LV)^+ \wedge V^\infty \wedge X &\rightarrow LV^\infty \wedge V^\infty ; \\
([t, u], v, x) &\mapsto (\theta(c)([t, u], x), c(x)(v)) .
\end{aligned}$$

(iv) The isomorphism $T(\delta\xi)^* \wedge \kappa_U$ sends $F_c, F_0 \in \{X^+; LU^\infty\}_{\mathbb{Z}_2}$ to

$$\begin{aligned}
(T(\delta\xi)^* \wedge \kappa_U)(F_c) &= (F \wedge F) \Delta_{T(\xi)} = p , \\
(T(\delta\xi)^* \wedge \kappa_U)(F_0) &= \Delta_{U^\infty} F = q \in \{T(\xi); U^\infty \wedge U^\infty\}_{\mathbb{Z}_2} .
\end{aligned}$$

so that the stable relative differences are such that

$$\begin{aligned}
(T(\delta\xi)^* \wedge \kappa_U)(\theta'(c)) &= (T(\delta\xi)^* \wedge \kappa_U)(\delta'(F_c, F_0)) \\
&= \delta'(p, q) = h'_V(F) \in \{T(\xi); S(LV)^+ \wedge U^\infty \wedge U^\infty\}_{\mathbb{Z}_2} .
\end{aligned}$$

(v) Immediate from (iv).

(vi)+(vii) By construction.

□

Given a map $c : (X, Y) \rightarrow (O(V, U \oplus V), \{0\})$ define the \mathbb{Z}_2 -equivariant section of the trivial \mathbb{Z}_2 -equivariant bundle $\epsilon_{LU \oplus LV}$ over $S(LV) \times X$

$$c' : S(LV) \times X \rightarrow S(\epsilon_{LU \oplus LV}) = S(LV) \times X \times LU \oplus LV ; (v, x) \mapsto (v, x, c(x)(v)) .$$

The local obstruction of c is the rel $S(LV) \times Y$ \mathbb{Z}_2 -equivariant difference class

$$\begin{aligned} \theta(c) &= \delta(c', 0') \\ &\in \omega_{\mathbb{Z}_2}^{-1}(S(LV) \times (X, Y); -\epsilon_{LU \oplus LV}) = \{\Sigma S(LV)^+ \wedge X/Y; (LU \oplus LV)^\infty\}_{\mathbb{Z}_2} . \end{aligned}$$

Proposition 5.76. (Crabb [12, 5.13, 2.7])

(i) *The local obstruction determines a stable map*

$$\theta : O(V, U \oplus V) \rightarrow P(U \oplus V)/P(U)$$

inducing a function

$$\begin{aligned} \theta : [X/Y, O(V, U \oplus V)] &\rightarrow \{\Sigma S(LV)^+ \wedge X/Y; (LU \oplus LV)^\infty\}_{\mathbb{Z}_2} \\ &= \{X/Y; S(LU \oplus LV)/S(LU)\}_{\mathbb{Z}_2} \\ &= \{X/Y; S(LV)^+ \wedge LU^\infty\}_{\mathbb{Z}_2} \\ &= \{X/Y; P(U \oplus V)/P(U)\} \\ &= \omega_{\mathbb{Z}_2}^{-1}(S(LV) \times (X, Y); -\epsilon_{LU \oplus LV}) . \end{aligned}$$

The local obstruction of a map $c : X/Y \rightarrow O(V, U \oplus V)$ is a stable \mathbb{Z}_2 -equivariant map $\theta(c) : X/Y \rightarrow S(LV)^+ \wedge LU^\infty$ such that the composite with the \mathbb{Z}_2 -equivariant map

$$s_{LV} \wedge 1 : S(LV)^+ \wedge LU^\infty \rightarrow S^0 \wedge LU^\infty = LU^\infty$$

is $F_c - F_0 \in \{X/Y; LU^\infty\}_{\mathbb{Z}_2}$, with $F_c : LV^\infty \wedge X/Y \rightarrow (LU \oplus LV)^\infty$ the adjoint map (3.52) and 0 the constant map

$$0 : X/Y \rightarrow O(V, U \oplus V) ; x \mapsto (v \mapsto (0, v)) .$$

(ii) *The composite stable map*

$$\theta \circ R : P(U \oplus V)/P(U) \xrightarrow{R} O(V, U \oplus V) \xrightarrow{\theta} P(U \oplus V)/P(U)$$

is the identity, so that

$$\theta \circ R : [X/Y; P(U \oplus V)/P(U)] \rightarrow \{X/Y; P(U \oplus V)/P(U)\}$$

is the stabilization map. In particular, since $(O(V, U \oplus V), P(U \oplus V)/P(U))$ is $2 \dim(U)$ -connected and $O(V, U \oplus V)$ is $(\dim(U) - 1)$ -connected,

$$\theta : \pi_i(O(V, U \oplus V)) \rightarrow \pi_i(P(U \oplus V)/P(U))$$

is an isomorphism for $i \leq \dim(U)$, and

$$\theta : \omega_i(O(V, U \oplus V)) \rightarrow \omega_i(P(U \oplus V)/P(U))$$

is an isomorphism for $i \leq 2 \dim(U) - 1$.

(iii) If $c : (X, Y) \rightarrow (O(V), \{0\})$ is a map such that $X \setminus Y$ is a manifold and the \mathbb{Z}_2 -equivariant section of the trivial $S(LV) \times S(LV)$ -bundle over $S(LV) \times X$

$$\begin{aligned} d = (-j, c) : S(LV) \times X &\rightarrow S(LV) \times X \times S(LV) \times S(LV) ; \\ (v, x) &\mapsto (v, x, -v, c(x)(v)) \end{aligned}$$

is transverse regular at $S(LV) \times X \times \Delta_{S(LV)}$ then

$$\begin{aligned} C &= d^{-1}(S(LV) \times X \times \Delta_{S(LV)}) \\ &= \{(v, x) \mid c(x)(v) = -v \in S(LV)\} \subset S(LV) \times (X \setminus Y) \end{aligned}$$

is a submanifold of codimension $\dim(V) - 1$ with normal bundle

$$\nu_{C \subset S(LV) \times (X \setminus Y)} = (d|_C)^* \nu_{\Delta_{S(LV)} \subset S(LV) \times S(LV)} = (d|_C)^* \tau_{S(LV)}$$

such that $\nu_{C \subset S(LV) \times (X \setminus Y)} \oplus \epsilon_{\mathbb{R}} \cong \epsilon_{LV}$, and

$$\begin{aligned} \theta(c) : \Sigma S(LV)^+ \wedge X/Y &\xrightarrow{\Sigma F} \\ \Sigma T(\nu_{C \subset S(LV) \times (X \setminus Y)}) &= C^+ \wedge LV^\infty \longrightarrow LV^\infty \end{aligned}$$

with $F : S(LV)^+ \wedge X/Y \rightarrow T(\nu_{C \subset S(LV) \times (X \setminus Y)})$ the adjunction Umkehr map of the codimension 0 embedding $E(\nu_{C \subset S(LV) \times (X \setminus Y)}) \subset S(LV) \times (X \setminus Y)$.

Proof. (i) The local obstruction of $c \in O(V, U \oplus V)$ is the \mathbb{Z}_2 -equivariant pointed map

$$\theta(c) = \delta(F_c, F_0) : \Sigma S(LV)^+ \rightarrow (LU \oplus LV)^\infty$$

with $F_c : LV^\infty \rightarrow (LU \oplus LV)^\infty$ the \mathbb{Z}_2 -equivariant adjoint (4.81) and $0 \in O(V, U \oplus V)$ the linear isometry

$$0 : V \rightarrow U \oplus V ; v \mapsto (0, v) .$$

For any $c : X/Y \rightarrow O(V, U \oplus V)$ the local obstruction defines a \mathbb{Z}_2 -equivariant map

$$\theta(c) : \Sigma S(LV)^+ \wedge X/Y \rightarrow (LU \oplus LV)^\infty ; (v, x) \mapsto c(x)(v) .$$

The \mathbb{Z}_2 -equivariant S -duality isomorphism

$$\{\Sigma S(LV)^+ \wedge X/Y; (LU \oplus LV)^\infty\}_{\mathbb{Z}_2} \rightarrow \{X/Y; S(LU \oplus LV)/S(LU)\}_{\mathbb{Z}_2}$$

sends $\theta(c)$ to the composite stable \mathbb{Z}_2 -equivariant map

$$\begin{aligned} (LU \oplus LV)^\infty \wedge X/Y &\xrightarrow{\alpha_{LU \oplus LV} \wedge 1} \Sigma S(LU \oplus LV)^+ \wedge X/Y \\ &\xrightarrow{\Delta \wedge 1} \Sigma S(LV)^+ \wedge S(LU \oplus V)/S(LU) \wedge X/Y \\ &\xrightarrow{1 \wedge \theta(c)} (LU \oplus LV)^\infty \wedge S(L(U \oplus V))/S(LU) \end{aligned}$$

with

$$\begin{aligned} \Delta : \Sigma S(LU \oplus LV)^+ &\rightarrow \Sigma S(LV)^+ \wedge S(LU \oplus LV)/S(LU) ; \\ (t, u, v) &\mapsto ((t, \frac{v}{\|v\|}), [u, v]) . \end{aligned}$$

(ii) Consider first the special case $U = \{0\}$, $V = \mathbb{R}$. The reflection map is

$$R : P(\mathbb{R})^+ = \{*, [1]\} \rightarrow O(\mathbb{R}) = \{\pm 1\} ; * \mapsto 1 , [1] \mapsto -1 .$$

By Example 5.74 the local obstruction stable map $\theta : O(\mathbb{R}) \rightarrow P(\mathbb{R})^+$ induces

$$\begin{aligned} \theta : [S^0, O(\mathbb{R})] &= \{\pm 1\} \rightarrow \{\Sigma S(L\mathbb{R})^+; L\mathbb{R}^\infty\}_{\mathbb{Z}_2} = \pi_0^S(P(\mathbb{R})^+) = \mathbb{Z} ; \\ c &\mapsto \delta(c, 1) = \begin{cases} 0 & \text{if } c = 1 \\ 1 & \text{if } c = -1 . \end{cases} \end{aligned}$$

Thus $\theta \circ R \simeq \text{id} : P(\mathbb{R})^+ \rightarrow P(\mathbb{R})^+$. For arbitrary V and any $v \in P(V)$ there is defined a commutative diagram

$$\begin{array}{ccccc} P(\mathbb{R})^+ & \xrightarrow{R} & O(\mathbb{R}) & \xrightarrow{\theta} & P(\mathbb{R})^+ \\ \downarrow v & & \downarrow v & & \downarrow v \\ P(V)^+ & \xrightarrow{R} & O(V) & \xrightarrow{\theta} & P(V)^+ \end{array}$$

so that $\theta \circ R \simeq \text{id} : P(V)^+ \rightarrow P(V)^+$. Similarly for arbitrary U, V .

(iii) This is a \mathbb{Z}_2 -equivariant special case of 3.50 (vi).

□

Proposition 5.77. (Crabb [12, 2.9, 2.10]) *Let U, V be finite-dimensional inner product spaces, and let $\xi : X \rightarrow BO(U \oplus V)$ be a $U \oplus V$ -bundle.*

(i) The \mathbb{Z}_2 -equivariant cofibration sequence

$$S(LV)^+ \rightarrow S^0 \rightarrow LV^\infty \rightarrow \Sigma S(LV)^+$$

induces a long exact sequence of stable \mathbb{Z}_2 -cohomotopy groups

$$\begin{aligned} \cdots \rightarrow \omega_{\mathbb{Z}_2}^0(X; \epsilon_{LV} - L\xi) &\xrightarrow{-\otimes \gamma^{\mathbb{Z}_2}(\epsilon_{LV})} \omega_{\mathbb{Z}_2}^0(X; -L\xi) \\ &\rightarrow \omega^0(X \times P(V); -\xi \times H_{\mathbb{R}}) \rightarrow \cdots \end{aligned}$$

with $H_{\mathbb{R}}$ the Hopf \mathbb{R} -bundle over $P(V)$ (4.55) and

$$\omega^0(X \times P(V); -\xi \times H_{\mathbb{R}}) = \{T(L\eta); LU^\infty \wedge \Sigma S(LV)^+ \wedge LW^\infty\}_{\mathbb{Z}_2}$$

for any W -bundle $\eta : X \rightarrow BO(W)$ such that $\xi \oplus \eta = \epsilon_{U \oplus V \oplus W}$.

(ii) If $\xi \cong \xi' \oplus \epsilon_V$ for some U -bundle $\xi' : X \rightarrow BO(U)$ then

$$\gamma^{\mathbb{Z}_2}(L\xi) = \gamma^{\mathbb{Z}_2}(L\xi') \otimes \gamma^{\mathbb{Z}_2}(\epsilon_{LV}) \in \text{im}(\omega_{\mathbb{Z}_2}^0(X; \epsilon_{LV} - L\xi) \rightarrow \omega_{\mathbb{Z}_2}^0(X; -L\xi))$$

(iii) If X is an m -dimensional CW complex and $m < 2\dim(U)$ then $\xi \cong \xi' \oplus \epsilon_V$ for some U -bundle $\xi' : X \rightarrow BO(U)$ if and only if

$$\begin{aligned} \gamma^{\mathbb{Z}_2}(L\xi) &\in \text{im}(-\otimes \gamma^{\mathbb{Z}_2}(\epsilon_{LV}) : \omega_{\mathbb{Z}_2}^0(X; \epsilon_{LV} - L\xi) \rightarrow \omega_{\mathbb{Z}_2}^0(X; -L\xi)) \\ &= \ker(\omega_{\mathbb{Z}_2}^0(X; -L\xi) \rightarrow \omega^0(X \times P(V); -\xi \times H_{\mathbb{R}})) . \end{aligned}$$

□

Definition 5.78. Let $H(V)$ be the space of homotopy equivalences $h : S(V) \rightarrow S(V)$, regarded as a pointed space with base point $1 : S(V) \rightarrow S(V)$.

□

Following Crabb [12, §3] we shall now factor the local obstruction stable map $\theta : O(V) \rightarrow P(V)^+$ through the inclusion $J : O(V) \subset H(V)$.

Definition 5.79. (i) Let $H^{\mathbb{Z}_2}(LV \oplus W) \subset H(LV \oplus W)$ be the subspace of \mathbb{Z}_2 -equivariant homotopy equivalences $h : S(LV \oplus W) \rightarrow S(LV \oplus W)$.

(ii) Let $H^{\mathbb{Z}_2}(LV; W) \subset H^{\mathbb{Z}_2}(LV \oplus W)$ be the subspace of \mathbb{Z}_2 -equivariant homotopy equivalences $h : S(LV \oplus W) \rightarrow S(LV \oplus W)$ which restrict to the identity $h|_W = 1 : S(W) \rightarrow S(W)$ on the \mathbb{Z}_2 -fixed point set.

□

For any inner product spaces V, W the homeomorphism defined in Proposition 2.11 (ii)

$$\lambda_{LV,W} : S(LV) * S(W) \rightarrow S(LV \oplus W)$$

is \mathbb{Z}_2 -equivariant. The spaces of \mathbb{Z}_2 -equivariant homotopy equivalences fit into a fibration sequence

$$H^{\mathbb{Z}_2}(LV; W) \longrightarrow H^{\mathbb{Z}_2}(LV \oplus W) \xrightarrow{\rho} H(W)$$

with ρ the \mathbb{Z}_2 -fixed point map, which is split by

$$\sigma : H(W) \rightarrow H^{\mathbb{Z}_2}(LV \oplus W) ; g \mapsto \lambda_{LV,W}(1_{LV} * g)(\lambda_{LV,W})^{-1} .$$

For any (reasonable) pointed space X there is induced a split short exact sequence of homotopy groups

$$0 \longrightarrow [X, H^{\mathbb{Z}_2}(LV; W)] \longrightarrow [X, H^{\mathbb{Z}_2}(LV \oplus W)] \xrightarrow{\rho} [X, H(W)] \longrightarrow 0$$

Any map $g : X \rightarrow H^{\mathbb{Z}_2}(LV \oplus W)$ such that $\rho(g) : X \rightarrow H(W)$ is null-homotopic can be compressed (up to homotopy) to a map

$$X \rightarrow H^{\mathbb{Z}_2}(LV; W) \subset H^{\mathbb{Z}_2}(LV \oplus W) .$$

An element $h \in H^{\mathbb{Z}_2}(LV; W)$ suspends to a pointed \mathbb{Z}_2 -equivariant homotopy equivalence

$$sh : sS(LV \oplus W) = (LV \oplus W)^\infty \rightarrow sS(LV \oplus W) = (LV \oplus W)^\infty$$

which is the identity on $W^\infty \subset (LV \oplus W)^\infty$. Given a map $h : X \rightarrow H^{\mathbb{Z}_2}(LV; W)$ let

$$F_0, F_h : (LV \oplus W)^\infty \wedge X \rightarrow (LV \oplus W)^\infty$$

be the \mathbb{Z}_2 -equivariant maps defined by

$$F_0(v, w, x) = (v, w), F_h(v, w, x) = sh(x)(v, w)$$

such that $F_0(0, w, x) = F_h(0, w, x) = (0, w)$. Use the \mathbb{Z}_2 -equivariant difference map

$$\delta(F_h, F_0) : \Sigma S(LV)^+ \wedge W^\infty \wedge X \rightarrow LV^\infty \wedge W^\infty$$

to define a function

$$\begin{aligned} \zeta^{\mathbb{Z}_2} : [X, H^{\mathbb{Z}_2}(LV; W)] &\rightarrow \{\Sigma S(LV)^+ \wedge X; LV^\infty\}_{\mathbb{Z}_2} = \{X; P(V)^+\} ; \\ h &\mapsto \delta(F_h, F_0) . \end{aligned}$$

The homeomorphism $\lambda_{V,V} : S(V) * S(V) \cong S(V \oplus V)$ is \mathbb{Z}_2 -equivariant with respect to the transposition involutions

$$\begin{aligned} T &: S(V) * S(V) \rightarrow S(V) * S(V) ; (v, t, w) \mapsto (w, 1 - t, v) , \\ T &: S(V \oplus V) \rightarrow S(V \oplus V) ; (x, y) \mapsto (y, x) \end{aligned}$$

with \mathbb{Z}_2 -fixed point sets

$$\begin{aligned} (S(V) * S(V))^{\mathbb{Z}_2} &= \{(v, 1/2, v) \mid v \in S(V)\} , \\ S(V \oplus V)^{\mathbb{Z}_2} &= \{(\frac{x}{\sqrt{2}}, \frac{x}{\sqrt{2}}) \mid x \in S(V)\} . \end{aligned}$$

The \mathbb{Z}_2 -equivariant linear isometry

$$\kappa_V / \sqrt{2} : LV \oplus V \cong V \oplus V ; (v, w) \mapsto (\frac{v+w}{\sqrt{2}}, \frac{-v+w}{\sqrt{2}})$$

restricts to a \mathbb{Z}_2 -equivariant homeomorphism

$$\kappa_V / \sqrt{2} : S(LV \oplus V) \rightarrow S(V \oplus V) .$$

The composite

$$\begin{aligned} \mu_V &= \lambda_{V,V}(\kappa_V / \sqrt{2})\lambda_{LV,V}^{-1} : \\ &S(LV) * S(V) \rightarrow S(LV \oplus V) \rightarrow S(V \oplus V) \rightarrow S(V) * S(V) \end{aligned}$$

is a \mathbb{Z}_2 -equivariant homeomorphism which is the identity on the \mathbb{Z}_2 -fixed point sets. For any homotopy equivalence $h : S(V) \rightarrow S(V)$ the diagram of \mathbb{Z}_2 -equivariant maps

$$\begin{array}{ccc} S(LV) * S(V) & \xrightarrow{\mu_V} & S(V) * S(V) \\ 1 * h \downarrow & & \downarrow h * h \\ S(LV) * S(V) & \xrightarrow{\mu_V} & S(V) * S(V) \end{array}$$

does not commute, but does restrict to a commutative diagram on the \mathbb{Z}_2 -fixed point subset $S(V) \subset S(LV) * S(V)$. Choose a homotopy inverse $h^{-1} : S(V) \rightarrow S(V)$ of h , and define a \mathbb{Z}_2 -equivariant homotopy equivalence

$$(1 * h^{-1})\mu_V^{-1}(h * h)\mu_V : S(LV \oplus V) \rightarrow S(LV \oplus V)$$

which on the \mathbb{Z}_2 -fixed point sets is homotopic to $1 : S(V) \rightarrow S(V)$.

Definition 5.80. (Crabb [12, p.20])

(i) The *doubling function* is defined by

$$S^2 : [X, H(V)] \rightarrow [X, H^{\mathbb{Z}_2}(LV; V)] ; h \mapsto \mu_V^{-1}(h * h)\mu_V .$$

(ii) The *reduced doubling function*

$$\bar{S}^2 : [X, H(V)] \rightarrow [X, H^{\mathbb{Z}_2}(LV; V)] ; h \mapsto (1 * h^{-1})\mu_V^{-1}(h * h)\mu_V$$

is defined using a continuous choice of homotopy inverse $h^{-1} : X \rightarrow H(V)$, with

$$\begin{aligned} \bar{S}^2(h) &= (1 * h^{-1})\mu_V^{-1}(h * h)\mu_V \\ &\in [X, H^{\mathbb{Z}_2}(LV; V)] = \ker(\rho : [X, H^{\mathbb{Z}_2}(LV \oplus V)] \rightarrow [X, H(V)]) . \end{aligned}$$

For $g : X \rightarrow O(V)$ and $h = Jg : X \rightarrow H(V)$ the actual inverse can be chosen in (ii), and

$$\bar{S}^2(Jg) = g \oplus 1 \in \text{im}(J : [X, O(V)] \rightarrow [X, H^{\mathbb{Z}_2}(LV; V)]) .$$

(iii) Define the function

$$\eta = \zeta^{\mathbb{Z}_2} \bar{S}^2 : [X, H(V)] \rightarrow [X, H^{\mathbb{Z}_2}(LV; V)] \rightarrow \{X; P(V)^+\} .$$

□

Proposition 5.81. (Crabb [12, pp.23,27]) *Let X be a pointed space.*

(i) *There is defined a commutative diagram*

$$\begin{array}{ccc} [X, O(V)] & \xrightarrow{J_{\mathbb{Z}_2} F} & [X, H^{\mathbb{Z}_2}(LV)] \\ \downarrow J & \searrow \theta & \downarrow \zeta^{\mathbb{Z}_2} \\ [X, H(V)] & \xrightarrow{\eta} & \{X; P(V)^+\} \end{array}$$

with

$$J_{\mathbb{Z}_2} F : [X, O(V)] \rightarrow [X, H^{\mathbb{Z}_2}(LV)] ; g \mapsto g|_{S(LV)}$$

the forgetful map.

(ii) *Passing to the direct limit over all finite-dimensional inner product spaces V there is defined a commutative square*

$$\begin{array}{ccc} \widetilde{KO}^{-1}(X) = \varinjlim_V [X, O(V)] & \xrightarrow{J_{\mathbb{Z}_2} F} & \varinjlim_V [X, H^{\mathbb{Z}_2}(LV \oplus V)] \\ \theta \downarrow & & \downarrow \\ \{X; P(\infty)^+\} = \varinjlim_V \{X; P(V)^+\} & \longrightarrow & \tilde{\omega}_{\mathbb{Z}_2}^0(X) \end{array}$$

with

$$\begin{aligned}
J_{\mathbb{Z}_2} F : \widetilde{KO}^{-1}(X) &= \varinjlim_V [X, O(V)] \rightarrow \varinjlim_V [X, H^{\mathbb{Z}_2}(LV \oplus V)] ; \\
&\quad g \mapsto (g \oplus 1)|_{S(LV \oplus V)} , \\
\varinjlim_V [X, H^{\mathbb{Z}_2}(LV \oplus V)] &\rightarrow \widetilde{\omega}_{\mathbb{Z}_2}^0(X) = \varinjlim_V \{(LV \oplus V)^\infty \wedge X; (LV \oplus V)^\infty\}_{\mathbb{Z}_2} ; \\
&\quad h \mapsto F_h - F_0 \ (F_0 = 1)
\end{aligned}$$

and $\{X; P(\infty)^+\} \rightarrow \widetilde{\omega}_{\mathbb{Z}_2}^0(X)$ the injection in the direct sum system

$$\begin{aligned}
\{X; P(\infty)^+\} = \{X; S(\infty)^+\}_{\mathbb{Z}_2} &\xrightarrow[\delta]{\gamma} \{X; S^0\}_{\mathbb{Z}_2} = \widetilde{\omega}_{\mathbb{Z}_2}^0(X) \\
&\xrightarrow[\sigma]{\rho} \{X; S^0\} = \widetilde{\omega}^0(X) .
\end{aligned}$$

□

Chapter 6

The double point theorem

We apply the geometric Hopf invariant to a homotopy theoretic treatment of the double points of maps.

The connection between the Hopf invariant and double points has been much studied already, cf. Boardman and Steer [4], Dax [15], Eccles [16], Haefliger [19], Haefliger and Steer [20], Hatcher and Quinn [21], Koschorke and Sanderson [37],[38], Vogel [66], Wood [76]

6.1 Framed manifolds

Every m -dimensional manifold M admits an embedding $M \subset V \oplus \mathbb{R}^m$ for some inner product space V (and certainly if $\dim(V) > m$) with a normal vector V -bundle $\nu_M : M \rightarrow BO(V)$. The embedding $M \subset V \oplus \mathbb{R}^m$ extends to an open embedding $E(\nu_M) \subset V \oplus \mathbb{R}^m$ with compactification Umkehr map

$$(V \oplus \mathbb{R}^m)^\infty = V^\infty \wedge S^m \rightarrow T(\nu_M) .$$

Definition 6.1. (i) A *framed m -dimensional manifold* (M, b) is an m -dimensional manifold M together with an embedding $M \subset V \oplus \mathbb{R}^m$ and with an isomorphism $b : \nu_M \cong \epsilon_V$, or equivalently with an extension to an open embedding $b : V \times M \subset V \oplus \mathbb{R}^m$.

(ii) The *Pontryagin-Thom map* of (M, b) is the composite

$$\alpha_M : (V \oplus \mathbb{R}^m)^\infty = V^\infty \wedge S^m \rightarrow T(\nu_M) \xrightarrow[\cong]{T(b)} T(\epsilon_V) = V^\infty \wedge M^+ ,$$

representing an element

$$\alpha_M \in \omega_m(M) = \varinjlim_V [V^\infty \wedge S^m, V^\infty \wedge M^+] .$$

(iii) For any space X let $\Omega_m^{fr}(X)$ be the *framed bordism* group of pairs (M, b, f) with (M, b) a m -dimensional framed manifold and $f : M \rightarrow X$ a map.

□

As is well-known:

Proposition 6.2. *For any space X the Pontryagin-Thom construction defines an isomorphism*

$$\Omega_m^{fr}(X) \xrightarrow{\cong} \omega_m(X) ; (M, b, f) \mapsto (1 \wedge f)\alpha_M .$$

In particular,

$$(M, 1) = \alpha_M \in \Omega_m^{fr}(M) = \omega_m(M) .$$

□

Let $\eta(\mathbb{R}^m)$ be the canonical \mathbb{R}^m -bundle over the Stiefel space $V_{m+k,k} = O(\mathbb{R}^k, \mathbb{R}^{m+k})$ of linear isometries $u : \mathbb{R}^k \rightarrow \mathbb{R}^{m+k}$, with

$$\begin{aligned} E(\eta(\mathbb{R}^m)) &= \{(u, x) \mid u \in O(\mathbb{R}^k, \mathbb{R}^{m+k}), x \in u(\mathbb{R}^k)^\perp\} , \\ \delta\eta(\mathbb{R}^m) : \eta(\mathbb{R}^m) \oplus \epsilon_{\mathbb{R}^k} &\cong \epsilon_{\mathbb{R}^{m+k}} . \end{aligned}$$

Definition 6.3. (Kervaire [31])

Let (M, b) be a framed m -dimensional manifold, with an embedding $e : M \subset \mathbb{R}^{m+k}$ and an isomorphism $b : \nu_M \cong \epsilon_{\mathbb{R}^k}$.

(i) The *generalized Gauss map* of (M, b)

$$c : M \rightarrow V_{m+k,k} ; x \mapsto (de_x b^{-1} : \mathbb{R}^k \cong (\nu_M)_x \hookrightarrow (\tau_{\mathbb{R}^{m+k}})_{e(x)} = \mathbb{R}^{m+k})$$

classifies the tangent bundle of M

$$c^* \eta(\mathbb{R}^m) = \tau_M : M \rightarrow BO(\mathbb{R}^m)$$

and the stable trivialization

$$\delta\tau_M : \tau_M \oplus \epsilon_{\mathbb{R}^k} \xrightarrow[\cong]{1 \oplus b^{-1}} \tau_M \oplus \nu_M = \tau_{\mathbb{R}^{m+k}}|_M = \epsilon_{\mathbb{R}^{m+k}} .$$

(ii) The *curvatura integra* of (M, b) is the image of the fundamental class $[M] \in H_m(M)$ under the generalized Gauss map

$$c_*[M] \in H_m(V_{m+k,k}) = Q_{(-)^m}(\mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } m \equiv 0 \pmod{2} \\ \mathbb{Z}_2 & \text{if } m \equiv 1 \pmod{2} \end{cases}$$

(assuming $k > 1$).

(iii) The *Hopf invariant* of (M, b) is

$$\text{Hopf}(M, b) = \begin{cases} 0 \in \mathbb{Z} & \text{if } m \equiv 0 \pmod{2} \\ \text{Hopf}(p\alpha_M) \in \mathbb{Z}_2 & \text{if } m \equiv 1 \pmod{2} \end{cases}$$

with $\alpha_M : S^{m+k} \rightarrow \Sigma^k M^+$ the Pontryagin-Thom map of $b : \mathbb{R}^k \times M \subset \mathbb{R}^{m+k}$, and $p : M \rightarrow \{*\}$ the projection. Note that $\text{Hopf}(M, b) = 0$ for $m \neq 1, 3, 7$.

(iv) The *semicharacteristic* of an m -dimensional manifold M is defined for odd m by

$$\chi_{1/2}(M) = \sum_{i=0}^{(m-1)/2} \dim_{\mathbb{Z}_2} H_i(M; \mathbb{Z}_2) \in \mathbb{Z}_2 .$$

□

By Proposition 5.76 (ii) the local obstruction function θ (5.72) defines an isomorphism

$$\theta : H_m(V_{m+k,k}) \xrightarrow{\cong} H_m(P(\mathbb{R}^{m+k})/P(\mathbb{R}^m)) = Q_{2m}^{[0,k-1]}(S^m \mathbb{Z})$$

so that the curvatura integra is the image of $[M] \in H_m(M)$ under the composite

$$\theta(c) : M \xrightarrow{c} V_{m+k,k} \xrightarrow{\theta} P(\mathbb{R}^{m+k})/P(\mathbb{R}^m) = S(L\mathbb{R}^k)^+ \wedge_{\mathbb{Z}_2} (L\mathbb{R}^m)^\infty ,$$

that is

$$\begin{aligned} c_*[M] &= \theta(c)_*[M] \in H_m(V_{m+k,k}) = Q_{2m}^{[0,k-1]}(S^m \mathbb{Z}) \\ &= \begin{cases} \mathbb{Z} & \text{if } m \text{ is even or if } k = 1 \\ \mathbb{Z}_2 & \text{if } m \text{ is odd and } k \geq 2 . \end{cases} \end{aligned}$$

The zero section maps

$$z_{\tau_M} : M \rightarrow T(\tau_M) , \quad z_{\eta(\mathbb{R}^m)} : V_{m+k,k} \rightarrow T(\eta(\mathbb{R}^m))$$

are such that (for any m) there is defined a commutative diagram

$$\begin{array}{ccc}
H_m(M) & \xrightarrow{c_*} & H_m(V_{m+k,k}) \\
\downarrow z_M & & \downarrow z_{\eta(\mathbb{R}^m)} \\
\dot{H}_m(T(\tau_M)) = \mathbb{Z} & \xrightarrow[\cong]{c_*} & \dot{H}_m(T(\eta(\mathbb{R}^m))) = \mathbb{Z}
\end{array}$$

with

$$z_{\tau_M}([M]) = \chi(M), \quad z_{\eta(\mathbb{R}^m)}([S^m]) = \chi(S^m) = 1 + (-)^m \in \mathbb{Z}.$$

Thus for even m

$$c_*[M] = \chi(M)/2 \in H_m(V_{m+k,k}) = \mathbb{Z},$$

as originally proved by Hopf [25]. Kervaire [31, 32] expressed the curvatura integra for odd m in terms of the semicharacteristic and the Hopf invariant:

$$c_*[M] = \chi_{1/2}(M) - \text{Hopf}(M) \in H_m(V_{m+k,k}).$$

We shall reprove this in Proposition 6.8 below, following the outline of Crabb [12, Thm. 8.4] and using the quadratic construction on

$$\alpha_M = (M, 1) \in \omega_m(M) = \Omega_m^{fr}(M)$$

which is a stable homotopy theoretic version of the quadratic refinement μ of the intersection form λ on $H^*(M)$.

Proposition 6.4. (Pontryagin [46] for $n = 1$, Kervaire and Milnor [35] for $(n-1)$ -connected M^{2n} , Browder [6], [7, §III.4] in general.)

Let (M, b) be an m -dimensional framed manifold, with an embedding $\mathbb{R}^k \times M \subset \mathbb{R}^{m+k}$ and Pontryagin-Thom map $\alpha_M : S^{m+k} \rightarrow \Sigma^k M^+$.

(i) Suppose that $m = 2n$, and define the ring

$$A_n = \begin{cases} \mathbb{Z} & \text{if } n \equiv 0 \pmod{2} \\ \mathbb{Z}_2 & \text{if } n \equiv 1 \pmod{2} \end{cases}$$

so that there is a $(-)^n$ -symmetric intersection pairing over A_n

$$\lambda : H^n(M; A_n) \times H^n(M; A_n) \rightarrow A_n; \quad (x, y) \mapsto \langle x \cup y, [M] \rangle$$

with

$$\lambda(x, y) = (-)^n \lambda(y, x) \in A_n.$$

The framing of M determines a $(-)^n$ -quadratic refinement of λ

$$\mu : H^n(M; A_n) \rightarrow Q_{(-)^n}(A_n) = \mathbb{Z}/\{1 - (-)^n\} ;$$

$$x \mapsto \begin{cases} \lambda(x, x)/2 & \text{if } n \equiv 0 \pmod{2} \\ \langle Sq_{xF}^{n+1}(\iota), [S^{2n}] \rangle & \text{if } n \equiv 1 \pmod{2} \end{cases} ,$$

with $\iota \in H^n(K(\mathbb{Z}_2, n); \mathbb{Z}_2) = \mathbb{Z}_2$ the generator, $xF \in \pi_{2n}^S(K(\mathbb{Z}_2, n))$, such that for all $x, y \in H^n(M; A_n)$, $a \in A_n$

$$\lambda(x, x) = \mu(x) + (-)^n \mu(x) \in A_n ,$$

$$\mu(ax) = a^2 \mu(x) , \lambda(x, y) = \mu(x + y) - \mu(x) - \mu(y) \in Q_{(-)^n}(A_n) .$$

(ii) Let $m = 2n$ as in (i), and suppose that $x \in H^n(M; A_n)$ is the Poincaré dual of the homology class $x[S^n] \in H_n(M; A_n)$ represented by an embedding $x : S^n \subset M$ (as is the case for all $x \in H^n(M; A_n)$ if M is $(n-1)$ -connected with $n \geq 3$), with normal bundle $\nu_x : S^n \rightarrow BO(\mathbb{R}^n)$ and geometric Umkehr map $G : M^+ \rightarrow T(\nu_x)$. The normal bundle of the composite $S^n \subset M^{2n} \subset \mathbb{R}^{2n+k}$ has a canonical trivialization (for large k)

$$\delta\nu_x : \nu_{S^n \subset \mathbb{R}^{2n+k}} = \nu_x \oplus \epsilon_{\mathbb{R}^k} \cong \epsilon_{\mathbb{R}^{n+k}} ,$$

such that the corresponding Pontryagin-Thom map

$$S^{2n+k} \xrightarrow{F} \Sigma^k M^+ \xrightarrow{\Sigma^k G} \Sigma^k T(\nu_x) \xrightarrow[\cong]{\delta\nu_x} \Sigma^k T(\epsilon^n) \rightarrow S^{n+k}$$

is null-homotopic. Then:

(a) The evaluation of the quadratic form μ on $x \in H^n(M; A_n)$ is

$$\mu(x) = (\delta\nu_x, \nu_x) \in \pi_n(V_{n+k, k}) = Q_{(-)^n}(A_n) .$$

(b) The exact sequence

$$\cdots \rightarrow \pi_n(O(\mathbb{R}^n)) \rightarrow \pi_n(O) \rightarrow Q_{(-)^n}(A_n) \rightarrow \pi_n(BO(\mathbb{R}^n)) \rightarrow \pi_n(BO) \rightarrow \cdots$$

is such that

$$Q_{(-)^n}(A_n) = \mathbb{Z}/\{1 + (-)^{n+1}\} \rightarrow \pi_n(BO(\mathbb{R}^n)) ; 1 \mapsto \tau_{S^n}$$

is injective for $n \notin \{1, 3, 7\}$ and is 0 for $n \in \{1, 3, 7\}$. Thus if $n \notin \{1, 3, 7\}$

$$\mu(x) = \begin{cases} \chi(\nu_x)/2 \in \mathbb{Z} = \ker(\pi_n(BO(\mathbb{R}^n)) \rightarrow \pi_n(BO)) & \text{if } n \equiv 0 \pmod{2} \\ \nu_x \in \mathbb{Z}_2 = \ker(\pi_n(BO(\mathbb{R}^n)) \rightarrow \pi_n(BO)) & \text{if } n \equiv 1 \pmod{2} \end{cases}$$

with $\mu(x) = 0$ if and only if $\nu_x \cong \epsilon_{\mathbb{R}^n}$.

(c) If $\nu_x = 0 \in \pi_n(BO(\mathbb{R}^n))$ (e.g. if $n \in \{1, 3, 7\}$ and ν_x is orientable) then for any choice of trivialization $\delta\nu'_x : \nu_x \cong \epsilon_{\mathbb{R}^n}$ the normal bundle of $S^n \subset M^{2n} \subset \mathbb{R}^{2n+k}$ has a trivialization

$$\delta\nu'_x \oplus \epsilon_{\mathbb{R}^k} : \nu_{S^n \subset \mathbb{R}^{2n+k}} = \nu_x \oplus \epsilon_{\mathbb{R}^k} \cong \epsilon_{\mathbb{R}^{n+k}} ,$$

with the Pontryagin-Thom map

$$S^{2n+k} \xrightarrow{F} \Sigma^k M^+ \xrightarrow{\Sigma^k G} \Sigma^k T(\nu_x) \xrightarrow[\cong]{\Sigma^k \delta\nu'_x} \Sigma^k T(\epsilon^n) \rightarrow S^{n+k}$$

such that

$$\mu(x) = \text{Hopf}(S^n, \delta\nu'_x \oplus \epsilon_{\mathbb{R}^k}) \in \text{im}(\pi_n(BO) \rightarrow Q_{(-)^n}(A_n)) .$$

(d) If $n \neq 2$ there exists a choice of trivialization $\delta\nu'_x : \nu_x \cong \epsilon_{\mathbb{R}^n}$ such that $x[S^n] \in H_n(M; A_n)$ can be killed by a framed surgery on M if and only if $\mu(x) = 0$.

(iii) For any m , the framing determines a bundle map $b : \nu_M \rightarrow \epsilon_{\mathbb{R}^k}$ over a degree 1 map $f : M \rightarrow S^m$, i.e. a normal map $(f, b) : M \rightarrow S^m$. The surgery obstruction of (f, b) is

$$\sigma_*(f, b) = \begin{cases} \text{signature}(H^n(M), \lambda)/8 \\ \text{Arf invariant}(H^n(M; \mathbb{Z}_2), \lambda, \mu) \\ 0 \end{cases}$$

$$\in L_m(\mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } m = 2n, \ n \equiv 0 \pmod{2} \\ \mathbb{Z}_2 & \text{if } m = 2n, \ n \equiv 1 \pmod{2} \\ 0 & \text{if } m \equiv 1 \pmod{2} . \end{cases}$$

□

Remark 6.5. In Ranicki [50] the quadratic function $\mu : H^n(M) \rightarrow Q_{(-)^n}(\mathbb{Z})$ of a $2n$ -dimensional framed manifold M was shown to be a special case of a quadratic structure on the chain complex $C(M)$ of an m -dimensional framed manifold M (for any m) obtained by the quadratic construction on the Pontryagin-Thom map $\alpha_M : V^\infty \wedge S^m \rightarrow V^\infty \wedge M^+$ of an embedding $V \times M \subset V \oplus \mathbb{R}^m$, namely

$$\psi = \psi(\alpha_M) \in H_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) = Q_m(C(M)) .$$

For any oriented m -dimensional manifold M there is defined a degree 1 map

$$f : M \rightarrow M/(M \setminus D^m) = D^m/S^{m-1} = S^m$$

such that $f : C(M) \rightarrow C(S^m)$ is a chain homotopy surjection split by the Umkehr chain map

$$f^! : C(S^m) \simeq C(S^m)^{m-*} \xrightarrow{f^*} C(M)^{m-*} \simeq C(M) ,$$

so that there is defined a chain homotopy direct sum system

$$C(S^m) \begin{array}{c} \xrightarrow{f^!} \\ \xleftarrow{f} \end{array} C(M) \begin{array}{c} \xrightarrow{e} \\ \xleftarrow{\quad} \end{array} \mathcal{C}(f^!)$$

with $e : C(M) \rightarrow \mathcal{C}(f^!)$ the inclusion in the algebraic mapping cone. A framing of M determines a bundle map $b : \nu_M \rightarrow \epsilon_V$ over f , i.e. a normal map $(f, b) : M \rightarrow S^m$. The S -dual of $T(b) : T(\nu_M) \rightarrow T(\epsilon_V)$ is a geometric Umkehr map $F : \Sigma^\infty(S^m)^+ \rightarrow \Sigma^\infty M^+$ with $(\Sigma^\infty f)F \simeq 1$, such that

$$\alpha_M : \Sigma^\infty(S^m) \rightarrow \Sigma^\infty(S^m)^+ \xrightarrow{F} \Sigma^\infty M^+ .$$

The geometric Umkehr F induces the chain Umkehr $f^!$. The quadratic structure

$$\psi = \psi(\alpha_M) \in H_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) = Q_m(C(M)) ,$$

is such that $(\mathcal{C}(f^!), e_{\%}(\psi))$ is an m -dimensional quadratic Poincaré complex over \mathbb{Z} representing the simply-connected surgery obstruction

$$\sigma_*(f, b) = (\mathcal{C}(f^!), e_{\%}(\psi)) \in L_m(\mathbb{Z}) .$$

□

For any space M Proposition 4.41 (ii) gives a direct sum system

$$\omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) \begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \omega_m^{\mathbb{Z}_2}(M \times M) \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \omega_m(M) .$$

The quadratic construction function (5.46)

$$\psi : \omega_m(M) \rightarrow \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) ; (\alpha : \Sigma^\infty \wedge S^m \rightarrow \Sigma^\infty M^+) \mapsto \psi(\alpha)$$

is such that

$$\begin{aligned} \gamma\psi(\alpha) &= (\alpha \wedge \alpha)\Delta_{S^m} - \Delta_M(\alpha) \in \omega_m^{\mathbb{Z}_2}(M \times M) , \\ \psi(\alpha + \beta) &= \psi(\alpha) + \psi(\beta) + (\alpha \wedge \beta)\Delta_{S^m} \in \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) \end{aligned}$$

with $(\alpha \wedge \beta)\Delta_{S^m}$ defined using the map induced by $S(L\mathbb{R}) \hookrightarrow S(\infty)$

$$\omega_m(M \times M) = \omega_m(S(L\mathbb{R}) \times_{\mathbb{Z}_2} (M \times M)) \rightarrow \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) .$$

Let $p : M \rightarrow \{*\}$ be the projection, and let

$q = (1 \times p \times p) : S(\infty) \times_{\mathbb{Z}_2} (M \times M) \rightarrow S(\infty) \times_{\mathbb{Z}_2} (\{*\} \times \{*\}) = P(\infty)$,

so that there is defined a commutative diagram

$$\begin{array}{ccc}
 \omega_m(M) & \xrightarrow{p_*} & \omega_m \\
 \downarrow \psi & & \downarrow \bar{P}^2 \\
 \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) & \xrightarrow{q_*} & \omega_m(P(\infty)) \\
 \downarrow H & & \downarrow H \\
 Q_m(C(M)) & \xrightarrow{q_*} & Q_m(\mathbb{Z}) = H_m(P(\infty))
 \end{array}$$

with H the Hurewicz maps, and \bar{P}^2 the reduced squaring operation (5.44 (ii)).

Proposition 6.6. (Crabb [12, p.45])

(i) *The quadratic construction on the Pontryagin-Thom map $\alpha_M : \Sigma^\infty S^m \rightarrow \Sigma^\infty M^+$ of an m -dimensional framed manifold (M, b) is a stable homotopy class*

$$\psi(\alpha_M) \in \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M))$$

with Hurewicz image the quadratic construction $\psi \in Q_m(C(M))$ of [50], defining a quadratic function

$$\begin{aligned}
 \psi : \Omega_m^{fr}(X) &= \omega_m(X) \rightarrow \omega_m(S(\infty) \times_{\mathbb{Z}_2} (X \times X)) ; \\
 (M, b, f : M \rightarrow X) &\mapsto (1 \times f \times f)_* \psi(\alpha_M)
 \end{aligned}$$

for any space X .

(ii) *The bordism class $\alpha_M = (M, 1) \in \omega_m(M) = \Omega_m^{fr}(M)$ has images*

$$\begin{aligned}
 p\alpha_M &= (M, b) \in \omega_m = \Omega_m^{fr} , \\
 \psi(\alpha_M) &\in \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) , \\
 H(\psi(\alpha_M)) &\in Q_m(C(M)) , \\
 q_* H\psi(\alpha_M) &= H\bar{P}^2(p\alpha_M) = \text{Hopf}(M) \\
 &\in Q_m(\mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } m = 0 \\ 0 & \text{if } m > 0 \text{ and } m \equiv 0 \pmod{2} \\ \mathbb{Z}_2 & \text{if } m \equiv 1 \pmod{2} . \end{cases}
 \end{aligned}$$

Proof. By construction. □

Proposition 6.7. (Milnor and Stasheff [43, §11], Crabb [12, p.92])

Let N be an n -dimensional manifold, so that $N \times N$ is a $2n$ -dimensional manifold, and $\Delta_N : N \hookrightarrow N \times N$ is the embedding of an n -dimensional submanifold with normal bundle

$$\nu_{\Delta_N} = \tau_N : N \rightarrow BO(\mathbb{R}^n) .$$

(i) If N is oriented then so is $N \times N$, and Δ_N represents a homology class $[N] \in H_n(N \times N)$ with a Poincaré dual $[N]^* \in H^n(N \times N)$. The $(-)^n$ -symmetric intersection form $\lambda_{N \times N}$ on $H^n(N \times N)$ is such that

$$\lambda_{N \times N}([N]^*, [N]^*) = \chi(N) \in H_n(N) = \mathbb{Z} ,$$

with $\chi(N) = 0$ for $n \equiv 1 \pmod{2}$.

(ii) If N is framed then so is $N \times N$, and the $(-)^n$ -quadratic function $\mu_{N \times N}$ on $H^n(N \times N)$ is such that

$$\mu_{N \times N}([N]^*) = \begin{cases} \chi(N)/2 & \text{if } n \equiv 0 \pmod{2} \\ \chi_{1/2}(N) & \text{if } n \equiv 1 \pmod{2} \end{cases} \in Q_{(-)^n}(\mathbb{Z})$$

Proof. (i) Let F be a field, and work with F -coefficients. By the Künneth formula

$$H^n(N \times N; F) = \sum_{i=0}^n H^i(N; F) \otimes_F H^{n-i}(N; F)$$

with

$$\lambda_{N \times N} \left(\sum_{j=1}^k x_j \otimes y_j, \sum_{j'=1}^{k'} x'_{j'} \otimes y'_{j'} \right) = \sum_{j=1}^k \sum_{j'=1}^{k'} \lambda_N(x_j, x'_{j'}) \lambda_N(y_j, y'_{j'}) \in F .$$

Choose a basis b_1, b_2, \dots, b_r for $H^*(N; F)$, and let $b_1^*, b_2^*, \dots, b_r^*$ be the dual basis for $H^*(N; F)$ with

$$\lambda(b_p, b_q^*) = \begin{cases} 1 & \text{if } p = q \\ 0 & \text{if } p \neq q . \end{cases}$$

See [43, pp. 127–130] for the proof that

$$\begin{aligned} [N]^* &= \sum_{q=1}^r (-)^{\dim(b_q)} b_q \otimes b_q^* \\ &\in H^n(N \times N; F) = \sum_{i=0}^n H^i(N; F) \otimes_F H^{n-i}(N; F) \end{aligned}$$

with

$$\lambda_{N \times N}([N]^*, [N]^*) = \sum_{q=1}^r (-)^{\dim(b_q)} = \chi(N) \in \text{im}(\mathbb{Z} \rightarrow F) .$$

(ii) For $n \equiv 0 \pmod{2}$ it is immediate from (i) and

$$\lambda_{N \times N}([N]^*, [N]^*) = 2\mu_{N \times N}([N]^*) \in \mathbb{Z}$$

that $\mu_{N \times N}([N]^*) = \chi(N)/2$.

So suppose that $n \equiv 1 \pmod{2}$, and work with \mathbb{Z}_2 -coefficients. By the Künneth formula

$$H^n(N \times N; \mathbb{Z}_2) = \sum_{i=0}^n H^i(N; \mathbb{Z}_2) \otimes_{\mathbb{Z}_2} H^{n-i}(N; \mathbb{Z}_2)$$

with

$$\begin{aligned} \lambda_{N \times N} \left(\sum_{j=1}^k x_j \otimes y_j, \sum_{j'=1}^{k'} x'_{j'} \otimes y'_{j'} \right) &= \sum_{j=1}^k \sum_{j'=1}^{k'} \lambda_N(x_j, x'_{j'}) \lambda_N(y_j, y'_{j'}) \in \mathbb{Z}_2 , \\ \mu_{N \times N} \left(\sum_{j=1}^k x_j \otimes y_j \right) &= \sum_{1 \leq j < j' \leq k} \lambda_N(x_j, x_{j'}) \lambda_N(y_j, y_{j'}) \in Q_{(-)^n}(\mathbb{Z}_2) = \mathbb{Z}_2 . \end{aligned}$$

Continuing with the terminology of (i) (with $F = \mathbb{Z}_2$) we have

$$\mu_{N \times N}([N]^*) = \sum_{\dim(b_q) < \dim(b_q^*)} 1 = \chi_{1/2}(N) \in \mathbb{Z}_2 .$$

□

Proposition 6.8. (Kervaire [31, 32], Crabb [12, Lemma 8.5])

(i) Let (M^{2n}, b) be a $2n$ -dimensional framed manifold, so that there is given an embedding $M \subset \mathbb{R}^{2n+k}$ with a trivialized normal \mathbb{R}^k -bundle $b : \nu_{M \subset \mathbb{R}^{2n+k}} \cong \epsilon_{\mathbb{R}^k}$. Let $N^n \subset M^{2n}$ be an n -dimensional submanifold such that the normal bundle of the composite embedding $N \subset M \subset \mathbb{R}^{2n+k}$ has is equipped with a trivialization $a : \nu_{N \subset \mathbb{R}^{2n+k}} \cong \epsilon_{\mathbb{R}^{n+k}}$. Let $c : N \rightarrow V_{n+k, k}$ classify the normal \mathbb{R}^n -bundle $\nu_{N \subset M} : N \rightarrow BO(\mathbb{R}^n)$ with the corresponding stable trivialization

$$\delta \nu_{N \subset M} : \nu_{N \subset M} \oplus \epsilon_{\mathbb{R}^k} \cong \epsilon_{\mathbb{R}^{n+k}} .$$

The Hopf invariant of (N, a) is given by

$$\text{Hopf}(N, a) = \mu([N]^*) - c_*[N] \in H_n(V_{n+k, k}) = Q_{(-)^n}(\mathbb{Z})$$

with $\mu([N]^*)$ the evaluation of the quadratic function μ on the Poincaré dual $[N]^* \in H^n(M)$ of $[N] \in H_n(M)$. (Note that $\text{Hopf}(N, a) = 0$ for $n \neq 1, 3, 7$, or if $N = S^n$ with the canonical framing of the normal bundle of $S^n \subset M^{2n} \subset \mathbb{R}^{2n+k}$.)

(ii) Let (N, a) be an n -dimensional framed manifold, so that there is given an embedding $N \subset \mathbb{R}^{n+k}$ with a trivialized normal bundle. Then

$$\text{Hopf}(N, a) = \chi_{1/2}(N) - c_*[N] \in H_n(V_{n+k, k}) = Q_{(-)^n}(\mathbb{Z})$$

with $c : N \rightarrow V_{n+k, k}$ classifying the tangent \mathbb{R}^n -bundle $\tau_N : N \rightarrow BO(\mathbb{R}^n)$ with the corresponding stable isomorphism

$$a : \tau_N \oplus \epsilon_{\mathbb{R}^k} \cong \epsilon_{\mathbb{R}^{n+k}} .$$

(iii) As in (i) let $N^n \subset M^{2n} \subset \mathbb{R}^{2n+k}$ with trivializations $b : \nu_{M \subset \mathbb{R}^{2n+k}} \cong \epsilon_{\mathbb{R}^k}$, $a : \nu_{N \subset M^{2n}} \cong \epsilon_{\mathbb{R}^n}$, and let $c : N \rightarrow V_{2n+k, n+k}$ classify $\tau_N : N \rightarrow BO(\mathbb{R}^n)$ with the corresponding stable trivialization $\tau_N \oplus \epsilon_{\mathbb{R}^{n+k}} \cong \epsilon_{\mathbb{R}^{2n+k}}$. The evaluation of $\mu : H^n(M) \rightarrow Q_{(-)^n}(\mathbb{Z})$ on $[N]^* \in H^n(M)$ is

$$\begin{aligned} \mu([N]^*) &= \text{Hopf}(N, a) = \chi_{1/2}(N) - c_*[N] \quad (= 0 \text{ for } n \neq 1, 3, 7) \\ &\in H_n(V_{2n+k, n+k}) = Q_{(-)^n}(\mathbb{Z}) . \end{aligned}$$

In particular, if $N = S^n$ then $\chi_{1/2}(N) = 1$ and

$$\mu([S^n]^*) = \text{Hopf}(S^n, \text{std.}) = 1 - c_*[S^n] \in H_n(V_{2n+k, n+k}) = Q_{(-)^n}(\mathbb{Z}) .$$

Proof. (i) Let $V = \mathbb{R}^k$ and let

$$\alpha_M : V^\infty \wedge (\mathbb{R}^{2n})^\infty \rightarrow V^\infty \wedge M^+$$

be the Pontryagin-Thom map of $V \times M \subset V \oplus \mathbb{R}^{2n}$, so that $\psi(\alpha_M) \in \omega_{2n}(S(\infty) \times_{\mathbb{Z}_2} (M \times M))$ determines the $(-)^n$ -quadratic form μ on $H^n(M)$ refining the $(-)^n$ -symmetric intersection form λ . Let

$$\alpha_{N \subset M} : M^+ \rightarrow M/(M \setminus E(\nu_{N \subset M})) = T(\nu_{N \subset M})$$

be the Umkehr map of $N \subset M$. Apply the composition formula for the quadratic construction to the factorization

$$\begin{array}{ccc} & V^\infty \wedge M^+ & \\ \alpha_M \nearrow & \downarrow 1 \wedge \alpha_{N \subset M} & \\ V^\infty \wedge S^{2n} & V^\infty \wedge T(\nu_{N \subset M}) & \\ \alpha_N \searrow & \downarrow T(\delta \nu_{N \subset M}) & \\ & (V \oplus \mathbb{R}^n)^\infty \wedge N^+ & \end{array}$$

to obtain

$$\begin{aligned} \psi(\alpha_N) &= (T(\delta\nu_{N \subset M})(1 \wedge \alpha_{N \subset M}))\% \psi(\alpha_M) + \psi(T(\delta\nu_{N \subset M}))(1 \wedge \alpha_{N \subset M})\alpha_M \\ &: H_{2n}(S^{2n}) \rightarrow Q_{2n}(S^n C(M)) . \end{aligned}$$

Let $p : N \rightarrow \{*\}$ be the projection. Now

$$\begin{aligned} p\% \psi(\alpha_N)[S^{2n}] &= \text{Hopf}(N) , \\ p\%(T(\delta\nu_{N \subset M})(1 \wedge \alpha_{N \subset M}))\% \psi(\alpha_M)[S^{2n}] &= \mu([N]^*) , \\ p\% \psi(T(\delta\nu_{N \subset M}))(1 \wedge \alpha_{N \subset M})\alpha_M[S^{2n}] &= -c_*[N] \in Q_{2n}(S^n \mathbb{Z}) \quad (5.75) , \end{aligned}$$

so that

$$\text{Hopf}(N) = \mu([N]^*) - c_*[N] \in Q_{2n}(S^n \mathbb{Z}) = Q_{(-)^n}(\mathbb{Z}) .$$

(ii) Apply (i) with $\Delta_N : N \subset M = N \times N$, $\nu_{\Delta_N} = \tau_N$, using 6.7 to identify $\mu([N]^*) = \chi_{1/2}(N)$.

(iii) Immediate from (i), (ii) and Proposition 6.4 (ii) (c). □

Example 6.9. The embeddings

$$x : S^n \times \{*\} \subset S^n \times S^n , \quad y : \{*\} \times S^n \subset S^n \times S^n$$

have trivial normal \mathbb{R}^n -bundles $\nu_x = \nu_y = \epsilon^n$. For $n = 1, 3, 7$ there exist framings b of $S^n \times S^n \subset \mathbb{R}^{2n+k}$ such that the corresponding quadratic function

$$\mu_b : H^n(S^n \times S^n; \mathbb{Z}_2) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$$

takes values

$$\mu_b(x) = \mu_b(y) = 1 \in \mathbb{Z}_2$$

with the corresponding framings of $x, y : S^n \subset \mathbb{R}^{2n+k}$ such that

$$\text{Hopf}(S^n, x) = \text{Hopf}(S^n, y) = 1 \in \mathbb{Z} .$$

□

Example 6.10. Given a map $\omega : S^m \rightarrow O(k)$ the embedding

$$e_\omega : S^m \times D^k \hookrightarrow S^{m+k} = S^m \times D^k \cup D^{m+1} \times S^{k-1} ; (x, y) \mapsto (x, \omega(x)(y))$$

defines a submanifold $S^m \subset S^{m+k}$ with a framing b_ω such that the generalized Gauss map is given by

$$c : S^m \rightarrow V_{m+k,k} ; x \mapsto (y \mapsto (x, \omega(x)(y)))$$

and the Pontryagin-Thom map is given by the J -homomorphism

$$J(\omega) : S^{m+k} \rightarrow S^k ; (x, y) \mapsto \begin{cases} \omega(x)(y) & \text{if } (x, y) \in S^m \times D^k \\ * & \text{if } (x, y) \in D^{m+1} \times S^{k-1} . \end{cases}$$

The curvatura integra is

$$c_*[S^m] = \begin{cases} \chi(S^m)/2 = 1 \\ \chi_{1/2}(S^m) - \text{Hopf}(J(\omega)) = 1 - \text{Hopf}(J(\omega)) \end{cases}$$

$$\in H_m(V_{m+k,k}) = \begin{cases} \mathbb{Z} & \text{if } m \text{ is even} \\ \mathbb{Z}_2 & \text{if } m \text{ is odd} \end{cases}$$

with $\text{Hopf}(J(\omega)) = 0$ for $m \neq 1, 3, 7$.

□

An m -dimensional framed manifold M determines a normal map $(f, b) : M \rightarrow S^m$ with a surgery obstruction $\sigma_*(f, b) \in L_m(\mathbb{Z})$. Following Crabb [12] we shall now obtain a geometric realization of the surgery obstruction by expressing the quadratic construction

$$\psi(\alpha_M) \in \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M))$$

(in Example 6.14 below) as a bordism class of an m -dimensional framed manifold N with a map $g : N \rightarrow S(\infty) \times_{\mathbb{Z}_2} (M \times M)$ such that

$$(N, g) = (M, \Delta_M) - (S^m, (\alpha_M \wedge \alpha_M) \Delta_{S^m}) = -\psi(\alpha_M)$$

$$\in \ker(\rho : \omega_m^{\mathbb{Z}_2}(M \times M) \rightarrow \omega_m(M)) = \omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) ,$$

and

$$(\overline{N}, \overline{g}) = (M, \Delta_M) \in \omega_m(M \times M)$$

with $\overline{N} = g^*(S(\infty) \times (M \times M))$ the canonical double cover of N and

$$\overline{g} : \overline{N} \rightarrow S(\infty) \times (M \times M) \rightarrow M \times M .$$

The construction uses the \mathbb{Z}_2 -equivariant S -duality analogue of:

Proposition 6.11. *An m -dimensional framed manifold (M, b) is self- S -dual, in the sense that the composite*

$$\Delta \alpha_M : (V \oplus \mathbb{R}^m)^\infty \rightarrow V^\infty \wedge M^+ \rightarrow M^+ \wedge T(\epsilon_V) = M^+ \wedge (V^\infty \wedge M^+)$$

is an S -duality map between M^+ and $V^\infty \wedge M^+$. Cap product with the stable homotopy fundamental class

$$[M] = (M, 1) = \alpha_M \in \omega_m(M)$$

defines Poincaré duality isomorphisms in stable homotopy

$$[M] \cap - : \omega^*(M) \cong \omega_{m-*}(M) .$$

In particular $[M] \cap - : \omega^0(M) \cong \omega_m(M)$ sends $1 \in \omega^0(M)$ to $[M] \in \omega_m(M)$. \square

And now for the \mathbb{Z}_2 -analogues. Every m -dimensional \mathbb{Z}_2 -manifold M admits a \mathbb{Z}_2 -equivariant embedding $i : M \subset LV \oplus W \oplus \mathbb{R}^m$ for some inner product spaces V, W with a normal \mathbb{Z}_2 -equivariant vector $LV \oplus W$ -bundle ν_M , and i extends to an open \mathbb{Z}_2 -equivariant embedding $i : E(\nu_M) \subset LV \oplus W \oplus \mathbb{R}^m$.

Definition 6.12. (i) An m -dimensional \mathbb{Z}_2 -manifold M is \mathbb{Z}_2 -framed if there is given a \mathbb{Z}_2 -embedding $i : M \subset LV \oplus W \oplus \mathbb{R}^m$ with an isomorphism $\nu_M \cong \epsilon_{LV \oplus W}$, in which case i extends to an open \mathbb{Z}_2 -equivariant embedding

$$i : LV \oplus W \times M \subset LV \oplus W \oplus \mathbb{R}^m .$$

(ii) For any \mathbb{Z}_2 -space let $\Omega_m^{\mathbb{Z}_2\text{-}fr}(X)$ be the \mathbb{Z}_2 -framed bordism group of pairs (M, f) with M an m -dimensional \mathbb{Z}_2 -framed \mathbb{Z}_2 -manifold and $f : M \rightarrow X$ a \mathbb{Z}_2 -equivariant map.

(iii) The \mathbb{Z}_2 -equivariant Pontryagin-Thom map of a \mathbb{Z}_2 -framed \mathbb{Z}_2 -manifold M is the compactification Umkehr map of $i : LV \oplus W \times M \subset LV \oplus W \oplus \mathbb{R}^m$, a stable \mathbb{Z}_2 -equivariant map

$$\begin{aligned} \alpha_M^{\mathbb{Z}_2} : (LV \oplus W \oplus \mathbb{R}^m)^\infty &= LV^\infty \wedge W^\infty \wedge S^m \\ &\rightarrow T(\epsilon_{LV \oplus W}) = LV^\infty \wedge W^\infty \wedge M^+ \end{aligned}$$

representing an element

$$\alpha_M^{\mathbb{Z}_2} \in \omega_m^{\mathbb{Z}_2}(M) = \varinjlim_{V, W} [(LV \oplus W \oplus \mathbb{R}^m)^\infty, LV^\infty \wedge W^\infty \wedge M^+]_{\mathbb{Z}_2}$$

\square

Proposition 6.13. (i) For any \mathbb{Z}_2 -space X the \mathbb{Z}_2 -equivariant Pontryagin-Thom construction defines an isomorphism

$$\Omega_m^{\mathbb{Z}_2\text{-}fr}(X) \xrightarrow{\cong} \omega_m^{\mathbb{Z}_2}(X) ; (M, f) \mapsto (1 \wedge f) \alpha_M^{\mathbb{Z}_2} .$$

In particular,

$$(M, 1) = \alpha_M^{\mathbb{Z}_2} \in \Omega_m^{\mathbb{Z}_2\text{-}fr}(M) = \omega_m^{\mathbb{Z}_2}(M) .$$

(ii) A \mathbb{Z}_2 -framed m -dimensional \mathbb{Z}_2 -manifold M is \mathbb{Z}_2 -equivariantly self- S -dual, in the sense that the composite

$$\begin{aligned} \Delta \alpha_M^{\mathbb{Z}_2} : (LV \oplus W \oplus \mathbb{R}^m)^\infty &\rightarrow LV \oplus W^\infty \wedge M^+ \\ &\rightarrow M^+ \wedge T(\epsilon_{LV \oplus W}) = M^+ \wedge (LV^\infty \wedge W^\infty \wedge M^+) \end{aligned}$$

is a \mathbb{Z}_2 -equivariant S -duality map. Cap product with the \mathbb{Z}_2 -equivariant stable homotopy fundamental class

$$[M] = (M, 1) = \alpha_M^{\mathbb{Z}_2} \in \omega_m^{\mathbb{Z}_2}(M)$$

defines Poincaré duality isomorphisms in stable homotopy

$$[M] \cap - : \omega_{\mathbb{Z}_2}^*(M) \cong \omega_{m-*}^{\mathbb{Z}_2}(M) .$$

In particular $[M] \cap - : \omega_{\mathbb{Z}_2}^0(M) \cong \omega_m^{\mathbb{Z}_2}(M)$ sends $1 \in \omega_{\mathbb{Z}_2}^0(M)$ to $[M] \in \Omega_m^{\mathbb{Z}_2}(M)$.

(iii) For any space X there is defined a direct sum system

$$\omega_m(S(\infty) \times_{\mathbb{Z}_2} (X \times X)) \begin{array}{c} \xrightarrow{\gamma} \\ \xleftarrow{\delta} \end{array} \omega_m^{\mathbb{Z}_2}(X \times X) \begin{array}{c} \xrightarrow{\rho} \\ \xleftarrow{\sigma} \end{array} \omega_m(X)$$

with

$$\begin{aligned} \rho : \omega_m^{\mathbb{Z}_2}(X \times X) &\rightarrow \omega_m(X) ; \\ (M, f : M \rightarrow X \times X) &\mapsto (M^{\mathbb{Z}_2}, f^{\mathbb{Z}_2} : M^{\mathbb{Z}_2} \rightarrow X) , \\ \gamma : \omega_m(S(\infty) \times_{\mathbb{Z}_2} (X \times X)) &\rightarrow \omega_m^{\mathbb{Z}_2}(X \times X) ; \\ (N, g : N \rightarrow S(\infty) \times_{\mathbb{Z}_2} (X \times X)) &\mapsto (\overline{N}, \overline{g}) , \quad \overline{N} = g^*(S(\infty) \times X \times X) , \\ \sigma : \omega_m(X) &\rightarrow \omega_m^{\mathbb{Z}_2}(X \times X) ; \\ (P, h : P \rightarrow X) &\mapsto (P, \Delta_X h : P \rightarrow X \times X) , \\ \delta : \omega_m^{\mathbb{Z}_2}(X \times X) &= \{S^m; X^+ \wedge X^+\}_{\mathbb{Z}_2} \rightarrow \omega_m(S(\infty) \times_{\mathbb{Z}_2} (X \times X)) ; \\ (F : V^\infty \wedge LW^\infty \wedge S^m \rightarrow V^\infty \wedge LW^\infty \wedge X^+ \wedge X^+) &\mapsto \delta(F, \sigma\rho(F)) . \end{aligned}$$

Proof. (i)+(ii) By Proposition 4.103.

(iii) Immediate from Proposition 4.41 (ii).

For any framed m -dimensional manifold (M, b) with embedding $V \times M \subset V \oplus \mathbb{R}^m$ and Pontryagin-Thom map $\alpha_M : (V \oplus \mathbb{R}^m)^\infty \rightarrow (V \times M)^\infty$ the product $M \times M$ is a $2m$ -dimensional \mathbb{Z}_2 -manifold with a \mathbb{Z}_2 -equivariant embedding

$$\begin{aligned}
(\kappa_{V \oplus \mathbb{R}^m})^{-1}(i \times i)(\kappa_V \times 1) &: (LV \oplus V) \times (M \times M) \cong V \times V \times M \times M \\
&\rightarrow (V \oplus \mathbb{R}^m) \times (V \oplus \mathbb{R}^m) \cong (LV \oplus L\mathbb{R}^m) \times (V \oplus \mathbb{R}^m)
\end{aligned}$$

and \mathbb{Z}_2 -equivariant Pontryagin-Thom map

$$\begin{aligned}
(\kappa_V \wedge 1)(\alpha_M \wedge \alpha_M)(\kappa_{V \oplus \mathbb{R}^m})^{-1} &: (LV \oplus L\mathbb{R}^m)^\infty \wedge (V \oplus \mathbb{R}^m)^\infty \\
&\rightarrow LV^\infty \wedge V^\infty \wedge M^+ \wedge M^+ .
\end{aligned}$$

The image of

$$\begin{aligned}
[M \times M, b \times b] &= (\kappa_V \wedge 1)(\alpha_M \wedge \alpha_M)(\kappa_{V \oplus \mathbb{R}^m})^{-1} \\
&\in \omega_{2m,m}(M \times M) = \{(L\mathbb{R}^m \oplus \mathbb{R}^m)^\infty, M^+ \wedge M^+\}_{\mathbb{Z}_2}
\end{aligned}$$

under

$$0_{L\mathbb{R}^m}^* : \omega_{2m,m}(M \times M) \rightarrow \omega_m^{\mathbb{Z}_2}(M \times M) = \omega_m(M) \oplus \omega_m(S(\infty) \times_{\mathbb{Z}_2}(M \times M))$$


is of the form

$$\begin{aligned}
0_{L\mathbb{R}^m}^*[M \times M, b \times b] &= ((M, b), (N, a)) \\
&\in \omega_m^{\mathbb{Z}_2}(M \times M) = \omega_m(M) \oplus \omega_m(S(\infty) \times_{\mathbb{Z}_2}(M \times M))
\end{aligned}$$

with (N, a) the framed m -dimensional manifold with a map $g : N \rightarrow S(\infty) \times_{\mathbb{Z}_2}(M \times M)$ constructed from M by Crabb [12, p.47]. Note that there is defined an exact sequence

$$\begin{aligned}
\rightarrow A_1 &= \omega_{2m,m}(S(L\mathbb{R}^m) \times M \times M) \rightarrow A_2 = \omega_{2m,m}(S(L\mathbb{R}^m \oplus L\mathbb{R}(\infty)) \times M \times M) \\
\rightarrow A_3 &= \omega_m(S(\infty) \times_{\mathbb{Z}_2}(M \times M)) \rightarrow A_4 = \omega_{2m-1,m-1}(S(L\mathbb{R}^m) \times M \times M)
\end{aligned}$$

which fits into a commutative braid of exact sequences



Example 6.14. (Crabb [12, pp. 44-48]) If (M, b) is a framed m -dimensional manifold with an open embedding $i : V \times M \subset V \oplus \mathbb{R}^m$ the Pontryagin-Thom map

$$\alpha_M : (V \oplus \mathbb{R}^m)^\infty = V^\infty \wedge S^m \rightarrow (V \times M)^\infty = V^\infty \wedge M^+$$

defines the stable homotopy fundamental class

$$[M] = \alpha_M \in \omega_m(M) = \{S^m; M^+\}.$$

The product \mathbb{Z}_2 -manifold $M \times M$ has stable \mathbb{Z}_2 -equivariant homotopy fundamental class

$$[M \times M] = \alpha_M \wedge \alpha_M \in \omega_{2m, m}^{\mathbb{Z}_2}(M \times M) = \{(L\mathbb{R}^m)^\infty \wedge S^m; M^+ \wedge M^+\}_{\mathbb{Z}_2}.$$

The quadratic construction on α_M

$$\psi(\alpha_M) \in \omega_m(S(LV) \times_{\mathbb{Z}_2} (M \times M))$$

has the following geometric interpretation, with $n = \dim(V)$. Define a $(2m + n)$ -dimensional manifold with boundary

$$(W, \partial W) = (D(LV), S(LV)) \times M \times M,$$

with \mathbb{Z}_2 -action

$$T : W \rightarrow W ; (v, x, y) \mapsto (-v, y, x)$$

(which is free on ∂W), with a stable \mathbb{Z}_2 -equivariant homotopy fundamental class

$$[W]_{\alpha_M \wedge \alpha_M} = \alpha_M \wedge \alpha_M \in \omega_m^{\mathbb{Z}_2}(W, \partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) = \omega_{2m, m}(M \times M).$$

Consider the stable \mathbb{Z}_2 -equivariant cohomotopy exact sequence associated to the trivial \mathbb{Z}_2 -bundle $\epsilon_{LV \oplus L\mathbb{R}^m}$ over W

$$\begin{aligned} \cdots \rightarrow \omega_{\mathbb{Z}_2}^{-1}(\partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) &\rightarrow \omega_{\mathbb{Z}_2}^0(W, \partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) \rightarrow \omega_{\mathbb{Z}_2}^0(W; -\epsilon_{LV \oplus L\mathbb{R}^m}) \\ &\rightarrow \omega_{\mathbb{Z}_2}^0(\partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) \rightarrow \cdots \end{aligned}$$

There are two reasons for the stable \mathbb{Z}_2 -equivariant cohomotopy Euler class

$$\gamma^{\mathbb{Z}_2}(\epsilon_{LV \oplus L\mathbb{R}^m}) \in \omega_{\mathbb{Z}_2}^0(W; -\epsilon_{LV \oplus L\mathbb{R}^m}) = \{(M \times M)^+; (LV \oplus L\mathbb{R}^m)^\infty\}_{\mathbb{Z}_2}$$

to restrict to 0 on ∂W , namely the \mathbb{Z}_2 -equivariant sections of the sphere bundle of $\epsilon_{LV \oplus L\mathbb{R}^m}$

$$s_1, s_2 : \partial W \rightarrow S(\epsilon_{LV \oplus L\mathbb{R}^m}) = S(LV \oplus L\mathbb{R}^m) \times \partial W$$

defined by

$$s_1(v, x, y) = \left(\frac{i(v, x) - i(-v, y)}{\|i(v, x) - i(-v, y)\|}, (v, x, y) \right), \quad s_2(v, x, y) = (v, (v, x, y)).$$

The \mathbb{Z}_2 -equivariant map

$$\begin{aligned} (s_1, -s_2) : \partial W &\rightarrow S(\epsilon_{LV \oplus L\mathbb{R}^m}) \times_{\partial W} S(\epsilon_{LV \oplus L\mathbb{R}^m}) \\ &= S(LV \oplus L\mathbb{R}^m) \times S(LV \oplus L\mathbb{R}^m) \times \partial W; \\ (v, x, y) &\mapsto (s_1(v, x, y), -s_2(v, x, y), (v, x, y)) \end{aligned}$$

is \mathbb{Z}_2 -equivariant homotopic to a \mathbb{Z}_2 -equivariant map

$$(t_1, -t_2) : \partial W \rightarrow S(\epsilon_{LV \oplus L\mathbb{R}^m}) \times_{\partial W} S(\epsilon_{LV \oplus L\mathbb{R}^m})$$

which is \mathbb{Z}_2 -equivariantly transverse regular at $\Delta_{S(LV \oplus L\mathbb{R}^m)} \times \partial W$. The inverse image

$$Y = (t_1, -t_2)^{-1}(\Delta_{S(LV \oplus L\mathbb{R}^m)} \times \partial W) \subset \partial W$$

is a framed m -dimensional manifold with a free \mathbb{Z}_2 -action, and with a \mathbb{Z}_2 -equivariant map $Y \rightarrow \partial W$, such that Y is nonequivariantly framed cobordant to M . The m -dimensional manifold

$$N = Y/\mathbb{Z}_2$$

is equipped with a map $g : N \rightarrow \partial W/\mathbb{Z}_2 = S(LV) \times_{\mathbb{Z}_2} (M \times M)$. The \mathbb{Z}_2 -equivariant Poincaré duality defines an isomorphism of exact sequences

$$\begin{array}{ccccc} \omega_{\mathbb{Z}_2}^{-1}(\partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) & \longrightarrow & \omega_{\mathbb{Z}_2}^0(W, \partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) & \longrightarrow & \omega_{\mathbb{Z}_2}^0(W; -\epsilon_{LV \oplus L\mathbb{R}^m}) \\ [\partial W]_{\alpha_M \wedge \alpha_M} \cap - \Big| \cong & & [W]_{\alpha_M \wedge \alpha_M} \cap - \Big| \cong & & [W]_{\alpha_M \wedge \alpha_M} \cap - \Big| \cong \\ \omega_m^{\mathbb{Z}_2}(\partial W) & \longrightarrow & \omega_m^{\mathbb{Z}_2}(W) & \longrightarrow & \omega_m^{\mathbb{Z}_2}(W, \partial W) \end{array}$$

and

$$\begin{aligned} \delta(s_1, s_2) &= \psi_V(\alpha_M) = (N, g) \\ \in \omega_{\mathbb{Z}_2}^{-1}(\partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) &= \{\Sigma \partial W^+; (LV \oplus L\mathbb{R}^m)^\infty\}_{\mathbb{Z}_2} \\ &= \omega_m^{\mathbb{Z}_2}(\partial W) = \omega_m(S(LV) \times_{\mathbb{Z}_2} (M \times M)). \end{aligned}$$

Note that for any $v \in S(LV)$ the \mathbb{Z}_2 -equivariant map

$$v \times 1 \times 1 : S(L\mathbb{R}) \times M \times M \rightarrow S(LV) \times M \times M$$

sends $\delta(s_1, s_2)$ to the element

$$(v \times 1 \times 1)^* \delta(s_1, s_2) \in \omega_{\mathbb{Z}_2}^{-1}(S(L\mathbb{R}) \times M \times M; -\epsilon_{LV \oplus L\mathbb{R}^m}) = \{M \times M; S(V \oplus \mathbb{R}^m)\}$$

represented by the map

$$M \times M \rightarrow S(V \oplus \mathbb{R}^m) = S^{m+n-1} ; (x, y) \mapsto \frac{i(v, x) - i(-v, y)}{\|i(v, x) - i(-v, y)\|} .$$

The rel ∂W \mathbb{Z}_2 -equivariant Euler classes are

$$\begin{aligned} \gamma^{\mathbb{Z}_2}(\epsilon_{LV \oplus L\mathbb{R}^m}, s_1) &= (\alpha_M \wedge \alpha_M) \Delta_{S^m} , \quad \gamma^{\mathbb{Z}_2}(\epsilon_{LV \oplus L\mathbb{R}^m}, s_2) = \Delta_M \alpha_M \\ &\in \omega_{\mathbb{Z}_2}^0(W, \partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) = \omega_m^{\mathbb{Z}_2}(W) = \omega_m^{\mathbb{Z}_2}(M \times M) \end{aligned}$$

and $\delta(s_1, s_2)$ has image

$$\begin{aligned} \gamma^{\mathbb{Z}_2}(\epsilon_{LV \oplus L\mathbb{R}^m}, s_1) - \gamma^{\mathbb{Z}_2}(\epsilon_{LV \oplus L\mathbb{R}^m}, s_2) &= (\alpha_M \wedge \alpha_M) \Delta_{S^m} - \Delta_M \alpha_M \\ &\in \text{im}(\omega_{\mathbb{Z}_2}^{-1}(\partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) \rightarrow \omega_{\mathbb{Z}_2}^0(W, \partial W; -\epsilon_{LV \oplus L\mathbb{R}^m})) \\ &= \ker(\omega_{\mathbb{Z}_2}^0(W, \partial W; -\epsilon_{LV \oplus L\mathbb{R}^m}) \rightarrow \omega_{\mathbb{Z}_2}^0(W; -\epsilon_{LV \oplus L\mathbb{R}^m})) \subset \omega_{2m, m}(M \times M) . \end{aligned}$$

Also, passing to the limit as $\dim(V) \rightarrow \infty$ note that

$$\begin{aligned} \lim_{\dim(V)} \psi_V(\alpha_M) &= (\alpha_M \wedge \alpha_M) \Delta_{S^m} - \Delta_M \alpha_M \\ &\in \text{im}(\omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) \rightarrow \omega_{2m, m}^{\mathbb{Z}_2}(M \times M)) \\ &= \ker(\rho : \omega_{2m, m}^{\mathbb{Z}_2}(M \times M) \rightarrow \omega_m(M)) \end{aligned}$$

is determined by $(\alpha_M \wedge \alpha_M) \Delta_{S^m}, \Delta_M \alpha_M \in \omega_m^{\mathbb{Z}_2}(M \times M)$, since

$$\omega_m(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) \rightarrow \omega_{2m, m}^{\mathbb{Z}_2}(M \times M)$$

is a split injection. □

6.2 Double points

Definition 6.15. (i) The *fibrewise product* of maps $f_1 : X_1 \rightarrow Y$, $f_2 : X_2 \rightarrow Y$ is the map

$$f_1 \times_Y f_2 : X_1 \times_Y X_2 \rightarrow Y ; (x_1, x_2) \mapsto f(x_1) = f(x_2)$$

with

$$X_1 \times_Y X_2 = \{(x_1, x_2) \in X_1 \times X_2 \mid f_1(x_1) = f_2(x_2) \in Y\}$$

the pullback

$$\begin{array}{ccc}
X_1 \times_Y X_2 & \longrightarrow & X_1 \\
\downarrow & \searrow f_1 \times_Y f_2 & \downarrow f_1 \\
X_2 & \xrightarrow{f_2} & Y
\end{array}$$

(ii) The *square* of a map $f : X \rightarrow Y$ is the \mathbb{Z}_2 -equivariant map

$$f \times f : X \times X \rightarrow Y \times Y ; (x_1, x_2) \mapsto (f(x_1), f(x_2))$$

with $T \in \mathbb{Z}_2$ acting on $X \times X$ by

$$T : X \times X \rightarrow X \times X ; (x_1, x_2) \mapsto (x_2, x_1)$$

and similarly for $Y \times Y$.

(iii) The *double point space* of $f : X \rightarrow Y$ is the \mathbb{Z}_2 -space

$$X \times_Y X = (f \times f)^{-1}(Y) = \{(x_1, x_2) \in X \times X \mid f(x_1) = f(x_2) \in Y\} \subset X \times X .$$

The diagonal embedding

$$\Delta_X : X \rightarrow X \times_Y X ; x \mapsto (x, x)$$

is \mathbb{Z}_2 -equivariant, with image the fixed point set

$$(X \times_Y X)^{\mathbb{Z}_2} = X = \{(x, x) \mid x \in X\} .$$

The map

$$f \times_Y f = (f \times f)| : X \times_Y X \rightarrow Y ; (x_1, x_2) \mapsto f(x_1) = f(x_2) .$$

is also \mathbb{Z}_2 -equivariant, with $(f \times_Y f)\Delta_X = f : X \rightarrow Y$.

(iv) Let $\tilde{Y} \rightarrow Y$ be a regular cover of Y with group of covering translations π , and given a map $f : X \rightarrow Y$ let

$$\tilde{X} = X \times_Y \tilde{Y} = f^* \tilde{Y} = \{(x, \tilde{y}) \in X \times \tilde{Y} \mid f(x) = [\tilde{y}] \in Y\}$$

be the pullback cover of X . The *assembly map* is the inclusion

$$A : X \times_Y X \rightarrow \tilde{X} \times_{\pi} \tilde{X} ; (x_1, x_2) \mapsto [(x_1, \tilde{y}), (x_2, \tilde{y})]$$

with $\tilde{y} \in \tilde{Y}$ any lift of $y = f(x_1) = f(x_2) \in Y$.

(v) The *ordered double point space* of $f : X \rightarrow Y$ is the free \mathbb{Z}_2 -space

$$D_2(f) = \{(x_1, x_2) \in X \times X \mid x_1 \neq x_2 \in X, f(x_1) = f(x_2) \in Y\}$$

with

$$X \times_Y X = X \sqcup D_2(f) .$$

(vi) The *unordered double point set* of $f : X \rightarrow Y$ is the quotient space

$$D_2[f] = D_2(f)/\mathbb{Z}_2 .$$

The projection $D_2(f) \rightarrow D_2[f]$ is a double cover. \square

Remark 6.16. The diagonal $Y \subseteq Y \times Y$ is a closed subspace (since Y is Hausdorff), so that $X \times_Y X = (f \times f)^{-1}(Y) \subseteq X \times X$ is also closed, and $X \subseteq X \times_Y X$ is also closed (since X is Hausdorff). We shall always assume that $X \subseteq X \times_Y X$ is also open, so that

$$X \times_Y X = X \sqcup D_2(f)$$

as a topological space, with one-point compactification

$$(X \times_Y X)^\infty = X^\infty \vee D_2(f)^\infty .$$

\square

We shall be concerned with the double points of a map $f : X \rightarrow Y$ which extends to an embedding $V \times X \hookrightarrow V \times Y$ (in the manner of a codimension 0 immersion of manifolds) :

Definition 6.17. (i) An *embedding* of a map $f : X \rightarrow Y$ is an open embedding of the type

$$e : V \times X \hookrightarrow V \times Y ; (v, x) \mapsto (g(v, x), f(x))$$

for an inner product space V , so that the diagram

$$\begin{array}{ccc} V \times X & \xhookrightarrow{e} & V \times Y \\ \text{proj.} \downarrow & & \downarrow \text{proj.} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. For each $y \in Y$ the restriction of e defines an embedding

$$e| : V \times f^{-1}(y) \rightarrow V ; (v, x) \mapsto g(v, x) .$$

The pair $e = (g, f)$ is an *embedded map*.

(ii) The *compactification Umkehr* of an embedded map $e = (g, f)$ is the compactification Umkehr given by 3.18

$$F : V^\infty \wedge Y^\infty \rightarrow V^\infty \wedge X^\infty ;$$

$$(w, y) \mapsto \begin{cases} (v, x) & \text{if } (w, y) = e(v, x) \in e(V \times X) \\ * & \text{otherwise.} \end{cases}$$

□

Let $e = (g, f) : V \times X \hookrightarrow V \times Y$ be an embedded map, with compactification Umkehr map $F : V^\infty \wedge Y^\infty \rightarrow V^\infty \wedge X^\infty$. We shall now construct a *local geometric Hopf map*

$$h_V(F)_Y = F_1 \vee \delta(F_2, F_3) : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty$$

$$\rightarrow LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty = LV^\infty \wedge V^\infty \wedge (D_2(f)^\infty \vee X^\infty)$$

with

$$F_1 : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge D_2(f)^\infty$$

the Umkehr map of a \mathbb{Z}_2 -equivariant embedding

$$e_1 : LV \times V \times D_2(f) \hookrightarrow (LV \setminus \{0\}) \times V \times Y ,$$

and

$$\delta(F_2, F_3) : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty$$

the relative difference of the Umkehr maps

$$F_2, F_3 : LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty$$

of two \mathbb{Z}_2 -equivariant embeddings $e_2, e_3 : LV \times V \times X \hookrightarrow LV \times V \times Y$ of $f : X \rightarrow Y$ such that

$$e_2|_{\{0\}} = e_3|_{\{0\}} = e : \{0\} \times V \times X \hookrightarrow \{0\} \times V \times Y .$$

The geometric Hopf invariant map is the assembly of the local geometric Hopf map

$$h_V(F) = (1 \wedge A)h_V(F)_Y :$$

$$\Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge (X \times X)^\infty$$

with $\tilde{Y} = Y$, $\pi = \{1\}$, $A : X \times_Y X \subset X \times X$.

Terminology 6.18 Given a map $f : X \rightarrow Y$ and an embedding $e = (g, f) : V \times X \hookrightarrow V \times Y$ define a \mathbb{Z}_2 -equivariant embedding of $f \times f : X \times X \rightarrow Y \times Y$

$$\begin{aligned}
e \times e &= (g \times g, f \times f) : \\
(V \times V) \times (X \times X) &\hookrightarrow (V \times V) \times (Y \times Y) ; \\
(u, v, x, y) &\mapsto (g(u, x), g(v, y), f(x), f(y))
\end{aligned}$$

with \mathbb{Z}_2 -equivariant compactification Umkehr map

$$F \wedge F : V^\infty \wedge V^\infty \wedge Y^\infty \wedge Y^\infty \rightarrow V^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty .$$

The conjugate of $e \times e$ by $\kappa_V \times 1$ is a \mathbb{Z}_2 -equivariant embedding of $f : X \rightarrow Y$

$$\begin{aligned}
e' &= (\kappa_V \times 1)^{-1}(e \times e)(\kappa_V \times 1) : LV \times V \times X \times X \hookrightarrow LV \times V \times Y \times Y ; \\
(u, v, x, y) &\mapsto (g(u + v, x) - g(-u + v, x))/2, (g(u + v, x) + g(-u + v, x))/2, f(x), f(y) .
\end{aligned}$$

with \mathbb{Z}_2 -equivariant compactification Umkehr map

$$\begin{aligned}
F' &= (\kappa_V \wedge 1)(F \wedge F)(\kappa_V \wedge 1)^{-1} : \\
LV^\infty \wedge V^\infty \wedge Y^\infty \wedge Y^\infty &\rightarrow LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty .
\end{aligned}$$

The restriction of e' defines a \mathbb{Z}_2 -equivariant open embedding

$$\begin{aligned}
e_1 &= e'| : LV \times V \times D_2(f) \hookrightarrow (LV \setminus \{0\}) \times V \times Y ; (u, v, x, y) \mapsto \\
&(g(u + v, x) - g(-u + v, x))/2, (g(u + v, x) + g(-u + v, x))/2, f(x) = f(y)
\end{aligned}$$

with a \mathbb{Z}_2 -equivariant compactification Umkehr map

$$F_1 : (LV \setminus \{0\})^\infty \wedge V^\infty \wedge Y^\infty = \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge D_2(f)^\infty .$$

The components of the \mathbb{Z}_2 -equivariant embeddings of $f : X \rightarrow Y$

$$e_2, e_3 : LV \times V \times X \hookrightarrow LV \times V \times Y$$

defined by

$$\begin{aligned}
e_2(u, v, x) &= (1 \times e)(u, v, x) = (u, g(v, x), f(x)) , \\
e_3(u, v, x) &= e'(u, v, x) \\
&= (g(u + v, x) - g(-u + v, x))/2, (g(u + v, x) + g(-u + v, x))/2, f(x)
\end{aligned}$$

have \mathbb{Z}_2 -equivariant compactification Umkehr maps

$$F_2 = 1 \wedge F, F_3 : LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty$$

with

$$\begin{aligned}
e_2| &= e_3| = e : \{0\} \times V \times X \hookrightarrow \{0\} \times V \times Y , \\
F_2| &= F_3| : \{0\}^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty .
\end{aligned}$$

□

Theorem 6.19. (*Double Point Theorem*)

For an embedding $e = (g, f) : V \times X \hookrightarrow V \times Y$ of a map $f : X \rightarrow Y$ the geometric Hopf invariant $h_V(F)$ of the Umkehr map $F : V^\infty \wedge Y^\infty \rightarrow V^\infty \wedge X^\infty$ is given up to natural \mathbb{Z}_2 -equivariant homotopy by the composite

$$\begin{aligned} h_V(F) &= (1 \wedge A)h_V(F)_Y : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \\ &\rightarrow LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty \rightarrow LV^\infty \wedge V^\infty \wedge (X \times X)^\infty \end{aligned}$$

with

$$\begin{aligned} h_V(F)_Y &= (1 \wedge i)F_1 + (1 \wedge \Delta_X)\delta(F_2, F_3) : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty &\rightarrow \\ LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty &= LV^\infty \wedge V^\infty \wedge (D_2(f)^\infty \vee X^\infty) , \\ i &= \text{inclusion} : D_2(f) \rightarrow X \times_Y X ; (x_1, x_2) \mapsto (x_1, x_2) , \\ \Delta_X &= \text{diagonal} : X \rightarrow X \times_Y X ; x \mapsto (x, x) , \\ A &= \text{assembly} = \Delta_X \sqcup i : X \times_Y X = X \sqcup D_2(f) \rightarrow X \times X . \end{aligned}$$

Proof. By definition

$$h_V(F) = \delta(p, q) : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty$$

with $p = (1 \wedge \Delta_{X^\infty})F_2$, $q = F'(1 \wedge \Delta_{Y^\infty})$ the composites in the (noncommutative) square of \mathbb{Z}_2 -equivariant maps

$$\begin{array}{ccc} LV^\infty \wedge V^\infty \wedge Y^\infty & \xrightarrow{1 \wedge \Delta_{Y^\infty}} & LV^\infty \wedge V^\infty \wedge Y^\infty \wedge Y^\infty \\ \downarrow F_2 & & \downarrow F' \\ LV^\infty \wedge V^\infty \wedge X^\infty & \xrightarrow{1 \wedge \Delta_{X^\infty}} & LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty \end{array}$$

with

$$p| = q| : \{0\}^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty .$$

Consider the commutative square of \mathbb{Z}_2 -equivariant open embeddings and inclusions

$$\begin{array}{ccc} LV \times V \times (X \times_Y X) & \xrightarrow{1 \times (f \times_Y f) = e_1 \sqcup e_3} & LV \times V \times Y \\ \downarrow 1 \times A & & \downarrow 1 \times \Delta_Y \\ LV \times V \times X \times X & \xrightarrow{e'} & LV \times V \times Y \times Y \end{array}$$

which determines a commutative square of \mathbb{Z}_2 -equivariant Umkehr maps and inclusions

$$\begin{array}{ccc}
 LV^\infty \wedge V^\infty \wedge Y^\infty & \xrightarrow{F_{13}} & LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty \\
 \downarrow 1 \wedge \Delta_{Y^\infty} & & \downarrow 1 \wedge A \\
 LV^\infty \wedge V^\infty \wedge Y^\infty \wedge Y^\infty & \xrightarrow{F'} & LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty
 \end{array}$$

with F_{13} the Umkehr map of $e_1 \sqcup e_3$. Thus

$$\begin{aligned}
 h_V(F) &= \delta(p, q) = \delta((1 \wedge \Delta_{X^\infty})F_2, F'(1 \wedge \Delta_{Y^\infty})) \\
 &= \delta((1 \wedge \Delta_{X^\infty})F_2, (1 \wedge (i \vee \Delta_{X^\infty}))F_{13}) : \\
 &\quad \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty .
 \end{aligned}$$

Proposition 2.23 (iv) gives a natural \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned}
 h_V(F) &\simeq (1 \wedge A \Delta_{X^\infty})\delta(F_2, F_3) + (1 \wedge A)\delta((1 \wedge \Delta_{X^\infty})F_3, (1 \wedge (i \vee \Delta_{X^\infty}))F_{13}) \\
 &= (1 \wedge A \Delta_{X^\infty})\delta(F_2, F_3) + \delta(p', q') : \\
 &\quad \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty
 \end{aligned}$$

with

$$\begin{aligned}
 p' &= (1 \wedge \Delta_{X^\infty})F_3, \quad q' = (1 \wedge (i \vee \Delta_{X^\infty}))F_{13} : \\
 &\quad LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty
 \end{aligned}$$

the composites in the (noncommutative) diagram of \mathbb{Z}_2 -equivariant maps

$$\begin{array}{ccc}
 LV^\infty \wedge V^\infty \wedge Y^\infty & \xrightarrow{F_{13}} & LV^\infty \wedge V^\infty \wedge (D_2(f)^\infty \vee X^\infty) \\
 \downarrow F_3 & & \parallel \\
 LV^\infty \wedge V^\infty \wedge X^\infty & \xrightarrow{1 \wedge \Delta_{X^\infty}} & LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty
 \end{array}$$

The neighbourhood

$$U = e_3(LV \times V \times Y) \cup \{*\} \subset LV^\infty \wedge V^\infty \wedge Y^\infty$$

of $0^+ \wedge V^\infty \wedge Y^\infty$ is such that for all $t \in LV^\infty$, $x \in V^\infty \wedge Y^\infty$

$$p'(t, x) = \begin{cases} q'(t, x) & \text{if } (t, x) \in U \\ * & \text{if } (t, x) \in \overline{(LV^\infty \wedge V^\infty \wedge Y^\infty) \setminus U} . \end{cases}$$

Proposition 2.23 (v) gives a natural \mathbb{Z}_2 -equivariant homotopy

$$\delta(p', q') \simeq q'' : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge (X \times_Y X)^\infty$$

with q'' defined by

$$q'' : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge (X \times_Y X)^\infty ;$$

$$(t, x) \mapsto \begin{cases} * & \text{if } (t, x) \in U \\ q'(t, x) & \text{if } (t, x) \in \overline{(LV^\infty \wedge V^\infty \wedge Y^\infty) \setminus U} . \end{cases}$$

But $q'' = (1 \wedge i)F_1$, giving the required result.

□

Similarly for the stable geometric Hopf invariant $h'_V(F) :$

Proposition 6.20. *Let $e = (g, f) : V \times X \hookrightarrow V \times Y$ be an embedded map*

$$\begin{array}{ccc} V \times X & \xrightarrow{e} & V \times Y \\ \text{proj.} \downarrow & & \downarrow \text{proj.} \\ X & \xrightarrow{f} & Y \end{array}$$

with Umkehr map $F : V^\infty \wedge Y^\infty \rightarrow V^\infty \wedge X^\infty$. The stable geometric Hopf invariant of F is given up to natural \mathbb{Z}_2 -equivariant homotopy by the sum

$$h'_V(F) = (1 \wedge A)h'_V(F)_Y : Y^\infty \rightarrow S(LV)^+ \wedge X^\infty \wedge X^\infty$$

with

$$h'_V(F)_Y = cF'_1 + (1 \wedge \Delta_{X^\infty})\delta'(F_2, F_3) : Y^\infty \rightarrow S(LV)^+ \wedge (X \times_Y X)^\infty$$

$$cF'_1 : Y^\infty \xrightarrow{F'_1} D_2(f)^\infty \xrightarrow{c} S(LV)^+ \wedge (X \times_Y X)^\infty$$

the composite of the \mathbb{Z}_2 -equivariant Umkehr map

$$F'_1 : LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge D_2(f)^\infty$$

of the \mathbb{Z}_2 -equivariant open embedding

$$e'_1 = (\kappa_V \times 1)^{-1}(e \times e)(\kappa_V \times 1) : LV \times V \times D_2(f) \hookrightarrow LV \times V \times Y ,$$

and the \mathbb{Z}_2 -equivariant map c defined by

$$c : D_2(f)^\infty \rightarrow S(LV)^+ \wedge (X \times_Y X)^\infty ; (x_1, x_2) \mapsto \left(\frac{g(0, x_1) - g(0, x_2)}{\|g(0, x_1) - g(0, x_2)\|}, x_1, x_2 \right),$$

and $\delta'(F_3, F_2) : Y^\infty \rightarrow S(LV)^+ \wedge X^\infty$ the stable relative difference of the \mathbb{Z}_2 -equivariant Umkehr maps

$$F_2, F_3 : LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty$$

of 6.18.

Proof. The \mathbb{Z}_2 -equivariant open embedding e'_1 is the composite

$$e'_1 : LV \times V \times D_2(f) \xrightarrow{e_1} (LV \setminus \{0\}) \times V \times Y \hookrightarrow LV \times V \times Y$$

with $e_1 = (\kappa_V \times 1)^{-1}(e \times e)(\kappa_V \times 1)$ as in 6.18, and the Umkehr of e'_1 is the composite

$$\begin{aligned} F'_1 &= F_1(\alpha_{LV} \wedge 1) : LV^\infty \wedge V^\infty \wedge Y^\infty \\ &\xrightarrow{\alpha_{LV} \wedge 1} (LV \setminus \{0\})^\infty \wedge V^\infty \wedge Y^\infty = \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \\ &\xrightarrow{F_1} LV^\infty \wedge V^\infty \wedge D_2(f)^\infty \end{aligned}$$

with F_1 the Umkehr map of e_1 as in 6.18. Theorem 6.19 gives a natural \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned} h_V(F) &\simeq (1 \wedge i)F_1(1 \wedge \Delta_{X^\infty}) + \delta(F_2, F_3) : \\ &\Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty \end{aligned}$$

with $i = \text{inclusion} : D_2(f) \rightarrow X \times_Y X$, so that there is defined a natural \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned} h'_V(F)_Y &= (1 \wedge h_V(F)_Y)(\Delta \alpha_{LV} \wedge 1) \\ &\simeq ((1 \wedge i)(1 \wedge F_1)(1 \wedge \Delta_{X^\infty}) + \delta(F_2, F_3))(\Delta \alpha_{LV} \wedge 1) : \\ &LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow S(LV)^+ \wedge LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty. \end{aligned}$$

Now

$$\begin{aligned}
& (1 \wedge i)(1 \wedge F_1)(\Delta \alpha_{LV} \wedge 1) : \\
& LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow S(LV)^+ \wedge LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty ; \\
& ((t, u), v, y) \mapsto \begin{cases} (u, v_1, v_2, x_1, x_2) \\ \quad \text{if } [t, u] = (g(v_1 + v_2, x_1) - g(-v_1 + v_2, x_2))/2 \in LV^\infty \setminus \{0\}, \\ \quad v = (g(v_1 + v_2, x_1) + g(-v_1 + v_2, x_2))/2 \in V, \\ \quad y = f(x_1) = f(x_2) \\ * \quad \text{otherwise} \end{cases}
\end{aligned}$$

and

$$\begin{aligned}
& (1 \wedge c)(1 \wedge F_1)(\alpha_{LV} \wedge 1) : \\
& LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow S(LV)^+ \wedge LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty ; \\
& ((t, u), v, y) \mapsto \begin{cases} (\frac{g(0, x_1) - g(0, x_2)}{\|g(0, x_1) - g(0, x_2)\|}, v_1, v_2, x_1, x_2) \\ \quad \text{if } [t, u] = (g(v_1 + v_2, x_1) - g(-v_1 + v_2, x_2))/2 \in LV^\infty \setminus \{0\}, \\ \quad v = (g(v_1 + v_2, x_1) + g(-v_1 + v_2, x_2))/2 \in V, \\ \quad y = f(x_1) = f(x_2) \\ * \quad \text{otherwise .} \end{cases}
\end{aligned}$$

The map

$$\begin{aligned}
& h : I \times LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow S(LV)^+ \wedge LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty ; \\
& (s, (t, u^+), v, y) \mapsto \begin{cases} (\frac{g(sv_1 + sv_2, x_1) - g(-sv_1 + sv_2, x_2)}{\|g(sv_1 + sv_2, x_1) - g(-sv_1 + sv_2, x_2)\|}, v_1, v_2, x_1, x_2) \\ \quad \text{if } [t, u] = (g(v_1 + v_2, x_1) - g(-v_1 + v_2, x_2))/2 \in LV^\infty \setminus \{0\} \\ \quad v = (g(v_1 + v_2, x_1) + g(-v_1 + v_2, x_2))/2 \in V, \\ \quad y = f(x_1) = f(x_2) \\ * \quad \text{otherwise} \end{cases}
\end{aligned}$$

defines a natural \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned}
h : (1 \wedge i)(1 \wedge F_1)(\Delta \alpha_{LV} \wedge 1) & \simeq (1 \wedge c)(1 \wedge F_1)(\alpha_{LV} \wedge 1) : \\
LV^\infty \wedge V^\infty \wedge Y^\infty & \rightarrow S(LV)^+ \wedge LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty .
\end{aligned}$$

The concatenation of the homotopies is a natural \mathbb{Z}_2 -equivariant homotopy

$$\begin{aligned}
h'_V(F)_Y & \simeq (1 \wedge c)(1 \wedge F_1)(\alpha_{LV} \wedge 1) = (1 \wedge c)(1 \wedge F'_1) : \\
LV^\infty \wedge V^\infty \wedge Y^\infty & \rightarrow S(LV)^+ \wedge LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty .
\end{aligned}$$

Finally, identify

$$\delta(F_2, F_3)(\Delta\alpha_{LV} \wedge 1) = \delta'(F_2, F_3) : LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty .$$

□

6.3 Positive embeddings

We shall now formulate a positivity condition on an embedded map

$$e = (g, f) : V \times X \hookrightarrow V \times Y$$

which ensures that the embedding term $(1 \wedge \Delta_{X^\infty})\delta(F_2, F_3)$ in the expression given by Theorem 6.19 for the geometric Hopf invariant $h_V(F)$ of the compactification Umkehr map $F : V^\infty \wedge Y^\infty \rightarrow V^\infty \wedge X^\infty$ is 0, so that $h_V(F)$ only depends on the double point \mathbb{Z}_2 -space $D_2(f)$ with $h_V(F) = (i \wedge 1)F_1$.

Definition 6.21. (i) A linear automorphism $a \in GL(V)$ of an inner product space V is *positive* if

$$\langle a(v), v \rangle > 0 \quad (v \neq 0 \in V) .$$

(ii) Let $GL^+(V) \subseteq GL(V)$ be the subgroup of the positive linear automorphisms.

(iii) Let

$$O^+(V) = O(V) \cap GL^+(V) \subseteq O(V)$$

be the subgroup of the positive orthogonal automorphisms.

Proposition 6.22. (i) *The topological group $GL^+(V)$ is contractible.*

(ii) *The following conditions on an orthogonal automorphism $a : V \rightarrow V$ are equivalent:*

(a) *a is positive,*

(b) *$1 + a : V \rightarrow V$ is a linear automorphism,*

(c) *-1 is not an eigenvalue of a .*

(iii) *The topological group $O^+(V)$ is contractible.*

Proof. (i) For any $a \in GL^+(V)$ and $t \in [0, 1]$ define the linear map

$$H_t(a) : V \rightarrow V ; v \mapsto tv + (1-t)a(v) .$$

For any $v \neq 0 \in V$

$$\langle H_t(a)(v), v \rangle = t\langle v, v \rangle + (1-t)\langle a(v), v \rangle > 0$$

so that $H_t(a)$ is a positive automorphism. It now follows from

$$H_0(a) = a , H_1(a) = \text{id.} : V \rightarrow V$$

that

$$H : I \times GL^+(V) \rightarrow GL^+(V) ; (t, a) \mapsto H_t(a)$$

defines a contraction of $GL^+(V)$ to $\{\text{id.}\} \subset GL^+(V)$.

(ii) Immediate from the definitions.

(iii) $O^+(V)$ is homeomorphic to the contractible space $\text{End}^-(V)$ of linear endomorphisms $b : V \rightarrow V$ such that

$$\langle b(u), v \rangle = -\langle u, b(v) \rangle > 0 \quad (u, v \in V) ,$$

with the Cayley parametrization (Weyl [70, II.10]) of positive orthogonal automorphisms defining inverse homeomorphisms

$$\begin{aligned} O^+(V) &\rightarrow \text{End}^-(V) ; a \mapsto (1+a)^{-1}(1-a) , \\ \text{End}^-(V) &\rightarrow O^+(V) ; b \mapsto (1+b)^{-1}(1-b) . \end{aligned}$$

□

Definition 6.23. An embedding $e = (g, f) : V \times X \hookrightarrow V \times Y$ of $f : X \rightarrow Y$ is *positive* if for each $x \in X$ the embedding

$$g_x : V \rightarrow V ; u \mapsto g(u, x)$$

is differentiable and the differentials $dg_x(v) : V \rightarrow V$ ($v \in V$) are positive linear automorphisms. □

Example 6.24. For any map $c : X \rightarrow O^+(V)$ the embedding

$$V \times X \hookrightarrow V \times X ; (v, x) \mapsto (c(x)(v), x)$$

is positive, with each

$$g_x : V \rightarrow V ; v \mapsto c(x)(v) \quad (x \in X)$$

a linear map with $dg_x = c(x) \in O^+(V)$.

□

The embeddings $g_x : V \rightarrow V$ ($x \in X$) in a positive embedding $e = (g, f) : V \times X \rightarrow V \times Y$ are orientation-preserving.

Theorem 6.25. *Let $e = (g, f) : V \times X \hookrightarrow V \times Y$ be a positively embedded map, with adjunction Umkehr map $F : V^\infty \wedge Y^\infty \rightarrow V^\infty \wedge X^\infty$.*

(i) *The local geometric Hopf invariant is given up to natural \mathbb{Z}_2 -equivariant homotopy by*

$$h_V(F)_Y = (1 \wedge i)F_1 : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty$$

with i = inclusion : $D_2(f) \rightarrow X \times_Y X$, and

$$F_1 : (LV \setminus \{0\})^\infty \wedge V^\infty \wedge Y^\infty = \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge D_2(f)^\infty$$

the compactification Umkehr of the \mathbb{Z}_2 -equivariant embedding

$$e_1 : LV \times V \times D_2(f) \hookrightarrow (LV \setminus \{0\}) \times V \times Y$$

of 6.19.

(ii) *The geometric Hopf invariant $h_V(F)$ is given up to natural \mathbb{Z}_2 -equivariant homotopy by*

$$h_V(F) = (1 \wedge A)h_V(F)_Y = (1 \wedge Ai)F_1 :$$

$$\Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \xrightarrow{h_V(F)_Y}$$

$$LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty \xrightarrow{1 \wedge A} LV^\infty \wedge V^\infty \wedge X^\infty \wedge X^\infty$$

with $A : X \times_Y X \subset X \times X$ the assembly.

Proof. (i) We shall use the positive property of the embedding $e = (g, f)$ to construct a \mathbb{Z}_2 -equivariant isotopy

$$E : e_3 \simeq e_2 : LV \times V \times X \rightarrow LV \times V \times Y$$

which is constant on $\{0\} \times V \times X$. This will give a \mathbb{Z}_2 -equivariant homotopy of the Umkehr maps

$$F_3 \simeq F_2 : LV^\infty \wedge V^\infty \wedge Y^\infty \rightarrow LV^\infty \wedge V^\infty \wedge X^\infty$$

which is constant on $\{0\}^+ \wedge V^\infty \wedge Y^\infty$, so that $\delta(F_3, F_2) \simeq \{*\}$ by Proposition 1.6 (i).

In the first instance, note that for any $\epsilon > 0$ and any \mathbb{Z}_2 -equivariant embedding $h : LV \times V \times X \hookrightarrow LV \times V \times Y$ there is defined a \mathbb{Z}_2 -equivariant isotopy

$$I \times LV \times V \times X \rightarrow LV \times V \times Y ; (t, u, v, x) \mapsto h\left(\frac{(1-t+t\epsilon)u}{1+t\|u\|}, v, x\right)$$

from h to the \mathbb{Z}_2 -equivariant embedding

$$\bar{h} : LV \times V \times X \hookrightarrow LV \times V \times Y ; (u, v, x) \mapsto h\left(\frac{\epsilon u}{1+\|u\|}, v, x\right)$$

which only involves the restriction

$$h| : D_\epsilon(0) \times V \times X \hookrightarrow LV \times V \times Y .$$

It therefore suffices to construct a \mathbb{Z}_2 -equivariant isotopy

$$e_3| \simeq e_2| : D_\epsilon(0) \times V \times X \hookrightarrow LV \times V \times Y$$

which is constant on $\{0\} \times V \times X$. For sufficiently small $\epsilon > 0$ and $u \in D_\epsilon(0)$ we have

$$\begin{aligned} e_2(u, v, x) &= ((g(u+v, x) - g(-u+v, x))/2, (g(u+v, x) + g(-u+v, x))/2, f(x)) \\ &\approx (u + dg_x(v)(u), g(v, x), f(x)) , \end{aligned}$$

$$e_3(u, v, x) = (u, g(v, x), f(x))$$

and

$$\begin{aligned} I \times D_\epsilon(0) \times V \times X &\hookrightarrow LV \times V \times Y ; \\ (t, u, v, x) &\mapsto (1-t)e_3(u, v, x) + te_2(u, v, x) \approx ((1-t)u + tdg_x(v)(u), g(v, x), f(x)) \end{aligned}$$

defines an isotopy $e_3| \simeq e_2|$ which is constant on $\{0\} \times V \times X$. The positive condition on e is used to ensure that the linear map

$$V \rightarrow V ; u \mapsto tu + (1-t)dg_x(v)(u)$$

is an automorphism for each $t \in [0, 1]$, exactly as in the proof of Lemma 6.22.

(ii) Immediate from (i) and 6.19.

□

Definition 6.26. Given maps $f_1 : X_1 \rightarrow Y$, $f_2 : X_2 \rightarrow Y$ write the compactification of the fibrewise product $X_1 \times_Y X_2$ of 6.15 as

$$(X_1 \times_Y X_2)^\infty = X_1^\infty \wedge_Y X_2^\infty .$$

□

Example 6.27. For compact X_1, X_2 the cartesian product of vector bundles $\xi_1 : X_1 \rightarrow BO(m_1)$, $\xi_2 : X_2 \rightarrow BO(m_2)$ is a vector bundle $\xi_1 \times \xi_2 : X_1 \times X_2 \rightarrow BO(m_1 + m_2)$ with total and Thom spaces

$$E(\xi_1 \times \xi_2) = E(\xi_1) \times E(\xi_2) , \quad T(\xi_1 \times \xi_2) = T(\xi_1) \wedge T(\xi_2) .$$

For maps $f_1 : X_1 \rightarrow Y$, $f_2 : X_2 \rightarrow Y$ the restriction of $\xi_1 \times \xi_2$ to $X_1 \times_Y X_2 \subseteq X_1 \times X_2$

$$\xi_1 \times_Y \xi_2 = (\xi_1 \times \xi_2)|_{X_1 \times_Y X_2} : X_1 \times_Y X_2 \rightarrow BO(m_1 + m_2)$$

is such that

$$E(\xi_1 \times_Y \xi_2) = E(\xi_1) \times_Y E(\xi_2) , \quad T(\xi_1 \times_Y \xi_2) = T(\xi_1) \wedge_Y T(\xi_2) .$$

□

Proposition 6.28. *Given an immersion $f : M^m \looparrowright N^n$ let $\nu_f : M \rightarrow BO(n - m)$ be the normal $(n - m)$ -plane bundle, so that f extends to a codimension 0 immersion*

$$f' : M' = E(\nu_f) \looparrowright N .$$

Assume that f has no triple points, and only transverse double points, so that the ordered double point space $M'' = D_2(f)$ is a closed $(2m - n)$ -dimensional manifold.

(i) *The immersion*

$$f \times_N f = f \sqcup f'' : M \times_N M = M \sqcup M'' \looparrowright N$$

has normal bundle $\nu_f \sqcup \nu_{f''}$, with

$$\begin{aligned} f'' : M'' \looparrowright N ; (x, y) &\mapsto f(x) = f(y) , \\ \nu_{f''} &= (\nu_f \times \nu_f)|_{M''} : M'' \rightarrow BO(2(n - m)) . \end{aligned}$$

The ordered double point space of f' is the n -dimensional manifold

$$D_2(f')^n = E(\nu_{f''}) = E((\nu_f \times_N \nu_f)|_{M''})$$

and

$$T(\nu_f) \wedge_N T(\nu_f) = T(\nu_f \sqcup \nu_{f''}) = T(\nu_f) \vee D_2(f')^\infty .$$

(ii) *For some finite-dimensional inner product space V the codimension 0 immersion*

$$1 \times f' : V \times M' \hookrightarrow V \times N ; (v, x) \mapsto (v, f'(x))$$

is regular homotopic to a positive embedding of f'

$$e = (g, f') : V \times M' \hookrightarrow V \times N ; (v, x) \mapsto (g(v, x), f'(x))$$

with Umkehr map

$$F : V^\infty \wedge N^\infty \rightarrow V^\infty \wedge (M')^\infty = V^\infty \wedge T(\nu_{f'})$$

and differentials of the form

$$\begin{aligned} de(v, x) &= \begin{pmatrix} dg_x(v) & \alpha(v, x) \\ 0 & df'(x) \end{pmatrix} : \tau_{(v, x)}(V \times E(\nu_{f'})) = V \oplus \tau_x(E(\nu_{f'})) \\ &\rightarrow \tau_{(g(v, x), f'(x))}(V \times N) = V \oplus \tau_{f'(x)}(N) \quad (g_x(v) = g(v, x)) \end{aligned}$$

with $dg_x(v) : V \rightarrow V$ positive linear automorphisms. The fibrewise geometric Hopf invariant of F is given up to natural \mathbb{Z}_2 -equivariant homotopy by

$$\begin{aligned} h_V(F)_N &= (1 \wedge i)F_1 : \Sigma S(LV)^+ \wedge V^\infty \wedge N^+ = (LV \setminus \{0\})^\infty \wedge V^\infty \wedge N^+ \\ &\rightarrow LV^\infty \wedge V^\infty \wedge T(\nu_{f''}) \rightarrow LV^\infty \wedge V^\infty \wedge T(\nu_f \times_N \nu_f) . \end{aligned}$$

with

$$i = \text{inclusion} : T(\nu_{f''}) = T((\nu_f \times_N \nu_f)|_{M''}) \rightarrow T(\nu_f \times_N \nu_f) = T(\nu_f) \wedge_N T(\nu_f) .$$

The geometric Hopf invariant of F is

$$h_V(F) = (1 \wedge Ai)F_1 : \Sigma S(LV)^+ \wedge V^\infty \wedge N^\infty \rightarrow LV^\infty \wedge V^\infty \wedge T(\nu_f) \wedge T(\nu_f)$$

with

$$A = \text{assembly} : T(\nu_f \times_N \nu_f) = T(\nu_f) \wedge_N T(\nu_f) \rightarrow T(\nu_f \times \nu_f) = T(\nu_f) \wedge T(\nu_f) .$$

Proof. (i) By construction.

(ii) The double point space $D_2(f) \subset M \times M$ is a compact manifold with free involution, and we can choose a smooth map $g_0 : M \rightarrow V$ for some V of large dimension such that

$$g_0(x) = -g_0(y) \in S(V) \quad ((x, y) \in D_2(f)) .$$

We can take g_0 to be zero outside a neighbourhood of $D_2(f) \subset M \times M$, so that there is defined an embedding

$$(g_0, f) : M \hookrightarrow V \times N ; x \mapsto (g_0(x), f(x)) .$$

Define

$$g : V \times M \rightarrow N ; (v, x) \mapsto g(v, x) = g_0(x) + v .$$

and let $D(V) = \{v \in V \mid \|v\| \leq 1\}$ be the unit ball in V . The map

$$(g, f) : V \times M \rightarrow V \times N ; (v, x) \mapsto (g(v, x), f(x))$$

restricts to an embedding of $D(V) \times M \hookrightarrow V \times N$, by the following argument: the derivative has maximal rank everywhere, and if

$$(g(v, x), f(x)) = (g(w, y), f(y)) \in V \times N$$

then for $(x, y) \in D_2(f)$ with $g(v, x) = g(w, y)$,

$$v - w = g_0(y) - g_0(x) , \quad g_0(x) = -g_0(y) , \quad \|v - w\| = 2$$

and $(v, x) = (w, y)$. Composing with the open embedding

$$j : V \rightarrow D(V) ; v \mapsto \frac{v}{1 + \|v\|}$$

gives an embedding

$$e : V \times M \hookrightarrow V \times N ; (v, x) \mapsto (jg(v, x), f(x))$$

which extends to a positive open embedding

$$e' : V \times M' \hookrightarrow V \times N ; (v, x) \mapsto (g'(v, x), f'(x)) .$$

Now apply Theorem 6.25. □

Example 6.29. (i) Let $M = N = \{*\}$, and $V = \mathbb{R}^k$ with the standard inner product

$$\langle (u_1, u_2, \dots, u_k), (v_1, v_2, \dots, v_k) \rangle = \sum_{i=1}^k u_i v_i \in \mathbb{R} .$$

For any $\lambda_1, \lambda_2, \dots, \lambda_k > 0$ and $c = (c_1, c_2, \dots, c_k) \in V$ the embedding

$$g : V \hookrightarrow V ; v = (v_1, v_2, \dots, v_k) \mapsto c + \frac{(\lambda_1 v_1, \lambda_2 v_2, \dots, \lambda_k v_k)}{1 + \|v\|}$$

is such that $e = (g, f) : V \times M \hookrightarrow V \times N$ is a positive embedding of the unique map $f : M \rightarrow N$, with

$$\begin{aligned}
dg_*(v) : V &\rightarrow V ; u = (u_1, u_2, \dots, u_k) \mapsto \frac{(\lambda_1 u_1, \lambda_2 u_2, \dots, \lambda_k u_k)}{(1 + \|v\|)^2} , \\
\langle dg_*(v)(u), u \rangle &= \frac{\sum_{i=1}^k \lambda_i u_i^2}{(1 + \|v\|)^2} > 0 \quad (u \neq 0 \in V) , \\
e(V \times M) &\subseteq D_\lambda(c) \times N
\end{aligned}$$

where

$$D_\lambda(c) = \{u = (u_1, u_2, \dots, u_k) \in V \mid |u_i - c_i| < \lambda_i, 1 \leq i \leq k\} \subset V .$$

(ii) A finite cover $f : \tilde{K} \rightarrow K$ admits positive embeddings $e = (g, f) : V \times \tilde{K} \hookrightarrow V \times K$ (cf. Example 3.8). For $k > 2 \dim(K)$ there exists an embedding $d : \tilde{K} \hookrightarrow V = \mathbb{R}^k$, and for each $x \in \tilde{K}$ it is possible to make a continuous choice of $\lambda_1(x), \lambda_2(x), \dots, \lambda_k(x) > 0$ with

$$D_{\lambda(x)}(d(x)) \cap D_{\lambda(y)}(d(y)) = \emptyset \quad (x \neq y \in \tilde{K}, f(x) = f(y) \in K) .$$

The map

$$g : V \times \tilde{K} \rightarrow V ; (v, x) \mapsto d(x) + \frac{(\lambda_1(x)v_1, \lambda_2(x)v_2, \dots, \lambda_k(x)v_k)}{1 + \|v\|}$$

determines a positive embedding $e = (g, f)$ with

$$e(V \times \{x\}) \subseteq D_{\lambda(x)}(d(x)) \times \{f(x)\} \quad (x \in \tilde{K}) .$$

The (stable) homotopy class of the compactification Umkehr map $F : V^\infty \wedge K^+ \rightarrow V^\infty \wedge \tilde{K}^+$ of e depends only on the homotopy class of the finite covering f . \square

Example 6.30. (i) Given a map $c : X \rightarrow O(V)$ define an embedding of $f = 1 : X \rightarrow X$

$$e = (g, 1) : V \times X \hookrightarrow V \times X ; (v, x) \mapsto (c(x)(v), x)$$

with adjunction Umkehr map

$$F : V^\infty \wedge X^+ \rightarrow V^\infty \wedge X^+ ; (v, x) \mapsto (c(x)^{-1}(v), x) .$$

The \mathbb{Z}_2 -equivariant embeddings $e_2, e_3 : LV \times V \times X \hookrightarrow LV \times V \times X$ in Theorem 6.19 are given by

$$\begin{aligned}
e_2(u, v, x) &= (u, g(v, x), f(x)) = (u, c(x)(v), x) , \\
e_3(u, v, x) &= ((g(u + v, x) - g(-u + v, x))/2, (g(u + v, x) + g(-u + v, x))/2, f(x)) \\
&= (c(x)(u), c(x)(v), x) ,
\end{aligned}$$

with compactification Umkehr maps

$$F_2 = 1 \wedge F, \quad F_3 = F \wedge F : LV^\infty \wedge V^\infty \wedge X^+ \rightarrow LV^\infty \wedge V^\infty \wedge X^+ \wedge X^+$$

such that

$$h_V(F) \simeq (1 \wedge \Delta_{X^+}) \delta(F_2, F_3) : \Sigma S(LV)^+ \wedge V^\infty \wedge X^+ \rightarrow LV^\infty \wedge V^\infty \wedge X^+ \wedge X^+$$

(cf. Proposition 5.75). Let $p : X^+ \rightarrow S^0$ be the projection, so that

$$pF = F_{c^{-1}} : LV^\infty \wedge X^+ \rightarrow LV^\infty$$

is the adjoint map of c^{-1} , and

$$-\theta(c) = \delta(1, F_{c^{-1}}) \in \{\Sigma S(LV)^+ \wedge X^+; LV^\infty\}_{\mathbb{Z}_2} = \omega_{\mathbb{Z}_2}^{-1}(S(LV) \times X; -\epsilon_{LV})$$

with

$$h_V(F_{c^{-1}}) = (F_{c^{-1}} \wedge -\theta(c)) \wedge \Delta_{X^+} \in \{\Sigma S(LV)^+ \wedge X^+; LV^\infty\}_{\mathbb{Z}_2}$$

as in Proposition 5.75.

(ii) As in Definition 6.21 let $O^+(V) \subset O(V)$ be the subgroup of the positive orthogonal automorphisms $a : V \rightarrow V$. If the map $c : X \rightarrow O(V)$ in (i) is such that $c(X) \subseteq O^+(V)$ then $h_V(F) \simeq *$, by Theorem 6.25. (Alternatively, note that the isotopy $e \simeq \text{id}$ defined by

$$V \times X \times I \rightarrow V \times X ; (v, x, t) \mapsto (tv + (1 - t)c(x)(v), x) ,$$

induces a homotopy $F \simeq \text{id}$.) □

Example 6.31. Given a manifold N , a space X , an inner product space V and a map $\rho : V^\infty \wedge N^+ \rightarrow V^\infty \wedge X^+$ it is possible to make ρ transverse at $\{0\} \times X \subset V^\infty \wedge X^+$, so that there exist a codimension 0 immersion $f : M = \rho^{-1}(\{0\} \times X) \looparrowright N$ with an embedding $e = (g, f) : V \times M \hookrightarrow V \times N$ and a map $h = \rho| : M \rightarrow X$ so that up to homotopy

$$\rho = (1 \wedge h)F : V^\infty \wedge N^+ \xrightarrow{F} V^\infty \wedge M^+ \xrightarrow{1 \wedge h} V^\infty \wedge X^+$$

with F the adjunction Umkehr of e . (In general, it is not possible to choose e to be a positive embedding – see (iii) below for an explicit example). The framing of $0 \times f$ determined by g differs from the canonical framing in (i) by a map $c : M \rightarrow O(V)$ such that

$$g(v, x) = g_0(c(x)(v), x) \in V \quad (v \in V, x \in M) .$$

The embeddings of f

$$e_0 = (g_0, f) , \quad e = (g, f) : V \times M \hookrightarrow V \times N$$

are such that $e = e_0(c, 1)$ with

$$(c, 1) : V \times M \rightarrow V \times M ; (v, x) \mapsto (c(x)(v), x) ,$$

so that the adjunction Umkehr maps $F_0, F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge M^+$ differ by the adjunction Umkehr of $(c, 1)^{-1}$

$$C : V^\infty \wedge M^+ \rightarrow V^\infty \wedge M^+ ; (v, x) \mapsto (c(x)(v), x)$$

(cf. Proposition 5.75) with

$$F : V^\infty \wedge N^+ \xrightarrow{F_0} V^\infty \wedge M^+ \xrightarrow{C} V^\infty \wedge M^+ .$$

By Theorem 6.25 and the composition formula of Proposition 5.33 (v) the geometric Hopf invariant of F is given by

$$\begin{aligned} h_V(F) &= h_V(CF_0) = h_V(C)(1 \wedge F_0) + (\kappa_V^{-1} \wedge 1)(C \wedge C)(\kappa_V \wedge 1)h_V(F_0) : \\ &\quad \Sigma S(LV)^+ \wedge V^\infty \wedge N^+ \rightarrow LV^\infty \wedge V^\infty \wedge M^+ \wedge M^+ . \end{aligned}$$

(iii) If $N = X = \{\text{pt.}\}$, $V = \mathbb{R}$ then

$$\rho = -1 : V^\infty \wedge N^+ = S^1 \rightarrow V^\infty \wedge X^+ = S^1 ; z \mapsto z^{-1}$$

is a map of degree -1 with $M = \rho^{-1}(\{0\} \times X) = \{*\}$, such that the corresponding embedding

$$e = (g, f) : V \times M = \mathbb{R} \hookrightarrow V \times N = \mathbb{R} ; v \mapsto -v$$

of the codimension 0 immersion $f : M \looparrowright N$ is not positive. The canonical (positive) embedding of f

$$e_0 = (g_0, f) : V \times M = \mathbb{R} \hookrightarrow V \times N = \mathbb{R} ; v \mapsto v$$

has adjunction Umkehr

$$F_0 = 1 : V^\infty \wedge N^+ = S^1 \rightarrow V^\infty \wedge M^+ = S^1 ,$$

and $e = (g, f)$ has adjunction Umkehr

$$F = F_0 C = \rho = -1 : V^\infty \wedge N^+ = S^1 \rightarrow V^\infty \wedge M^+ = S^1 ,$$

with $C : S^1 \rightarrow S^1$ the adjoint of $c = -1 : M = \{*\} \rightarrow O(\mathbb{R}) = \{\pm 1\}$. The geometric Hopf invariant of C is $h_{\mathbb{R}}(C) = 1 \in \mathbb{Z}$, by Example 5.36. \square

Proposition 6.32. *Let $e = (g, f) : \mathbb{R} \times M \hookrightarrow \mathbb{R} \times N$ be a positively embedded map, with adjunction Umkehr map*

$$F : \mathbb{R}^\infty \wedge N^+ = \Sigma N^+ \rightarrow \mathbb{R}^\infty \wedge M^+ = \Sigma M^+ .$$

(i) *The transposition \mathbb{Z}_2 -action on the set*

$$E = \mathbb{R} \times \mathbb{R} \times D_2(f)$$

is free, with a decomposition

$$E = E_+ \sqcup E_-$$

defined by

$$\begin{aligned} E_+ &= \{(v_1, v_2, x_1, x_2) \in \mathbb{R} \times \mathbb{R} \times M \times M \mid \\ &\quad g(v_1, x_1) > g(v_2, x_2) \in \mathbb{R}, x_1 \neq x_2 \in M, f(x_1) = f(x_2) \in N\} , \\ E_- &= \{(v_1, v_2, x_1, x_2) \in \mathbb{R} \times \mathbb{R} \times M \times M \mid \\ &\quad g(v_1, x_1) < g(v_2, x_2) \in \mathbb{R}, x_1 \neq x_2 \in M, f(x_1) = f(x_2) \in N\} . \end{aligned}$$

(ii) *The geometric Hopf invariant of F is given up to a natural homotopy by*

$$\begin{aligned} h_{\mathbb{R}}(F)_+ &= i_+ F_+ : (L\mathbb{R} \setminus \{0\})_+^\infty \wedge \mathbb{R}^\infty \wedge N^+ = \Sigma^2 N^+ \\ &\rightarrow L\mathbb{R}^\infty \wedge \mathbb{R}^\infty \wedge M^+ \wedge M^+ = \Sigma^2(M^+ \wedge M^+) \end{aligned}$$

(using Terminology 5.39) with

$$F_+ : (L\mathbb{R} \setminus \{0\})_+^\infty \wedge \mathbb{R}^\infty \wedge N^+ \rightarrow E_+^\infty$$

the adjunction Umkehr of the embedding

$$\begin{aligned} e_+ : E_+ &\hookrightarrow (L\mathbb{R} \setminus \{0\})_+ \times \mathbb{R} \times N ; (v_1, v_2, x_1, x_2) \mapsto (w_1, w_2, y) \\ (w_1 &= (g(v_1, x_1) - g(v_2, x_2))/2 \in (L\mathbb{R} \setminus \{0\})_+ , \\ w_2 &= (g(v_1, x_1) + g(v_2, x_2))/2 \in \mathbb{R} , y = f(x_1) = f(x_2) \in N) \end{aligned}$$

and

$$i_+ = \text{inclusion} : E_+ \rightarrow \mathbb{R} \times \mathbb{R} \times M \times M .$$

Proof. This is the special case $V = \mathbb{R}$ of Theorem 6.25.

□

Example 6.33. Let $f : M = \{1, 2\} \rightarrow N = \{*\}$ be the unique map, with ordered double point set

$$D_2(f) = (M \times M) \setminus \Delta(M) = \{(1, 2), (2, 1)\}.$$

Define a positive embedding $(e : \mathbb{R} \times M \hookrightarrow \mathbb{R} \times N, f : M \rightarrow N)$ of f by

$$e : \mathbb{R} \times \{1, 2\} \hookrightarrow \mathbb{R} ; \quad \begin{cases} (v, 1) \mapsto e^v \\ (v, 2) \mapsto -e^{-v} \end{cases}.$$

The compactification Umkehr of e is a sum map

$$F = \nabla = h_1 + h_2 : (\mathbb{R} \times N)^\infty = S^1 \rightarrow (\mathbb{R} \times M)^\infty = S^1 \vee S^1$$

with

$$h_i : S^1 \rightarrow S^1 \vee S^1 ; t \mapsto t_i \quad (i = 1, 2),$$

and by Proposition 6.32 the geometric Hopf invariant of F is determined up to \mathbb{Z}_2 -equivariant homotopy by the map

$$\begin{aligned} |h_{\mathbb{R}}(F)| &= h_1 \wedge h_2 : \\ \Sigma^2 N^+ &= S^2 = S^1 \wedge S^1 \rightarrow \Sigma^2(M^+ \wedge M^+) = (S^1 \vee S^1) \wedge (S^1 \vee S^1). \end{aligned}$$

□

6.4 Finite covers

In this section K is a finite CW complex. Every finite covering $f : \tilde{K} \rightarrow K$ admits a positive embedding $e = (g, f) : V \times \tilde{K} \hookrightarrow V \times K$ (6.29 (ii)), with a compactification Umkehr map $F : V^\infty \wedge K^+ \rightarrow V^\infty \wedge \tilde{K}^+$ (2.2).

For a finite covering $f : \tilde{K} \rightarrow K$ of degree d define a commutative square of finite coverings

$$\begin{array}{ccc} \tilde{K}' & \xrightarrow{\tilde{f}'} & \tilde{K} \\ \tilde{f} \downarrow & & \downarrow f \\ K' & \xrightarrow{f'} & K \end{array}$$

with

$$\begin{aligned}\tilde{K}' &= D_2(f) = \{(x_1, x_2) \in \tilde{K} \times \tilde{K} \mid x_1 \neq x_2 \in \tilde{K}, f(x_1) = f(x_2) \in K\}, \\ K' &= D_2[f] = \tilde{K}' / \{(x_1, x_2) \sim (x_2, x_1)\}\end{aligned}$$

so that the projections

$$\begin{aligned}\tilde{f} &: \tilde{K}' \rightarrow K' ; (x_1, x_2) \mapsto [x_1, x_2], \\ f' &: K' \rightarrow K ; [x_1, x_2] \mapsto f(x_1) = f(x_2), \\ \tilde{f}' &: \tilde{K}' \rightarrow \tilde{K} ; (x_1, x_2) \mapsto x_1\end{aligned}$$

are finite coverings of degree 2, $d(d-1)/2$, $d-1$ respectively. The composite

$$f\tilde{f}' = f'f : \tilde{K}' \rightarrow K ; (x_1, x_2) \mapsto f(x_1) = f(x_2)$$

is a finite covering of degree $d(d-1)$.

Proposition 6.34. *The geometric Hopf invariant of the compactification Umkehr map $F : V^\infty \wedge K^+ \rightarrow V^\infty \wedge \tilde{K}^+$ of a positively embedded finite covering $e = (g, f) : V \times \tilde{K} \hookrightarrow V \times K$ is given up to natural \mathbb{Z}_2 -equivariant homotopy by*

$$\begin{aligned}h_V(F) &= (1 \wedge i)F_1 : \Sigma S(LV)^+ \wedge V^\infty \wedge K^+ \\ &\rightarrow LV^\infty \wedge V^\infty \wedge (\tilde{K}')^+ \rightarrow LV^\infty \wedge V^\infty \wedge \tilde{K}^+ \wedge \tilde{K}^+\end{aligned}$$

with

$$F_1 : \Sigma S(LV)^+ \wedge V^\infty \wedge K^+ = (LV \setminus \{0\})^+ \wedge V^\infty \wedge K^+ \rightarrow LV^\infty \wedge V^\infty \wedge (\tilde{K}')^+$$

the Umkehr map of the \mathbb{Z}_2 -equivariant embedding

$$\begin{aligned}e_1 : LV \times V \times \tilde{K}' &\hookrightarrow (LV \setminus \{0\}) \times V \times K ; (v_1, v_2, x_1, x_2) \mapsto \\ &((g(v_1 + v_2, x_1) - g(-v_1 + v_2, x_2))/2, (g(v_1 + v_2, x_1) + g(-v_1 + v_2, x_2))/2), y) \\ &\quad (y = f(x_1) = f(x_2))\end{aligned}$$

and

$$i = \text{inclusion} : \tilde{K}' = D_2(f) \rightarrow \tilde{K} \times \tilde{K}.$$

Proof. A direct application of Theorem 6.25.

□

Example 6.35. For any $d \geq 1$ consider the finite covering of degree d

$$f : \tilde{K} = \{1, 2, \dots, d\} \rightarrow K = \{*\}$$

with ordered double point set

$$\tilde{K}' = D_2(f) = \{(i, j) \mid 1 \leq i, j \leq d, i \neq j\}$$

consisting of $d(d-1)$ points. For any positive embedding $e = (g, f) : \mathbb{R} \times \tilde{K} \hookrightarrow \mathbb{R} \times K = \mathbb{R}$ the Umkehr is given up to homotopy by the d -fold sum map

$$F = h_1 + h_2 + \cdots + h_d : \mathbb{R}^+ = S^1 = I/(0=1) \rightarrow \mathbb{R}^+ \wedge \tilde{K}^+ = \bigvee_d S^1 ;$$

$$t \mapsto (dt - i + 1)_i \text{ if } (i-1)/d \leq t \leq i/d$$

where

$$h_i : S^1 \rightarrow \bigvee_d S^1 ; t \mapsto t_i \text{ (} 1 \leq i \leq d \text{)} .$$

The geometric Hopf invariant of F (6.32) is given up to homotopy by

$$h_{\mathbb{R}}(F) = \sum_{1 \leq i < j \leq d} h_i \wedge h_j : S^1 \wedge S^1 \rightarrow \left(\bigvee_d S^1 \right) \wedge \left(\bigvee_d S^1 \right)$$

The geometric Hopf invariant of the composite

$$d : S^1 \xrightarrow{F} \bigvee_d S^1 \xrightarrow{1 \vee 1 \vee \cdots \vee 1} S^1$$

is thus

$$\begin{aligned} h_{\mathbb{R}}(d) &= ((1 \vee 1 \vee \cdots \vee 1) \wedge (1 \vee 1 \vee \cdots \vee 1)) h_{\mathbb{R}}(F) \\ &= d(d-1)/2 : S^2 \rightarrow S^2 . \end{aligned}$$

□

A double cover $f : \tilde{K} \rightarrow K$ of a finite CW complex K is classified by a map $c : K \rightarrow P(V)$ to the projective space $P(V) = S(LV)/\mathbb{Z}_2$ of a inner product space V , for sufficiently large $\dim(V)$.

Proposition 6.36. *For a finite cover $f : \tilde{K} \rightarrow K$ of degree d the quadratic construction on the compactification Umkehr map $F : V^\infty \wedge K^+ \rightarrow V^\infty \wedge \tilde{K}^+$ of a positive embedding $e = (g, f) : V \times \tilde{K} \hookrightarrow V \times K$ is given by*

$$\psi_V(F) : K^+ \xrightarrow{F'} K'^+ \xrightarrow{c} S(LV)^+ \wedge_{\mathbb{Z}_2} (\tilde{K} \times \tilde{K})^+$$

with F' the Umkehr map for the finite cover of degree $d(d-1)/2$

$$\begin{aligned} f' : K' &= \{(x, y) \in \tilde{K} \times \tilde{K} \mid x \neq y \in \tilde{K}, f(x) = f(y) \in K\} / \mathbb{Z}_2 \rightarrow K ; \\ [x, y] &\mapsto f(x) = f(y) \end{aligned}$$

and

$$c : K' \rightarrow S(LV) \times_{\mathbb{Z}_2} (\tilde{K} \times \tilde{K}) ; [x, y] \mapsto \left[\frac{g(0, x) - g(0, y)}{\|g(0, x) - g(0, y)\|}, x, y \right] .$$

For $d = 2$ F' is a homeomorphism

$$F' : K^+ \rightarrow (K')^+ ; f(x) \mapsto [x, Tx]$$

with $T : \tilde{K} \rightarrow \tilde{K}$ the covering translation, and

$$K \rightarrow P(V) ; f(x) \mapsto \frac{g(0, x) - g(0, Tx)}{\|g(0, x) - g(0, Tx)\|}$$

is a classifying map for the double cover $f : \tilde{K} \rightarrow K$. □

Proposition 6.37. Let $c : K \rightarrow P(V)$ be a map, classifying the double cover of K

$$\tilde{K} = c^*S(LV) = \{(v, x) \in S(LV) \times K \mid [v] = c(x) \in P(V)\}$$

with covering projection

$$f : \tilde{K} \rightarrow K ; (v, x) \mapsto x$$

and covering translation

$$T : \tilde{K} \rightarrow \tilde{K} ; (v, x) \mapsto (-v, x) .$$

(i) The map

$$g : V \times \tilde{K} \rightarrow V ; (u, (v, x)) \mapsto v + \frac{u}{1 + \|u\|}$$

determines a positive embedding

$$e = (g, f) : V \times \tilde{K} \hookrightarrow V \times K ; (u, (v, x)) \mapsto (v + \frac{u}{1 + \|u\|}, x)$$

with Umkehr map

$$F : V^\infty \wedge K^+ \rightarrow V^\infty \wedge \tilde{K}^+ ;$$

$$(u, x) \mapsto \begin{cases} (\frac{u-v}{1-\|u-v\|}, (v, x)) & \text{if } (v, x) \in \tilde{K}, \|u-v\| < 1 \\ (\frac{u+v}{1-\|u+v\|}, (-v, x)) & \text{if } (v, x) \in \tilde{K}, \|u+v\| < 1 \\ * & \text{otherwise} . \end{cases}$$

(ii) The restriction of e is a \mathbb{Z}_2 -equivariant embedding

$$e' = e| : LV \times \tilde{K} \hookrightarrow (LV \setminus \{0\}) \times K$$

with \mathbb{Z}_2 -equivariant Umkehr map

$$F' = F| : (LV \setminus \{0\})^+ \wedge K^+ = \Sigma S(LV)^+ \wedge K^+ \rightarrow LV^\infty \wedge \tilde{K}^+ .$$

(iii) For any subspace $K_0 \subseteq K$ let

$$f_0 = f| : \tilde{K}_0 = f^{-1}(K_0) \rightarrow K_0$$

be the pullback double cover of K_0 . The maps F, F' in (i), (ii) induce maps

$$\begin{aligned} F &: V^\infty \wedge K/K_0 \rightarrow V^\infty \wedge \tilde{K}/\tilde{K}_0 , \\ F' &: \Sigma S(LV)^+ \wedge K/K_0 \rightarrow LV^\infty \wedge \tilde{K}/\tilde{K}_0 \end{aligned}$$

with F' \mathbb{Z}_2 -equivariant, such that the geometric Hopf invariant of F is given up to natural \mathbb{Z}_2 -homotopy by

$$\begin{aligned} h_V(F) : \Sigma S(LV)^+ \wedge V^\infty \wedge K/K_0 &\xrightarrow{1 \wedge F'} LV^\infty \wedge V^\infty \wedge \tilde{K}/\tilde{K}_0 \\ &\xrightarrow{1 \wedge (1 \wedge T)\Delta_{\tilde{K}/\tilde{K}_0}} LV^\infty \wedge V^\infty \wedge \tilde{K}/\tilde{K}_0 \wedge \tilde{K}/\tilde{K}_0 . \end{aligned}$$

Proof. (i)+(ii) By construction.

(iii) By Proposition 6.34 $h_V(F)$ is given up to natural \mathbb{Z}_2 -equivariant homotopy to

$$\begin{aligned} h_V(F) : \Sigma S(LV)^+ \wedge V^\infty \wedge K/K_0 &\xrightarrow{F_1} LV^\infty \wedge V^\infty \wedge D_2(f)/D_2(f_0) \\ &\xrightarrow{1 \wedge i} LV^\infty \wedge V^\infty \wedge \tilde{K}/\tilde{K}_0 \wedge \tilde{K}/\tilde{K}_0 \end{aligned}$$

with F_1 the Umkehr map of the \mathbb{Z}_2 -equivariant open embedding

$$\begin{aligned} e_1 : LV \times V \times D_2(f) &\hookrightarrow (LV \setminus \{0\}) \times V \times K ; (u, v, x_1, x_2) \mapsto \\ &((g(u+v, x_1) - g(-u+v, x_2))/2, (g(u+v, x_1) + g(-u+v, x_2))/2, y) \\ &(y = f(x_1) = f(x_2) \in K) \end{aligned}$$

and $i : D_2(f) \rightarrow \tilde{K} \times \tilde{K}$ the inclusion. Use the \mathbb{Z}_2 -equivariant homeomorphism

$$\tilde{K} \rightarrow D_2(f) ; (v, x) \mapsto ((v, x), (-v, x))$$

to identify

$$i : D_2(f) = \tilde{K} \rightarrow \tilde{K} \times \tilde{K} ; (v, x) \mapsto ((v, x), (-v, x))$$

and to express e_1 as

$$\begin{aligned} e_1 : LV \times V \times \tilde{K} \hookrightarrow (LV \setminus \{0\}) \times V \times K ; (u_1, u_2, v, x) \mapsto \\ ((g(u_1 + u_2, v, x) - g(-u_1 + u_2, -v, x))/2, \\ (g(u_1 + u_2, v, x) + g(-u_1 + u_2, -v, x))/2, f(x)) . \end{aligned}$$

The \mathbb{Z}_2 -equivariant embeddings $e_1, 1 \times e' : LV \times V \times \tilde{K} \hookrightarrow (LV \setminus \{0\}) \times V \times K$ are such that

$$e_1| = (1 \times e')| : LV \times \{0\} \times \tilde{K} \hookrightarrow (LV \setminus \{0\}) \times V \times K ; (u, 0, v, x) \mapsto (g(u, v, x), 0, f(x)) ,$$

so that by the \mathbb{Z}_2 -equivariant tubular neighbourhood theorem (applied in the special case $K = \{*\}$, $\tilde{K} = S^0$) they are related by a natural \mathbb{Z}_2 -equivariant open isotopy $e_1 \simeq (1 \times e')$ inducing a natural \mathbb{Z}_2 -equivariant homotopy

$$F_1 \simeq 1 \wedge F' : \Sigma S(LV)^+ \wedge V^\infty \wedge K/K_0 \rightarrow LV^\infty \wedge V^\infty \wedge \tilde{K}/\tilde{K}_0 \wedge \tilde{K}/\tilde{K}_0 .$$

□

Remark 6.38. (Segal [57, 58]) For any unpointed space X let $C(K; X)$ denote the set of isomorphism classes of pairs $(f : \tilde{K} \rightarrow K, g : \tilde{K} \rightarrow X)$ with f a finite cover and $g \in [\tilde{K}, X]$. The set $C(K; X)$ is an abelian semigroup with respect to disjoint union of the spaces \tilde{K} , which is a contravariant homotopy-functor of K . The Umkehr construction defines a natural transformation of semi-group valued functors $C(; X) \rightarrow \{ ; X\}$ by

$$C(K; X) \rightarrow \{K^+; X^+\} ; (f, g) \mapsto gF$$

which is a ‘group completion’, i.e. is universal among transformations $T : C(; X) \rightarrow F$ where F is a representable abelian-group-valued homotopy-functor and T is a transformation of semigroup-valued functors (Barratt, Priddy and Quillen). As above, given a finite cover $f : \tilde{K} \rightarrow K$ of degree d and a map $g : \tilde{K} \rightarrow X$ there are defined a degree $d(d-1)/2$ cover $f' : K' \rightarrow K$ and a degree 2 cover $\tilde{f}' : \tilde{K}' \rightarrow \tilde{K}$, with

$$\tilde{K}' = \{(x_1, x_2) \in \tilde{K} \times \tilde{K} \mid x_1 \neq x_2 \in \tilde{K}, f(x_1) = f(x_2) \in K\} .$$

Let $c : K' \rightarrow P(\mathbb{R}(\infty))$ be a classifying map for \tilde{f}' , with \mathbb{Z}_2 -equivariant lift $\tilde{c} : \tilde{K}' \rightarrow S(\infty)$, and let $g' : K' \rightarrow S(\infty) \times_{\mathbb{Z}_2} (X \times X)$ be the quotient of the \mathbb{Z}_2 -equivariant map

$$g' : \tilde{K}' \rightarrow S(\infty) \times (X \times X) ; (x_1, x_2) \mapsto (\tilde{c}(x_1, x_2), g(x_1), g(x_2)) .$$

The transformation of semigroup-valued functors

$$\theta^2 : C(K; X) \rightarrow C(K; S(\infty) \times_{\mathbb{Z}_2} (X \times X)) ; (f, g) \mapsto (f', g')$$

determines a transformation of abelian-group-valued functors

$$\theta^2 : \{K^+; X^+\} \rightarrow \{K^+; (S(\infty) \times_{\mathbb{Z}_2} (X \times X))^+\} .$$

In Example 6.58 below the corresponding stable map

$$\theta^2 : X^+ \rightarrow (S(\infty) \times_{\mathbb{Z}_2} (X \times X))^+$$

will be expressed in terms of the geometric Hopf invariant of the evaluation stable map $\Omega^\infty \Sigma^\infty X^+ \rightarrow X^+$.

□

6.5 Function spaces

We shall now use the quadratic construction $\psi_{\mathbb{R}^k}(F)$ and the double point theorem for finite covers to split off the quadratic part of the function spaces $\Omega^k \Sigma^k X$ for $k = 1, 2, \dots, \infty$, with X a pointed (but not necessarily connected) space.

The geometric Hopf invariant defines a map of function spaces

$$h_V : \text{map}(V^\infty \wedge X, V^\infty \wedge Y) \rightarrow \text{map}^{\mathbb{Z}_2}(\Sigma S(LV)^+ \wedge V^\infty \wedge X, LV^\infty \wedge V^\infty \wedge Y \wedge Y)$$

for any pointed spaces X, Y and inner product space V .

Definition 6.39. The pointed space

$$Q_V(X) = \text{map}(V^\infty, V^\infty \wedge X)$$

is the function space of maps $\omega : V^\infty \rightarrow V^\infty \wedge X$.

□

Proposition 6.40. (i) A map $f : Y \rightarrow Q_V(X)$ is essentially the same as a pointed map $F : V^\infty \wedge Y \rightarrow V^\infty \wedge X$, via the adjoint map construction

$$F = \text{adj}(f) : V^\infty \wedge Y \rightarrow V^\infty \wedge X ; (v, y) \mapsto f(y)(v) .$$

The function

$$[Y, Q_V(X)] \rightarrow [V^\infty \wedge Y, V^\infty \wedge X] ; f \mapsto F$$

is a bijection.

(ii) The inclusion

$$i : X \rightarrow Q_V(X) ; x \mapsto (v \mapsto (v, x))$$

has adjoint the identity map

$$\text{adj}(i) = 1 : V^\infty \wedge X \rightarrow V^\infty \wedge X .$$

(iii) The identity map $1_V : Q_V(X) \rightarrow Q_V(X)$ has adjoint the evaluation map

$$e_V = \text{adj}(1_V) : V^\infty \wedge Q_V(X) \rightarrow V^\infty \wedge X ; (v, \omega) \mapsto \omega(v) ,$$

such that for f, F as in (i)

$$F : V^\infty \wedge Y \xrightarrow{1 \wedge f} V^\infty \wedge Q_V(X) \xrightarrow{e_V} V^\infty \wedge X ,$$

and

$$\psi_V(F) : Y \xrightarrow{f} Q_V(X) \xrightarrow{\psi_V(e_V)} S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) .$$

(iv) For $V \neq \{0\}$ $Q_V(X)$ is an H -space via

$$Q_V(X) \times Q_V(X) \rightarrow Q_V(X) ; (f, g) \mapsto f + g = (f \vee g) \nabla$$

with $\nabla : V^\infty \rightarrow V^\infty \vee V^\infty$ any sum map. For $\dim(V) \geq 2$ $Q_V(X)$ is a homotopy commutative H -space.

(v) For any pointed space Y the quadratic construction defines a function

$$\psi_V : [Y, Q_V(X)] = [V^\infty \wedge Y, V^\infty \wedge X] \rightarrow \{Y; S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X)\} ; F \mapsto \psi_V(F)$$

which is induced by a stable map

$$\psi_V : Q_V(X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) .$$

□

Example 6.41. Let $V = \mathbb{R}^k$, with $k \geq 1$.

(i) If X is connected then so is $Q_V(X)$, with

$$\pi_n(Q_V(X)) = \pi_{n+k}(\Sigma^k X) .$$

(ii) For $X = S^0$

$$Q_V(S^0) = \text{map}(V^\infty, V^\infty) = \Omega^k S^k ,$$

with a bijection

$$\pi_0(Q_V(S^0)) \rightarrow \mathbb{Z} ; (\omega : V^\infty \rightarrow V^\infty) \mapsto \text{degree}(\omega) .$$

The quadratic construction is given by

$$\psi_V : \pi_0(Q_V(S^0)) = \mathbb{Z} \rightarrow \pi_0^S(P(V)^+) = \mathbb{Z} ; d \mapsto \frac{d(d-1)}{2} .$$

Proposition 5.75 gives a homotopy commutative diagram of stable maps

$$\begin{array}{ccc} O(V) & \xrightarrow{\theta \wedge J} & P(V)^+ \wedge Q_V(S^0) \\ J \downarrow & & \downarrow 1 \wedge e_V \\ Q_V(S^0) & \xrightarrow{\psi_V} & P(V)^+ \wedge S^0 = P(V)^+ \end{array}$$

given on the path-components by

$$\begin{array}{ccc} \pi_0(O(V)) = \{1, -1\} & \xrightarrow{1} & \pi_0(P(V)^+ \wedge Q_V(S^0)) = \{1, -1\} \\ \updownarrow & & \updownarrow \\ \pi_0(Q_V(S^0)) = \mathbb{Z} & \xrightarrow{\psi_V} & \pi_0^S(P(V)^+) = \mathbb{Z} . \end{array}$$

By Proposition 5.76 (ii) $\theta R \simeq \text{id}$., so there is defined a homotopy commutative diagram of stable maps

$$\begin{array}{ccc} P(V)^+ & \xrightarrow{1 \wedge JR} & P(V)^+ \wedge Q_V(S^0) \\ JR \downarrow & & \downarrow 1 \wedge e_V \\ Q_V(S^0) & \xrightarrow{\psi_V} & P(V)^+ \wedge S^0 = P(V)^+ \end{array}$$

□

Example 6.42. (i) Let K be a finite CW complex with a map $c : K \rightarrow P(V)$, classifying a double cover $f : \tilde{K} = c^*S(LV) \rightarrow K$, with Umkehr map $F : V^\infty \wedge K^+ \rightarrow V^\infty \wedge \tilde{K}^+$. The composite stable map

$$K^+ \xrightarrow{\text{adj}(F)} Q_V(\tilde{K}^+) \xrightarrow{\psi_V} S(LV)^+ \wedge_{\mathbb{Z}_2} (\tilde{K}^+ \wedge \tilde{K}^+)$$

is represented by

$$K \rightarrow S(LV) \times_{\mathbb{Z}_2} (\tilde{K} \times \tilde{K}) ; f(x) \mapsto [\tilde{c}(x), x, Tx]$$

with $\tilde{c} : \tilde{K} \rightarrow S(LV)$ a \mathbb{Z}_2 -equivariant lift of c and $T : \tilde{K} \rightarrow \tilde{K}$ the covering translation (by Proposition 5.49 (vii)).

(ii) In particular, (i) applies to the canonical double cover $f : \tilde{K} = S(LV) \rightarrow K = P(V)$ classified by $c = 1 : K \rightarrow P(V)$, with Umkehr map

$$F : V^\infty \wedge P(V)^+ \rightarrow V^\infty \wedge S(LV)^+ ;$$

$$(u, [v]) \mapsto \begin{cases} \left(\frac{u-v}{1-\|u-v\|}, v \right) & \text{if } v \in S(LV), \|u-v\| < 1 \\ \left(\frac{u+v}{1-\|u+v\|}, -v \right) & \text{if } v \in S(LV), \|u+v\| < 1 \\ * & \text{otherwise .} \end{cases}$$

Let $p : S(LV)^+ \rightarrow S^0$ be the projection. It follows from (i) that the map

$$D_V = (1 \wedge p)F : V^\infty \wedge P(V)^+ \rightarrow V^\infty ;$$

$$(u, [v]) \mapsto \begin{cases} \frac{u-v}{1-\|u-v\|} & \text{if } v \in S(LV), \|u-v\| < 1 \\ \frac{u+v}{1-\|u+v\|} & \text{if } v \in S(LV), \|u+v\| < 1 \\ * & \text{otherwise} \end{cases}$$

is such that the composite stable map

$$P(V)^+ \xrightarrow{\text{adj}(D_V)} Q_V(S^0) \xrightarrow{\psi_V} P(V)^+$$

is the identity. □

Given maps $f_1 : A \rightarrow B_1$, $f_2 : A \rightarrow B_2$ define the *homotopy pushout* to be the double mapping cone

$$\mathcal{C}(f_1, f_2) = (A \times I \sqcup B_1 \sqcup B_2) / \{(a, 0) \sim f_1(a), (a, 1) \sim f_2(a) \mid a \in A\}$$

which fits into a square

$$\begin{array}{ccc} A & \xrightarrow{f_1} & B_1 \\ f_2 \downarrow & & \downarrow i_1 \\ B_2 & \xrightarrow{i_2} & \mathcal{C}(f_1, f_2) \end{array}$$

with a canonical homotopy

$$j : i_1 f_1 \simeq i_2 f_2 : A \rightarrow \mathcal{C}(f_1, f_2) .$$

The homotopy pushout has the universal property that for any maps $g_1 : B_1 \rightarrow C$, $g_2 : B_2 \rightarrow C$ and homotopy $h : g_1 f_1 \simeq g_2 f_2 : A \rightarrow C$ there is a unique map $(g_1, g_2, h) : \mathcal{C}(f_1, f_2) \rightarrow C$ such that

$$g_1 = (g_1, g_2, h)i_1 , \quad g_2 = (g_1, g_2, h)i_2 , \quad h = (g_1, g_2, h)j .$$

Definition 6.43. (i) Given a pointed space X and an inner product space V let

$$J_{2,V}(X) = \mathcal{C}(f_1, f_2)$$

be the pointed space defined by the homotopy pushout of the maps

$$\begin{aligned} f_1 &: S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times \{*\} \cup \{*\} \times X) \rightarrow X ; \quad (v, x, *) = (-v, *, x) \mapsto x , \\ f_2 &= \text{inclusion} : S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times \{*\} \cup \{*\} \times X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) \end{aligned}$$

with base point $* \in X$, so that there are defined a homotopy commutative diagram

$$\begin{array}{ccc} S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times \{*\} \cup \{*\} \times X) & \xrightarrow{f_1} & X \\ f_2 \downarrow & & \downarrow i_1 \\ S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) & \xrightarrow{i_2} & J_{2,V}(X) \end{array}$$

and a cofibration sequence

$$X \rightarrow J_{2,V}(X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) .$$

(It is assumed that the base point $* \in X$ is nondegenerate).

(ii) The *Dyer-Lashof map* is

$$d_V : J_{2,V}(X) \rightarrow Q_V(X) ; \quad \begin{cases} x \mapsto ((v, x) \mapsto (v, x)) \\ [v, x, y] \mapsto (i(x) \vee i(y)) \nabla_v \end{cases}$$

with $\nabla_v : V^\infty \rightarrow V^\infty \vee V^\infty$ the sum map defined by the Umkehr map of the open embedding

$$V \times \{v, -v\} \hookrightarrow V ; \quad (u, \pm v) \mapsto \pm v + \frac{u}{1 + \|u\|}$$

(as in 2.14 (ii)), so that

$$d_V[v, x, y] : V^\infty \rightarrow V^\infty \wedge X ; u \mapsto \begin{cases} (\frac{u-v}{1-\|u-v\|}, x) & \text{if } \|u-v\| < 1 \\ (\frac{u+v}{1-\|u+v\|}, y) & \text{if } \|u+v\| < 1 \\ * & \text{otherwise .} \end{cases}$$

Write the adjoint map as

$$D_V = \text{adj}(d_V) : V^\infty \wedge J_{2,V}(X) \rightarrow V^\infty \wedge X .$$

(iii) For an unpointed space Y let

$$\begin{aligned} d_V^+ &: (S(LV) \times_{\mathbb{Z}_2} (Y \times Y))^+ \hookrightarrow J_{2,V}(Y^+) \xrightarrow{d_V} Q_V(Y^+) , \\ D_V^+ &= \text{adj}(d_V^+) : V^\infty \wedge (S(LV) \times_{\mathbb{Z}_2} (Y \times Y))^+ \rightarrow V^\infty \wedge Y^+ . \end{aligned}$$

□

Remark 6.44. (i) The map

$$d_{\mathbb{R}^k} : J_{2,\mathbb{R}^k}(X) \rightarrow Q_{\mathbb{R}^k}(X) = \Omega^k \Sigma^k X \quad (1 \leq k \leq \infty)$$

is essentially the same as the James map J of 5.59 (for $k = 1$) and Dyer-Lashof [18] (for $k \geq 1$).

(ii) For a connected pointed space X $Q_{\mathbb{R}^k}(X)$ is connected and the inclusion $i : X \hookrightarrow Q_{\mathbb{R}^k}(X)$ induces the k -fold suspension map in the homotopy groups

$$E^k : \pi_n(X) \rightarrow \pi_n(Q_{\mathbb{R}^k}(X)) = \pi_{n+k}(\Sigma^k X) ; f \mapsto \Sigma^k f .$$

$J_{2,\mathbb{R}^k}(X)$ is the second stage of the combinatorial model for $\Omega^k \Sigma^k X$ constructed by James [28] for $k = 1$ and by some combination of Boardman-Vogt, Milgram, May, Segal, Snaith for $2 \leq k \leq \infty$, where

$$\begin{aligned} Q_{\mathbb{R}^\infty}(X) &= \varinjlim_k Q_{\mathbb{R}^k} X = \Omega^\infty \Sigma^\infty X , \\ \pi_n(Q_{\mathbb{R}^\infty} X) &= \{S^n; X\} = \pi_n^S(X) \quad (n \geq 0) , \\ \dot{H}_*(Q_{\mathbb{R}^k}(X)) &= \bigoplus_{n=1}^{\infty} \dot{H}_*(C(k, n)^+ \wedge_{\Sigma_n} (\bigwedge_n X)) , \\ \dot{H}_*(QX) &= \bigoplus_{n=1}^{\infty} \dot{H}_*((E\Sigma_n)^+ \wedge_{\Sigma_n} (\bigwedge_n X)) \end{aligned}$$

with $C(k, n)$ the Σ_n -equivariant configuration space of embeddings $\{1, 2, \dots, n\} \hookrightarrow \mathbb{R}^k$, and $E\Sigma_n = \varinjlim_k C(k, n)$ a contractible space with a free Σ_n -action. In particular, for $k = 1$

$$C(1, n) \simeq_{\Sigma_n} \Sigma_n , \quad C(1, n)^+ \wedge_{\Sigma_n} (\bigwedge_n X) \simeq \bigwedge_n X ,$$

and for $n = 2$

$$C(k, 2) \simeq_{\Sigma_2} S^{k-1}.$$

The quadratic construction on the evaluation map

$$e_{\mathbb{R}^k} : \Sigma^k(\Omega^k \Sigma^k X) \rightarrow \Sigma^k X ; (s, \omega) \mapsto \omega(s)$$

induces the projection

$$\psi_{\mathbb{R}^k}(e_{\mathbb{R}^k}) : \dot{H}_*(\Omega^k \Sigma^k X) \rightarrow \dot{H}_*((S^{k-1})^+ \wedge_{\mathbb{Z}_2} (X \wedge X)).$$

The quadratic construction on a map $F : \Sigma^k Y \rightarrow \Sigma^k X$ with adjoint

$$\text{adj}(F) : Y \rightarrow \Omega^k \Sigma^k X ; y \mapsto (s \mapsto F(s, y))$$

is given by the composite

$$\psi_{\mathbb{R}^k}(F) : \dot{H}_*(Y) \xrightarrow{\text{adj}(F)_*} \dot{H}_*(\Omega^k \Sigma^k X) \xrightarrow{\psi_{\mathbb{R}^k}(e_{\mathbb{R}^k})} \dot{H}_*((S^{k-1})^+ \wedge_{\mathbb{Z}_2} (X \wedge X)).$$

(iii) If X is an $(m-1)$ -connected pointed space the map $d_{\mathbb{R}} : J_{2,\mathbb{R}}(X) \rightarrow \Omega \Sigma X$ is $(3m-2)$ -connected, and for $n \leq 3m-2$ the homotopy exact sequence

$$\begin{aligned} \dots \longrightarrow \pi_n(X) \longrightarrow \pi_n(J_{2,\mathbb{R}}(X)) \longrightarrow \pi_n(J_{2,\mathbb{R}}(X), X) \\ \longrightarrow \pi_{n-1}(X) \longrightarrow \pi_{n-1}(J_{2,\mathbb{R}}(X)) \longrightarrow \dots \end{aligned}$$

becomes the *EHP* exact sequence

$$\begin{aligned} \dots \longrightarrow \pi_n(X) \xrightarrow{E} \pi_{n+1}(\Sigma X) \xrightarrow{H} \pi_n(X \wedge X) \\ \xrightarrow{P} \pi_{n-1}(X) \xrightarrow{E} \pi_n(\Sigma X) \longrightarrow \dots \end{aligned}$$

with H the Hopf invariant map (Whitehead [71], Milgram [42]). The homotopy class of a map $f : S^{n-1} \rightarrow X$ together with a null-homotopy $g : \Sigma f \simeq * : S^n \rightarrow \Sigma X$ is an element $(f, g) \in \pi_n(X \wedge X)$. The Hurewicz image $h(f, g) \in \dot{H}_n(X \wedge X)$ has the following description in terms of the quadratic construction. Let $Y = X \cup_f D^n$, and use g to define a stable map

$$F : \Sigma S^n = S^{n+1} \rightarrow \Sigma X \vee S^{n+1} \simeq \Sigma X \cup_{\Sigma f} D^{n+1} \simeq \Sigma Y.$$

The quadratic construction $\psi_{\mathbb{R}}(F) : \dot{H}_n(S^n) \rightarrow \dot{H}_n(Y \wedge Y)$ sends $[S^n] \in \dot{H}_n(S^n)$ to

$$\psi_{\mathbb{R}}(F)[S^n] = h(f, g) \in \dot{H}_n(Y \wedge Y) = \dot{H}_n(X \wedge X).$$

In particular, if $f = * : S^{n-1} \rightarrow X$ then a null-homotopy $g : \Sigma f \simeq *$ is just a map $g : \Sigma(S^n) = S^{n+1} \rightarrow \Sigma X$, and $F = g : S^{n+1} \rightarrow \Sigma X$, with the Hurewicz

image of $H(g) \in \pi_n(X \wedge X)$ given by $\psi_{\mathbb{R}}(g)[S^m] \in \dot{H}_n(X \wedge X)$. For $X = S^m$, $n = 2m$

$$\begin{aligned} H &= \text{Hopf invariant} : \pi_{2m+1}(S^{m+1}) \rightarrow \pi_{2m}(S^m \wedge S^m) = \mathbb{Z} ; \\ (g : S^{2m+1} = \Sigma(S^{2m}) \rightarrow S^{m+1} = \Sigma(S^m)) &\mapsto h_{\mathbb{R}}(g) , \\ P : \mathbb{Z} \rightarrow \pi_{2m-1}(S^m) ; 1 &\mapsto J(\tau_{S^m}) = [\iota, \iota] \end{aligned}$$

with $\tau_{S^m} \in \pi_m(BSO(m)) = \pi_{m-1}(SO(m))$.

(iv) For a connected unpointed space Y there is defined a bijection

$$\pi_0(Q_{\mathbb{R}^k}(Y^+)) = \pi_k(\Sigma^k Y^+) \rightarrow \mathbb{Z} ; (\omega : S^k \rightarrow \Sigma^k Y^+) \mapsto \text{degree}(\omega) .$$

The Dyer-Lashof map

$$d_{\mathbb{R}^k}^+ : J_{2, \mathbb{R}^k}^+(Y) = Y^+ \sqcup S^{k-1} \times_{\mathbb{Z}_2} (Y \times Y) \hookrightarrow J_{2, \mathbb{R}^k}(Y^+) \xrightarrow{d_{\mathbb{R}^k}} Q_{\mathbb{R}^k}(Y^+)$$

sends Y to the degree 1 component of $Q_{\mathbb{R}^k}(Y^+)$ and $S^{k-1} \times_{\mathbb{Z}_2} (Y \times Y)$ to the degree 2 component. $J_{2, \mathbb{R}^k}^+(Y)$ is the second stage of the combinatorial model for $Q_{\mathbb{R}^k}(Y^+)$ provided by the work of Barratt-Eccles, Priddy, Quillen etc., which expresses $H_*(Q_{\mathbb{R}^k}(Y^+))$ for $1 \leq k < \infty$ (resp. $k = \infty$) as the ‘group completion’ of

$$\bigoplus_{n=1}^{\infty} \dot{H}_*(C(k, n) \times_{\Sigma_n} (\prod_n Y)) \text{ (resp. } \bigoplus_{n=1}^{\infty} \dot{H}_*(E\Sigma_n \times_{\Sigma_n} (\prod_n Y))) .$$

□

Proposition 6.45. *The geometric Hopf invariant of the Dyer-Lashof stable map $D_V : V^\infty \wedge J_{2, V}(X) \rightarrow V^\infty \wedge X$*

$$h_V(D_V) : \Sigma S(LV)^+ \wedge V^\infty \wedge J_{2, V}(X) \rightarrow LV^\infty \wedge V^\infty \wedge X \wedge X$$

is such that up to natural \mathbb{Z}_2 -equivariant homotopy

$$h_V(D_V)(1 \wedge i_1) = * : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge X \wedge X ,$$

$$h_V(D_V)(1 \wedge i_2) = 1 \wedge E_V :$$

$$\Sigma S(LV)^+ \wedge V^\infty \wedge S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) \rightarrow LV^\infty \wedge V^\infty \wedge X \wedge X$$

with $i_1 : X \rightarrow J_{2, V}(X)$, $i_2 : S(LV)^+ \times_{\mathbb{Z}_2} (X \times X) \rightarrow J_{2, V}(X)$ as in 6.43, and

$$E_V : \Sigma S(LV)^+ \wedge S(LV)^+ \times_{\mathbb{Z}_2} (X \times X) \rightarrow LV^\infty \wedge X \wedge X ;$$

$$(u, [v, x, y]) \mapsto \begin{cases} (\frac{u-v}{1-\|u-v\|}, x, y) & \text{if } \|u-v\| < 1 \\ (\frac{u+v}{1-\|u+v\|}, y, x) & \text{if } \|u+v\| < 1 \\ * & \text{otherwise} , \end{cases}$$

using the canonical homeomorphism $\Sigma S(LV)^+ \cong (LV \setminus \{0\})^\infty$ to regard the elements $u \neq * \in \Sigma S(LV)^+$ as elements $u \neq 0 \in LV$.

Proof. The double cover

$$f : \tilde{K} = S(LV) \times X \times X \rightarrow K = S(LV) \times_{\mathbb{Z}_2} (X \times X) ; (v, x, y) \mapsto [v, x, y]$$

is classified by the projection $c : K \rightarrow S(LV)/\mathbb{Z}_2 = P(V)$. As in Proposition 6.37 define a positive embedding of f

$$e = (g, f) : V \times \tilde{K} \hookrightarrow V \times K ; (u, (v, x, y)) \mapsto (v + \frac{u}{1+\|u\|}, [v, x, y]) .$$

The Umkehr map of f determined by e

$$F : V^\infty \wedge K/K_0 \rightarrow V^\infty \wedge \tilde{K}/\tilde{K}_0 ;$$

$$(u, [v, x, y]) \mapsto \begin{cases} (\frac{u-v}{1-\|u-v\|}, v, x, y) & \text{if } \|u-v\| < 1 \\ (\frac{u+v}{1-\|u+v\|}, -v, y, x) & \text{if } \|u+v\| < 1 \\ * & \text{otherwise} \end{cases}$$

is such that there is defined a homotopy commutative diagram of stable maps

$$\begin{array}{ccc} S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times \{*\} \cup \{*\} \times X) & \longrightarrow & X \\ \downarrow & & \downarrow i_1 \\ S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) & \xrightarrow{i_2} & J_{2,V}(X) \\ \downarrow F & & \searrow 1 \\ S(LV)^+ \wedge (X \times X) & \xrightarrow{p} & X \\ & & \nearrow D_V \end{array}$$

with

$$p : S(LV)^+ \wedge (X \times X) \rightarrow X ; (v, x, y) \mapsto x .$$

Now apply Proposition 6.37 with

$$K_0 = P(V) = S(LV) \times_{\mathbb{Z}_2} (*, *) \subset K , \tilde{K}_0 = S(LV) \times (*, *) \subset \tilde{K}$$

such that

$$K/K_0 = S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) , \quad \tilde{K}/\tilde{K}_0 = S(LV)^+ \wedge (X \times X) .$$

The geometric Hopf invariant $h_V(F)$ fits into a \mathbb{Z}_2 -equivariant homotopy commutative diagram of stable \mathbb{Z}_2 -equivariant homotopy commutative maps

$$\begin{array}{ccc} \Sigma S(LV)^+ \wedge K/K_0 & \xrightarrow{1 \wedge i_2} & \Sigma S(LV)^+ \wedge J_{2,V}(X) \\ h_V(F) \downarrow & & \downarrow h_V(D_V) \\ LV^\infty \wedge \tilde{K}/\tilde{K}_0 \wedge \tilde{K}/\tilde{K}_0 & \xrightarrow{1 \wedge p \wedge p} & LV^\infty \wedge X \wedge X \end{array}$$

with

$$\begin{aligned} h_V(F) : \Sigma S(LV)^+ \wedge V^\infty \wedge K/K_0 &\xrightarrow{1 \wedge F'} LV^\infty \wedge V^\infty \wedge \tilde{K}/\tilde{K}_0 \\ &\xrightarrow{1 \wedge (1 \wedge T)\Delta_{\tilde{K}/\tilde{K}_0}} LV^\infty \wedge V^\infty \wedge \tilde{K}/\tilde{K}_0 \wedge \tilde{K}/\tilde{K}_0 \end{aligned}$$

where

$$F' = F| : (LV \setminus \{0\})^\infty \wedge K/K_0 = \Sigma S(LV)^+ \wedge K/K_0 \rightarrow V^\infty \wedge \tilde{K}/\tilde{K}_0$$

is the \mathbb{Z}_2 -equivariant Umkehr map of the \mathbb{Z}_2 -equivariant embedding

$$e' = e| : LV \times \tilde{K} \hookrightarrow (LV \setminus \{0\}) \times K .$$

Finally, note that it follows from

$$\begin{aligned} D_V i_1 &= 1 : V^\infty \wedge X \rightarrow V^\infty \wedge X , \\ D_V i_2 &= (1 \wedge p)F : V^\infty \wedge K/K_0 \rightarrow V^\infty \wedge X \end{aligned}$$

that

$$\begin{aligned} h_V(D_V)(1 \wedge i_1) &= h_V(1) = * : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge X &\rightarrow LV^\infty \wedge V^\infty \wedge X \wedge X , \\ h_V(D_V)(1 \wedge i_2) &= (1 \wedge p \wedge p)h_V(F) = 1 \wedge E_V : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) &\rightarrow LV^\infty \wedge V^\infty \wedge X \wedge X . \end{aligned}$$

□

Proposition 6.46. (i) *For a pointed space X the quadratic constructions on the maps*

$$\begin{aligned} D_V &: V^\infty \wedge J_{2,V}(X) \rightarrow V^\infty \wedge X, \\ e_V &: V^\infty \wedge Q_V(X) \rightarrow V^\infty \wedge X \end{aligned}$$

are stable maps

$$\begin{aligned} \psi_V(D_V) &: J_{2,V}(X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X), \\ \psi_V(e_V) &: Q_V(X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) \end{aligned}$$

such that up to stable homotopy

$$\psi_V(D_V) = \psi_V(e_V)d_V = \text{projection} : J_{2,V}(X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X).$$

(ii) For an unpointed space Y the quadratic constructions on the maps

$$\begin{aligned} D_V^+ &: V^\infty \wedge (S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \times Y)^+) \rightarrow V^\infty \wedge Y^+, \\ e_V &: V^\infty \wedge Q_V(Y^+) \rightarrow V^\infty \wedge Y^+ \end{aligned}$$

are such that

$$\psi_V(D_V^+) = \psi_V(e_V)d_V^+ = 1 : (S(LV) \times_{\mathbb{Z}_2} (Y \times Y))^+ \rightarrow (S(LV) \times_{\mathbb{Z}_2} (Y \times Y))^+.$$

Proof. (i) By the proof of Proposition 6.45 there is defined a commutative diagram of stable maps

$$\begin{array}{ccccc} S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times \{*\} \cup \{*\} \times X) & \xrightarrow{\quad} & & & X \\ \downarrow & & \nearrow i_1 & & \parallel \\ S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) & \xrightarrow{i_2} & J_{2,V}(X) & \xrightarrow{d_V} & Q_V(X) \\ \downarrow F & & \searrow & \nearrow i & \downarrow e_V \\ S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) & \xrightarrow{p} & & & X \\ & & \nearrow D_V & & \parallel \\ & & & & 1 \end{array}$$

with

$$F : S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) \rightarrow S(LV)^+ \wedge (X \times X)$$

the Umkehr for the double cover

$$f : S(LV) \times (X \times X) \rightarrow S(LV) \times_{\mathbb{Z}_2} (X \times X)$$

and

$$p : S(LV)^+ \wedge (X \times X) \rightarrow X ; (v, x, y) \mapsto x.$$

Now

$$\psi_V(D_V)i_1 = \psi_V(1) = * : X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X),$$

and by the finite cover formula of Proposition 5.49 (vii) applied to F

$$\begin{aligned} \psi_V(D_V)i_2 &= (p \wedge p)\psi_V(F) = \text{projection} : \\ &S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) , \end{aligned}$$

so that by composition formula of 5.49 (vi) applied to $D_V = e_V d_V$

$$\psi_V(D_V) = \psi_V(e_V)d_V = \text{projection} : J_{2,V}(X) \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) .$$

(ii) Immediate from (i), with

$$\begin{aligned} X &= Y^+ = Y \sqcup \{*\} , \\ X \wedge X &= (Y \times Y)^+ \subset X \times X = (Y \times Y)^+ \sqcup (Y \times \{*\}) \sqcup \{*\} \times Y , \\ S(LV)^+ \wedge_{\mathbb{Z}_2} (X \wedge X) &= (S(LV) \times_{\mathbb{Z}_2} (Y \times Y))^+ \subset S(LV)^+ \wedge_{\mathbb{Z}_2} (X \times X) . \end{aligned}$$

□

Example 6.47. As in Proposition 6.37 let K be a space with a map $c : K \rightarrow P(V)$, classifying the double cover

$$\tilde{K} = c^*S(LV) = \{(v, x) \in S(LV) \times K \mid [v] = c(x) \in P(V)\}$$

of K . The Umkehr map $F : V^\infty \wedge K^+ \rightarrow V^\infty \wedge \tilde{K}^+$ is such that there is defined a commutative square

$$\begin{array}{ccc} K^+ & \xrightarrow{c} & P(V)^+ \\ \text{adj}(F) \downarrow & & \downarrow D_V^+ \\ Q_V(\tilde{K}^+) & \xrightarrow{p} & Q_V(S^0) \end{array}$$

and up to stable homotopy

$$c : K^+ \xrightarrow{p \text{ adj}(F)} Q_V(S^0) \xrightarrow{\psi_V} P(V)^+ .$$

□

Remark 6.48. The Dyer-Lashof map $d_V^+ : P(V)^+ \rightarrow O(V)$ is related to the reflection map $R : P(V)^+ \rightarrow O(V)$ by the identity (up to stable homotopy)

$$d_V^+ = 1 - JR : P(V)^+ \rightarrow Q_V(S^0) .$$

It is enough to verify the special case $V = \mathbb{R}$ directly, the general case following by naturality, working as in the proof of $\theta \circ R = 1$ in Proposition 5.76 (ii) with the commutative diagrams

$$\begin{array}{ccc}
P(\mathbb{R})^+ & \xrightarrow{d_{\mathbb{R}}^+ = 1 - JR} & Q_{\mathbb{R}}(S^0) \\
v \downarrow & & \downarrow v \\
P(V)^+ & \xrightarrow{d_V^+ = 1 - JR} & Q_V(S^0)
\end{array}$$

defined for $v \in P(V)$.

□

6.6 Embeddings and immersions

We shall now consider the application of the geometric Hopf invariant of §5.3 to an embedding-immersion pair

$$(e \times f : M \hookrightarrow V \times N, f : M \looparrowright N)$$

of manifolds M, N , with an Umkehr map $F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f)$. The basic result (6.54) applies the Double Point Theorem 6.19 to identify the quadratic construction

$$\psi_V(F) : N^+ \looparrowright S(LV)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))$$

with the Pontryagin-Thom map of the immersion $D_2(f)^{2m-n} \looparrowright N$ of the double point manifold.

Definition 6.49. An (m, n, j) -dimensional embedding-immersion pair

$$(e \times f : M \hookrightarrow V \times N, f : M \looparrowright N)$$

consists of an embedding $e \times f$ and an immersion f , with M an m -dimensional manifold, N an n -dimensional manifold, and V a j -dimensional inner product space. □

It will always be assumed that f only has transverse self-intersections, so that the double point sets $D_2(f)$, $D_2[f]$ (6.15) are $(2m - n)$ -dimensional manifolds.

The immersion $f : M \looparrowright N$ has a normal bundle $\nu_f : M \rightarrow BO(n - m)$, and f extends to a codimension 0 immersion $E(f) : E(\nu_f) \looparrowright N$ of the total space of ν_f . The embedding $e \times f : M \hookrightarrow V \times N$ is regular homotopic to the composite $M \looparrowright N \hookrightarrow V \times N$ of f and the embedding

$$N \hookrightarrow V \times N ; x \mapsto (0, x)$$

which has trivial normal bundle $\epsilon^j : N \rightarrow BO(j)$. It follows that the normal bundle of $e \times f$ is

$$\nu_{e \times f} = \nu_f \oplus \epsilon^j : M \rightarrow BO(n - m + j) .$$

The product immersion $1 \times E(f) : V \times E(\nu_f) \looparrowright V \times N$ is regular homotopic to an embedding $E(e \times f) : V \times E(\nu_f) \hookrightarrow V \times N$ of the immersion $E(f) : E(\nu_f) \looparrowright N$.

Definition 6.50. The *adjunction Umkehr map* (or *Pontryagin-Thom map*) of an (m, n, j) -dimensional embedding-immersion pair $(e \times f, f)$ is the adjunction Umkehr map of $E(e \times f)$

$$F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f) ; (a, x) \mapsto \begin{cases} (b, y) & \text{if } (a, x) = E(e \times f)(b, y) \\ * & \text{otherwise .} \end{cases}$$

□

Example 6.51. A finite cover $f : M \rightarrow N$ of an n -dimensional manifold is an immersion with normal bundle $\nu_f : M \rightarrow BO(0) = \{*\}$. An embedding $e : V \times M \hookrightarrow V \times N$ of f in the sense of 6.17 with $\dim(V) = j$ determines an (n, n, j) -dimensional embedding-immersion pair $(e_0 \times f, f)$ with

$$e_0 : M \rightarrow V ; x \mapsto v \text{ if } e(0, x) = (v, f(x)) .$$

The Umkehr map given by 6.50 is the same as the Umkehr map given by 6.17

$$F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f) = V^\infty \wedge M^+ .$$

□

The construction of 6.50 will also be applied to an embedding $f : M \hookrightarrow V \times N$ when M is a disjoint union of manifolds of different dimensions.

Remark 6.52. By the embedding theorem of Whitney [72] for any immersion $f : M^m \looparrowright N^n$ and $j > 2m - n$ there exists a map $e : M \rightarrow V = \mathbb{R}^j$ such that $e(x) \neq e(y)$ whenever $f(x) = f(y)$ and $x \neq y$, i.e. such that

$$e \times f : M \hookrightarrow V \times N ; x \mapsto (e(x), f(x))$$

is an embedding, and $(e \times f, f)$ is an (m, n, j) -dimensional embedding-immersion pair. □

Definition 6.53. The *square* of an (m, n, j) -dimensional embedding-immersion pair $(e \times f : M \hookrightarrow V \times N, f : M \looparrowright N)$ is the \mathbb{Z}_2 -equivariant $(2m, 2n, 2j)$ -dimensional embedding-immersion pair

$$(g : M \times M \hookrightarrow (LV \oplus V) \times N \times N, f \times f : M \times M \looparrowright N \times N)$$

with

$$\begin{aligned} g &= (e \times f) \times (e \times f) : M \times M \hookrightarrow (LV \oplus V) \times (N \times N) ; \\ (x, y) &\mapsto (\tfrac{1}{2}(e(x) - e(y)), \tfrac{1}{2}(e(x) + e(y)), f(x), f(y)) . \end{aligned}$$

□

The embedding g has normal bundle

$$\nu_g = (\nu_f \oplus \epsilon^j) \times (\nu_f \oplus \epsilon^j) : M \times M \rightarrow BO(2(n - m + j))$$

and \mathbb{Z}_2 -equivariant Umkehr map

$$F \wedge F : LV^\infty \wedge V^\infty \wedge N^+ \wedge N^+ \rightarrow T(\nu_g) = LV^\infty \wedge V^\infty \wedge T(\nu_f) \wedge T(\nu_f) .$$

The $(2m - n)$ -dimensional ordered double point manifold

$$\begin{aligned} D_2(f) &= \{(x, y) \in M \times M \mid x \neq y \in M, f(x) = f(y) \in N\} \\ &= \{(x, y) \in M \times M \mid e(x) \neq e(y) \in V, f(x) = f(y) \in N\} \end{aligned}$$

fits into a pullback square of \mathbb{Z}_2 -equivariant embeddings

$$\begin{array}{ccc} M \sqcup D_2(f) & \xhookrightarrow{g_1 \sqcup g_2} & (LV \oplus V) \times N \\ \Delta_M \sqcup i_2 \downarrow & & \downarrow 1 \times \Delta_N \\ M \times M & \xhookrightarrow{g} & (LV \oplus V) \times N \times N \end{array}$$

with

$$\begin{aligned} g_1 &: M \hookrightarrow (LV \oplus V) \times N ; x \mapsto (0, e(x), f(x)) , \\ g_2 &: D_2(f) \hookrightarrow (LV \oplus V) \times N ; (x, y) \mapsto g(x, y) , \\ i_2 &: D_2(f) \hookrightarrow M \times M ; (x, y) \mapsto (x, y) , \end{aligned}$$

and g_2 restricts to a \mathbb{Z}_2 -equivariant embedding

$$g_3 : D_2(f) \hookrightarrow (LV \setminus \{0\} \times V) \times N .$$

The normal bundles are given by

$$\begin{aligned}\nu_{g_1} &= \nu_f \oplus \epsilon^j \oplus L\epsilon^j : M \rightarrow BO(n-m+2j) , \\ \nu_{g_2} &= \nu_{g_3} = ((\nu_f \oplus \epsilon^j) \times (\nu_f \oplus \epsilon^j))| : D_2(f) \rightarrow BO(2n-2m+2j) .\end{aligned}$$

Proposition 6.54. *Let $(e \times f : M \hookrightarrow V \times N, f : M \looparrowright N)$ be an (m, n, j) -dimensional embedding-immersion pair. Write the \mathbb{Z}_2 -equivariant Umkehr maps of g_1, g_2, g_3 as*

$$\begin{aligned}G_1 : LV^\infty \wedge V^\infty \wedge N^+ &\rightarrow T(\nu_{g_1}) = LV^\infty \wedge V^\infty \wedge T(\nu_f) , \\ G_2 : LV^\infty \wedge V^\infty \wedge N^+ &\rightarrow T(\nu_{g_2}) = LV^\infty \wedge V^\infty \wedge T(\nu_f \times \nu_f|_{D_2(f)}) , \\ G_3 : (LV \setminus \{0\})^\infty \wedge V^\infty \wedge N^+ &= \Sigma S(LV)^+ \wedge V^\infty \wedge N^+ \\ &\rightarrow T(\nu_{g_3}) = LV^\infty \wedge V^\infty \wedge T(\nu_f \times \nu_f|_{D_2(f)}) .\end{aligned}$$

(i) *The Umkehr maps for g_1, g_2, g_3 are such that*

$$G_1 = 1 \wedge F , \quad G_2 = G_3(\alpha_{LV} \wedge 1) .$$

(ii) *There is defined a homotopy*

$$h_V(F) \simeq i_2 G_3 : \Sigma S(LV)^+ \wedge V^\infty \wedge N^+ \rightarrow LV^\infty \wedge V^\infty \wedge T(\nu_f) \wedge T(\nu_f)$$

with $h_V(F) = \delta(p, q)$ the geometric Hopf invariant of F , i.e. the relative difference of the maps

$$\begin{aligned}p &= (1 \wedge \Delta_{T(\nu_f)})(1 \wedge F) , \quad q = (F \wedge F)(1 \wedge \Delta_N) : \\ &LV^\infty \wedge V^\infty \wedge N^+ \rightarrow LV^\infty \wedge V^\infty \wedge T(\nu_f) \wedge T(\nu_f) .\end{aligned}$$

which agree on $0^+ \wedge V^\infty \wedge N^+$.

Proof. (i) Immediate from the expressions of g_1, g_2 as composites

$$\begin{aligned}g_1 : M &\xrightarrow{e \times f} V \times N \hookrightarrow (LV \oplus V) \times N , \\ g_2 : D_2(f) &\xrightarrow{g_3} (LV \setminus \{0\}) \times V \times N \hookrightarrow (LV \oplus V) \times N .\end{aligned}$$

(ii) This is a special case of Theorem 6.25. □

Definition 6.55. The *(second) extended power* of a k -plane bundle $\alpha : M \rightarrow BO(k)$ with respect to an inner product space V

$$e_2(\alpha) : S(LV) \times_{\mathbb{Z}_2} (M \times M) \rightarrow BO(2k)$$

is the $2k$ -plane bundle with total space

$$E(e_2(\alpha)) = S(LV) \times_{\mathbb{Z}_2} (E(\alpha) \times E(\alpha)) .$$

□

The Thom space of the extended power bundle is given by

$$T(e_2(\alpha)) = S(LV)^+ \wedge_{\mathbb{Z}_2} (T(\alpha) \wedge T(\alpha)) .$$

Example 6.56. For the trivial k -plane bundle over a space M

$$\alpha = \epsilon^k , \quad E(\alpha) = V \times M , \quad T(\alpha) = \Sigma^k M^+$$

with $V = \mathbb{R}^k$. The isomorphism of $\mathbb{R}[\mathbb{Z}_2]$ -modules

$$\kappa_V : LV \oplus V \rightarrow V \oplus V ; (v, w) \mapsto (v + w, -v + w)$$

can be used to identify

$$e_2(\epsilon^k) = k\lambda \oplus \epsilon^k : S(LV) \times_{\mathbb{Z}_2} (M \times M) \rightarrow BO(2k)$$

where $\lambda : S(LV) \times_{\mathbb{Z}_2} (M \times M) \rightarrow BO(1)$ is the line bundle with

$$\begin{aligned} E(\lambda) &= S(LV) \times_{\mathbb{Z}_2} (L\mathbb{R} \times M \times M) , \\ T(\lambda) &= (S(LV \oplus \mathbb{R})/S(LV)) \wedge_{\mathbb{Z}_2} (M^+ \wedge M^+) , \end{aligned}$$

and

$$\begin{aligned} E(e_2(\epsilon^k)) &= S(LV) \times_{\mathbb{Z}_2} ((V \times M) \times (V \times M)) , \\ T(e_2(\epsilon^k)) &= \Sigma^k ((S(LV \oplus V)/S(LV)) \wedge_{\mathbb{Z}_2} (M^+ \wedge M^+)) . \end{aligned}$$

In the special case $k = 1$, $V = \mathbb{R}$

$$\begin{aligned} S(LV) &= S^0 , \quad \lambda = \epsilon , \\ e_2(\epsilon^k) &= \epsilon^{2k} : S(LV) \times_{\mathbb{Z}_2} (M \times M) = M \times M \rightarrow BO(2k) , \\ T(e_2(\epsilon^k)) &= \Sigma^k M^+ \wedge \Sigma^k M^+ . \end{aligned}$$

□

Proposition 6.57. *Let $(e \times f : M \hookrightarrow V \times N, f : M \looparrowright N)$ be an (m, n, j) -dimensional embedding-immersion pair, and write the $(2m - n)$ -dimensional unordered double point manifold as*

$$M' = D_2[f] .$$

(i) *The immersion*

$$f' : M' \looparrowright N ; [x, y] \mapsto f(x) = f(y)$$

has normal bundle

$$\nu_{f'} = c^* e_2(\nu_f) : M' \rightarrow BO(2(n-m))$$

with

$$c : M' \rightarrow S(LV) \times_{\mathbb{Z}_2} (M \times M) ; [x, y] \mapsto \left[\frac{e(x) - e(y)}{\|e(x) - e(y)\|}, x, y \right] .$$

(ii) The quadratic construction (5.46) on the Umkehr map $F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f)$ is given by the composite

$$\psi_V(F) : N^+ \xrightarrow{F'} T(\nu_{f'}) \xrightarrow{T(c)} T(e_2(\nu_f)) = S(LV)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))$$

with $F' : N^+ \rightarrow T(\nu_{f'})$ the Umkehr stable map of f' .

(iii) The quadratic construction $\psi_V(F)$ sends the fundamental class $[N] \in H_n(N)$ to the image of the fundamental class $[M'] \in H_{2m-n}(M'; \mathbb{Z}^{w'})$

$$\begin{aligned} \psi_V(F)_*[N] &= c_*[M'] \\ &\in \tilde{H}_n(S(LV)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))) = H_{2m-n}(S(LV) \times_{\mathbb{Z}_2} (M \times M); \mathbb{Z}^{w'}) \end{aligned}$$

where $\mathbb{Z}^{w'}$ refers to \mathbb{Z} twisted by the orientation character $w' = w_1(e_2(\nu_f))$.

Proof. This is just a matter of looking at the duality construction for the sphere $S(LV)$ and the homotopy $h_V(F) \simeq i_2 G_3$ of 6.54 (ii). \square

Example 6.58. (i) A finite covering $f : M \rightarrow N$ of an n -dimensional manifold N is a framed immersion of the n -dimensional manifold M , and there exists a map $e : M \rightarrow \mathbb{R}^j$ such that $e \times f : M \rightarrow \mathbb{R}^j \times N$ is an embedding, defining an (n, n, j) -dimensional embedding-immersion pair

$$(e \times f : M \hookrightarrow \mathbb{R}^j \times N, f : M \twoheadrightarrow N)$$

with an Umkehr stable map $F : \Sigma^j N^+ \rightarrow \Sigma^j M^+$. The quadratic construction on F is the composite

$$\psi_{\mathbb{R}^j}(F) : N^+ \xrightarrow{F'} M'^+ \xrightarrow{c} (S(L\mathbb{R}^j) \times_{\mathbb{Z}_2} (M \times M))^+$$

with $F' : N^+ \rightarrow M'^+$ the Umkehr stable map of the finite covering map

$$f' : M' = D_2[f] \rightarrow N ; [x, y] \mapsto f(x) = f(y)$$

and

$$c : M' \rightarrow S(L\mathbb{R}^j) \times_{\mathbb{Z}_2} (M \times M) ; [x, y] \mapsto \left[\frac{e(x) - e(y)}{\|e(x) - e(y)\|}, x, y \right] .$$

(ii) For a double covering $f : M \rightarrow N$ with covering translation $T : M \rightarrow M$ there is defined a canonical homeomorphism

$$N \rightarrow M' ; f(x) \mapsto [x, Tx] \quad (x \in M) ,$$

with

$$f' = 1 : M' = N \rightarrow N .$$

For any $e : M \rightarrow \mathbb{R}^j$ with $e \times f : M \rightarrow \mathbb{R}^j \times N$ an embedding the quadratic construction on the Umkehr map $F : \Sigma^j N^+ \rightarrow \Sigma^j M^+$ is given by

$$\psi_V(F) = c : N^+ \rightarrow (S^{j-1})^+ \wedge_{\mathbb{Z}_2} (M^+ \wedge M^+)$$

with

$$c : M' = N \rightarrow S^{j-1} \times_{\mathbb{Z}_2} (M \times M) ; f(x) \mapsto \left[\frac{e(x) - e(Tx)}{\|e(x) - e(Tx)\|}, x, Tx \right] .$$

(iii) The stable homotopy operation of Segal [57]

$$\theta^2 : \pi_S^0(X) = \{X; S^0\} \rightarrow \pi_S^0(X; B\Sigma_2) = \{X; (B\Sigma_2)^+\}$$

is such that for any $F : \Sigma^j X \rightarrow S^j$

$$\theta^2(F) : X \xrightarrow{\psi_{\mathbb{R}^j}(F)} (S(L\mathbb{R}^j)/\mathbb{Z}_2)^+ = P(\mathbb{R}^j)^+ \hookrightarrow (B\Sigma_2)^+$$

with

$$B\Sigma_2 = P(\mathbb{R}(\infty)) = \bigcup_{j=1}^{\infty} P(\mathbb{R}^j) .$$

□

Remark 6.59. (i) A stable map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ induces a morphism of direct sum systems

$$\begin{array}{ccccc} \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (X \wedge X)\} & \xrightleftharpoons[\delta]{\gamma} & \{X; X \wedge X\}_{\mathbb{Z}_2} & \xrightleftharpoons[\sigma]{\rho} & \{X; X\} \\ \downarrow (1 \wedge F \wedge F)_* & & \downarrow (F \wedge F)_* & & \downarrow F_* \\ \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\} & \xrightleftharpoons[\delta]{\gamma} & \{X; Y \wedge Y\}_{\mathbb{Z}_2} & \xrightleftharpoons[\sigma]{\rho} & \{X; Y\} \end{array}$$

with

$$(F \wedge F)_* = \begin{pmatrix} (1 \wedge F \wedge F)_* h_\infty(F) \\ 0 & F_* \end{pmatrix} :$$

$$\{X; X \wedge X\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (X \wedge X)\} \oplus \{X; X\} \rightarrow$$

$$\{X; Y \wedge Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\} \oplus \{X; Y\} ,$$

where $h_\infty(F) = h_{\mathbb{R}^\infty}(F) \in \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\}$. The image of

$$\Delta_X = (0, 1) \in \{X; X \wedge X\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (X \wedge X)\} \oplus \{X; X\}$$

is

$$(F \wedge F)_*(\Delta_X) = (h_\infty(F), F) \in$$

$$\{X; Y \wedge Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\} \oplus \{X; Y\} .$$

(ii) Now suppose given an (m, n, j) -dimensional embedding-immersion pair $(e \times f : M \hookrightarrow V \times N, f : M \looparrowright N)$ with Umkehr map $F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f)$. The induced morphism of direct sum systems

$$\begin{array}{ccccc} \Omega_n(S(\infty) \times_{\mathbb{Z}_2} (N \times N)) & \xrightleftharpoons[\delta]{\gamma} & \Omega_{n,n}^{\mathbb{Z}_2}(N \times N) & \xrightleftharpoons[\sigma]{\rho} & \Omega_n(N) \\ \downarrow (1 \wedge F \wedge F)_* & & \downarrow (F \wedge F)_* & & \downarrow F_* \\ \tilde{\Omega}_n(S(\infty)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))) & \xrightleftharpoons[\delta]{\gamma} & \tilde{\Omega}_{n,n}^{\mathbb{Z}_2}(T(\nu_f) \wedge T(\nu_f)) & \xrightleftharpoons[\sigma]{\rho} & \tilde{\Omega}_n(T(\nu_f)) \\ \parallel & & \parallel & & \parallel \\ \Omega_{2m-n}(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) & \xrightleftharpoons[\delta]{\gamma} & \Omega_{2m-n,m}^{\mathbb{Z}_2}(M \times M) & \xrightleftharpoons[\sigma]{\rho} & \Omega_m(M) \end{array}$$

is such that

$$(F \wedge F)_* = \begin{pmatrix} (1 \wedge F \wedge F)_* h_\infty(F) \\ 0 & F_* \end{pmatrix} :$$

$$\Omega_{n,n}^{\mathbb{Z}_2}(N \times N) = \Omega_n(S(\infty) \times_{\mathbb{Z}_2} (N \times N)) \oplus \Omega_n(N) \rightarrow$$

$$\tilde{\Omega}_{n,n}^{\mathbb{Z}_2}(T(\nu_f) \wedge T(\nu_f)) = \tilde{\Omega}_n(S(\infty)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))) \oplus \tilde{\Omega}_n(T(\nu_f))$$

$$= \Omega_{2m-n,m}^{\mathbb{Z}_2}(M \times M) = \Omega_{2m-n}(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) \oplus \Omega_m(M) .$$

The image of

$$(N, \Delta_N) = (0, (N, 1)) \in \Omega_{n,n}(N \times N) = \Omega_n(S(\infty) \times_{\mathbb{Z}_2} (N \times N)) \oplus \Omega_n(N)$$

is

$$(F \wedge F)_*(N, \Delta_N) = ((D_2[f], i_2), (M, 1)) \in \\ \Omega_{2m-n, m}^{\mathbb{Z}_2}(M \times M) = \Omega_{2m-n}(S(\infty) \times_{\mathbb{Z}_2} (M \times M)) \oplus \Omega_m(M) .$$

(iii) For any pointed spaces X, Y there is defined a nonadditive function of direct sum systems

$$\begin{array}{ccccc} \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} Y\} & \xrightleftharpoons[\delta]{\gamma} & \{X; Y\}_{\mathbb{Z}_2} & \xrightleftharpoons[\sigma]{\rho} & \{X; Y\} \\ \downarrow 1 \wedge P^2 & & \downarrow P^2 & & \parallel \\ \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\} & \xrightleftharpoons[\delta]{\gamma} & \{X; Y \wedge Y\}_{\mathbb{Z}_2} & \xrightleftharpoons[\sigma]{\rho} & \{X; Y\} \end{array}$$

with

$$\begin{aligned} P^2 &: \{X; Y\}_{\mathbb{Z}_2} \rightarrow \{X; Y \wedge Y\}_{\mathbb{Z}_2} ; F \mapsto (F \wedge F) \Delta_X , \\ P^2 &= \begin{pmatrix} 1 \wedge P^2 & h_\infty(E) \\ 0 & 1 \end{pmatrix} : \\ \{X; Y\}_{\mathbb{Z}_2} &= \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} Y\} \oplus \{X; Y\} \rightarrow \\ \{X; Y \wedge Y\}_{\mathbb{Z}_2} &= \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\} \oplus \{X; Y\} , \end{aligned}$$

using $\{X; Y\} = [X, \Omega^\infty \Sigma^\infty Y]$ and the geometric Hopf invariant

$$h_\infty(E) \in \{\Omega^\infty \Sigma^\infty Y; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\}$$

of the evaluation stable map

$$E : \Sigma^\infty(\Omega^\infty \Sigma^\infty Y) \rightarrow \Sigma^\infty Y$$

(Proposition 6.40). For any stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ the image of

$$F = (0, F) \in \{X; Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} Y\} \oplus \{X; Y\}$$

is

$$\begin{aligned} P^2(F) &= (h_\infty(E)(F), F) = (h_\infty(F), F) \\ &\in \{X; Y \wedge Y\}_{\mathbb{Z}_2} = \{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\} \oplus \{X; Y\} . \end{aligned}$$

(iv) For an immersion $f : M^m \looparrowright N^n$ consider the nonadditive function of direct sum systems

$$\begin{array}{ccccc}
\{N^+; S(\infty) \wedge_{\mathbb{Z}_2} T(\nu_f)\} & \xrightleftharpoons[\delta]{\gamma} & \{N^+; T(\nu_f)\}_{\mathbb{Z}_2} & \xrightleftharpoons[\sigma]{\rho} & \{N^+; T(\nu_f)\} \\
\downarrow 1 \wedge P^2 & & \downarrow P^2 & & \parallel \\
\{N^+; S(\infty)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))\} & \xrightleftharpoons[\delta]{\gamma} & \{N^+; T(\nu_f) \wedge T(\nu_f)\}_{\mathbb{Z}_2} & \xrightleftharpoons[\sigma]{\rho} & \{N^+; T(\nu_f)\}
\end{array}$$

with

$$\begin{aligned}
P^2 &= \begin{pmatrix} 1 \wedge P^2 h_\infty(E) \\ 0 & 1 \end{pmatrix} : \\
\{N^+; T(\nu_f)\}_{\mathbb{Z}_2} &= \{N^+; S(\infty) \wedge_{\mathbb{Z}_2} T(\nu_f)\} \oplus \{N^+; T(\nu_f)\} \rightarrow \\
\{N^+; T(\nu_f) \wedge T(\nu_f)\}_{\mathbb{Z}_2} &= \{N^+; S(\infty)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))\} \oplus \{N^+; T(\nu_f)\} .
\end{aligned}$$

The image of the Umkehr map $F : \Sigma^\infty N^+ \rightarrow \Sigma^\infty T(\nu_f)$

$$F = (0, F) \in \{N^+; T(\nu_f)\}_{\mathbb{Z}_2} = \{N^+; S(\infty) \wedge_{\mathbb{Z}_2} T(\nu_f)\} \oplus \{N^+; T(\nu_f)\}$$

is

$$\begin{aligned}
P^2(F) &= (h_\infty(F), F) = (D_2[f], F) \in \\
\{N^+; T(\nu_f) \wedge T(\nu_f)\}_{\mathbb{Z}_2} &= \{N^+; S(\infty)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))\} \oplus \{N^+; T(\nu_f)\} ,
\end{aligned}$$

with

$$h_\infty(F) = D_2[f] \in \{N^+; S(\infty)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))\}$$

a regular homotopy invariant of the immersion $f : N \looparrowright M$, and

$$\begin{aligned}
[F] &= [h_\infty(F)] = [D_2[f]] \\
&\in \text{coker}(P^2 : \{N^+; T(\nu_f)\}_{\mathbb{Z}_2} \rightarrow \{N^+; T(\nu_f) \wedge T(\nu_f)\}_{\mathbb{Z}_2}) \\
&= \text{coker}(1 \wedge P^2 : \{N^+; S(\infty)^+ \wedge_{\mathbb{Z}_2} T(\nu_f)\} \rightarrow \\
&\quad \{N^+; S(\infty)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))\})
\end{aligned}$$

a homotopy invariant of f .

□

For any manifold A , bundle $\gamma : B \rightarrow BO(k)$ and integer $\ell \geq 0$ let $\Omega_\ell(A, B, \gamma)$ be the bordism group of pairs

$$(\text{immersion } L^\ell \looparrowright A, \text{ stable bundle map } (L, \nu_{L \looparrowright A}) \rightarrow (B, \gamma)) .$$

The Pontryagin-Thom construction identifies this bordism group with a stable homotopy group

$$\Omega_\ell(A, B, \gamma) = \{A^\infty, T(\gamma)\} .$$

Definition 6.60. The *double point Hopf invariant* of an (m, n, j) -dimensional embedding-immersion pair $(e \times f : M^m \hookrightarrow V \times N^n, f : M \looparrowright N)$ is the bordism class

$$(M', f', c) \in \Omega_{2m-n}(N, S(LV) \times_{\mathbb{Z}_2} (M \times M), e_2(\nu_f))$$

of the immersion of the $(2m - n)$ -dimensional unordered double point manifold

$$f' : M' = D_2[f] \looparrowright N ; [x, y] \mapsto f(x) = f(y)$$

with

$$c : M' \rightarrow S(LV) \times_{\mathbb{Z}_2} (M \times M) ; [x, y] \mapsto \left[\frac{e(x) - e(y)}{\|e(x) - e(y)\|}, x, y \right]. \quad \square$$

Proposition 6.61. The *double point Hopf invariant* is the stable homotopy class of the quadratic construction (5.46) on the Umkehr map $F : V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f)$

$$\psi_V(F) : N^+ \xrightarrow{F'} T(\nu_{f'}) \xrightarrow{T(c)} S(LV)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))$$

with $F' : N^+ \rightarrow T(\nu_{f'})$ the Umkehr stable map of f'

$$\begin{aligned} (M', f', c) &= \psi_V(F) \in \Omega_{2m-n}(N, S(LV) \times_{\mathbb{Z}_2} (M \times M), e_2(\nu_f)) \\ &= \{N^+, S(LV)^+ \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))\}. \end{aligned}$$

□

For any $j, k \geq 1$ the normal bundle of the standard embedding of real projective spaces $P(\mathbb{R}^j) \subset P(\mathbb{R}^{j+k})$ is the Whitney sum of k copies of the canonical line bundle $\lambda : P(\mathbb{R}^j) \rightarrow BO(1)$

$$\nu_{P(\mathbb{R}^j) \subset P(\mathbb{R}^{j+k})} = k\lambda : P(\mathbb{R}^j) \rightarrow BO(k)$$

with bundle projection

$$E(k\lambda) = S(L\mathbb{R}^j) \times_{\mathbb{Z}_2} L\mathbb{R}^k \rightarrow S(L\mathbb{R}^j)/\mathbb{Z}_2 = P(\mathbb{R}^j)$$

and Thom space the stunted projective space

$$T(k\lambda) = (S(L\mathbb{R}^{j+k})/S(L\mathbb{R}^k))/\mathbb{Z}_2 = P(\mathbb{R}^{j+k})/P(\mathbb{R}^k).$$

The bordism group of triples

(ℓ -dimensional manifold L^ℓ , map $L \rightarrow P(\mathbb{R}^j)$, stable bundle map $\nu_L \rightarrow k\lambda$)

is given by the Pontryagin-Thom isomorphism to be

$$\Omega_\ell(P(\mathbb{R}^j), k\lambda) = \pi_{k+\ell}^S(P(\mathbb{R}^{j+k})/P(\mathbb{R}^k)) .$$

In particular, for $j = 0$ this is

$$\Omega_\ell^{fr} = \pi_\ell^S .$$

The Stiefel manifold $V_{j,k}$ ($1 \leq k \leq j$) of orthonormal k -frames in \mathbb{R}^j fits into a fibration

$$V_{j,k} = O(j)/O(j-k) \rightarrow BO(j-k) \rightarrow BO(j) .$$

The canonical embedding

$$P(\mathbb{R}^j)/P(\mathbb{R}^{j-k}) \hookrightarrow V_{j,k} ; x = [x_1, x_2, \dots, x_j] \mapsto [\mathbf{v}_{j-k+1}, \mathbf{v}_{j-k+2}, \dots, \mathbf{v}_j]$$

is defined using the columns \mathbf{v}_ℓ of the orthogonal $j \times j$ -matrix

$$(\delta_{pq} - 2x_p x_q)_{1 \leq p, q \leq j} \left(\sum_{i=1}^j (x_i)^2 = 1 \right)$$

of the reflection in the hyperplane $x^\perp \subset \mathbb{P}(\mathbb{R}^j)$. The pair $(V_{j,k}, P(\mathbb{R}^j)/P(\mathbb{R}^{j-k}))$ is $2(j-k)$ -connected (James [30, p. 5]).

Remark 6.62. Let $N = S^n$.

(i) The double point Hopf invariant (6.60) of an (m, n, j) -dimensional embedding-immersion pair

$$(e \times f : M^m \hookrightarrow \mathbb{R}^j \times S^n, f : M \looparrowright S^n)$$

is the bordism class of the $(2m-n)$ -dimensional unordered double point manifold

$$\begin{aligned} (f' : M' \looparrowright S^n) &= \psi_{\mathbb{R}^j}(F|) \\ &\in \{S^n; (S^{j-1})^\infty \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))\} = \Omega_{2m-n}(S^{j-1} \times_{\mathbb{Z}_2} (M \times M), e_2(\nu_f)) \end{aligned}$$

with

$$f' : M' = D_2[f]^{2m-n} \looparrowright S^n ; [x, y] \mapsto f(x) = f(y)$$

and $F| : S^{n+j} \rightarrow \Sigma^j T(\nu_f)$ the restriction of the Umkehr map of $e \times f$

$$F : (\mathbb{R}^j \times S^n)^\infty = S^{n+j} \vee S^j \rightarrow \Sigma^j T(\nu_f) .$$

(ii) The bordism class of an (m, n, j) -dimensional embedding-immersion pair $(e \times f : M^m \hookrightarrow \mathbb{R}^j \times S^n, f : M \looparrowright S^n)$ with a framing $\delta\nu_f : \nu_f \cong \epsilon^{n-m}$ is the homotopy class

$$(e \times f : M^m \hookrightarrow \mathbb{R}^j \times S^n, f : M \looparrowright S^n, \delta\nu_f) = \phi \in \pi_{n+j}(S^{n-m+j})$$

of the Pontryagin-Thom map

$$\begin{aligned} \phi : S^{n+j} &\longrightarrow (\mathbb{R}^j \times S^n)^\infty = S^{n+j} \vee S^j \\ &\xrightarrow{F} \Sigma^j T(\nu_f) \xrightarrow{\Sigma^j T(\delta\nu_f)} \Sigma^{n-m+j} M^+ \rightarrow S^{n-m+j} . \end{aligned}$$

As in 6.57 (i) the normal bundle of the immersion $f : M' = D_2[f]^{2m-n} \looparrowright S^n$ is given by

$$\nu_{f'} = d^*(e_2(\epsilon^{n-m})) : M' \rightarrow BO(2(n-m))$$

with

$$d : M' \rightarrow P(\mathbb{R}^j) ; [x, y] \mapsto \left[\frac{e(x) - e(y)}{\|e(x) - e(y)\|} \right]$$

and

$$\begin{aligned} e_2(\epsilon^{n-m}) &= (n-m)\lambda \oplus \epsilon^{n-m} : \\ S(L\mathbb{R}^j) \times_{\mathbb{Z}_2} (\{\text{pt.}\} \times \{\text{pt.}\}) &= P(\mathbb{R}^j) \rightarrow BO(2(n-m)) \end{aligned}$$

with

$$\begin{aligned} E(e_2(\epsilon^{n-m})) &= S(L\mathbb{R}^j) \times_{\mathbb{Z}_2} (L\mathbb{R}^{n-m} \oplus \mathbb{R}^{n-m}) , \\ T(e_2(\epsilon^{n-m})) &= \Sigma^{n-m}(P(\mathbb{R}^{n-m+j})/P(\mathbb{R}^{n-m})) . \end{aligned}$$

The double point Hopf invariant of Koschorke and Sanderson [37] is the bordism class

$$\begin{aligned} (f' : M' \looparrowright S^n) &= \psi_{\mathbb{R}^j}(\phi) \\ &\in \Omega_{2m-n}(P(\mathbb{R}^j), (n-m)\lambda) = \pi_m^S(P(\mathbb{R}^{j+n-m})/P(\mathbb{R}^{n-m})) \end{aligned}$$

of the quadratic construction

$$\psi_{\mathbb{R}^j}(\phi) : S^n \looparrowright (S^{j-1})^\infty \wedge_{\mathbb{Z}_2} (S^{n-m} \wedge S^{n-m}) = \Sigma^{n-m}(P(\mathbb{R}^{j+n-m})/P(\mathbb{R}^{n-m})) .$$

For $n \leq 2(n-m)$ the double point Hopf invariant map

$$\begin{aligned} H^j : \pi_{n+j}(S^{n-m+j}) &\rightarrow \Omega_{2m-n}(P(\mathbb{R}^j), (n-m)\lambda) = \pi_m(V_{n-m+j,j}) ; \\ (e \times f : M^m \hookrightarrow \mathbb{R}^j \times S^n, f : M \looparrowright S^n, \delta\nu_f) &\mapsto (f' : M' \looparrowright S^n) = \psi_{\mathbb{R}^j}(\phi) \end{aligned}$$

fits into the *EHP* exact sequence of James [27]

$$\pi_n(S^{n-m}) \xrightarrow{E^j} \pi_{n+j}(S^{n-m+j}) \xrightarrow{H^j} \pi_m(V_{n-m+j,j}) \xrightarrow{P^j} \pi_{n-1}(S^{n-m}) \rightarrow \dots ,$$

with

$$\begin{aligned} \pi_m(V_{n-m+j,j}) &= \pi_m(P(\mathbb{R}^{j+n-m})/P(\mathbb{R}^{n-m})) \\ &= \pi_m^S(P(\mathbb{R}^{j+n-m})/P(\mathbb{R}^{n-m})) = \Omega_{2m-n}(P(\mathbb{R}^j), (n-m)\lambda) . \end{aligned}$$

The J -homomorphism defines a natural transformation of exact sequences

$$\begin{array}{ccccccc} \pi_m(SO(n-m)) & \rightarrow & \pi_m(SO(j+n-m)) & \rightarrow & \pi_m(V_{j+n-m,j}) & \rightarrow & \pi_{m-1}(SO(n-m)) \\ \downarrow J & & \downarrow J & & \parallel & & \downarrow J \\ \pi_n(S^{n-m}) & \xrightarrow{E^j} & \pi_{j+n}(S^{j+n-m}) & \xrightarrow{H^j} & \pi_m(V_{j+n-m,j}) & \xrightarrow{P^j} & \pi_{n-1}(S^{n-m}) \end{array}$$

(iii) For $n \leq 2(n-m)$, $j = 1$ the exact sequence in (ii) is just the EHP sequence

$$\pi_n(S^{n-m}) \xrightarrow{E} \pi_{n+1}(S^{n-m+1}) \xrightarrow{H} \pi_m(S^{n-m}) \xrightarrow{P} \pi_{n-1}(S^{n-m}) \rightarrow \dots,$$

with $V_{1+n-m,1} = S^{n-m}$, E the suspension map and H the classical Hopf invariant, interpreted as sending an $(m, n, 1)$ -dimensional embedding-immersion pair $(e \times f : M^m \hookrightarrow \mathbb{R} \times S^n, f : M \looparrowright S^n)$ with a framing $\delta\nu_f : \nu_f \cong \epsilon^{n-m}$ to the double point Hopf invariant

$$(f' : M' \looparrowright S^n) = \psi_{\mathbb{R}}(\phi) \in \Omega_{2m-n}^{fr} = \pi_m(V_{n-m+1,1}) = \pi_m(S^{n-m}) = \pi_{2m-n}^S.$$

For $n = 2m \geq 2$ the EHP sequence is

$$\pi_{2m}(S^m) \xrightarrow{E} \pi_{2m+1}(S^{m+1}) \xrightarrow{H} \pi_m(S^m) = \mathbb{Z} \xrightarrow{P} \pi_{2m-1}(S^m) \rightarrow \dots$$

In particular, for $m = 1$, $n = 2$ the generator $1 \in \pi_3(S^2)$ is the cobordism class of the $(1, 2, 1)$ -dimensional embedding-immersion pair

$$(e \times f : S^1 \hookrightarrow \mathbb{R} \times S^2, f : S^1 \looparrowright S^2)$$

with f the figure 8 immersion and $\delta\nu_f \cong \epsilon$ one of the two framings. The Hopf invariant 1 element

$$(e \times f : S^1 \hookrightarrow \mathbb{R} \times S^2, f : S^1 \looparrowright S^2) = \eta = 1 \in \pi_3(S^2) = \mathbb{Z}$$

is detected by the cobordism class of the 0-dimensional unordered double point manifold $M' = \{*\}$ of the immersion $f : M = S^1 \looparrowright S^2$

$$(f' : M' \looparrowright S^2) = \psi_{\mathbb{R}}(\phi) = 1 \in \Omega_0^{fr} = \pi_0^S = \mathbb{Z}.$$

□

6.7 Linking and self-linking

The original invariant of Hopf [10] was detected by the linking of k -dimensional submanifolds of S^{2k+1}

$$\begin{aligned} H &: \pi_{2k+1}(S^{k+1}) \rightarrow \mathbb{Z} ; \\ (\phi : S^{2k+1} \rightarrow S^{k+1}) &\mapsto \text{linking number}((M_1)^k \cup (M_2)^k \subset S^{2k+1}) \end{aligned}$$

with $M_1 = \phi^{-1}(x_1)$, $M_2 = \phi^{-1}(x_2) \subset S^{2k+1}$ for distinct regular values $x_1, x_2 \in S^{k+1}$ of ϕ . The homotopy-theoretic construction of the generalized Hopf invariant H of G.W. Whitehead [71] was interpreted by Kervaire [33] and Haefliger and Steer [20, p. 262] in terms of higher linking numbers. The Hopf invariant constructions λ_2, μ_2 of Boardman and Steer [4] (recalled in §§5.3,4 above) were motivated by the linking and self-linking of embedded submanifolds. We shall now interpret the constructions in terms of immersed submanifolds.

Let M_1, M_2, N be closed manifolds such that

$$\dim M_1 = m_1, \quad \dim M_2 = m_2, \quad \dim N = n,$$

and suppose given disjoint embeddings

$$e_i \times f_i : M_i \hookrightarrow \mathbb{R} \times N \quad (i = 1, 2)$$

which intersect transversely, with $f_i : M_i \looparrowright N$ immersions. The normal bundle of $e_i \times f_i : M_i \hookrightarrow \mathbb{R} \times N$ is

$$\nu_{e_i \times f_i} = \nu_{f_i} \oplus \epsilon : M_i \rightarrow BO(n - m_i + 1)$$

with $\nu_{f_i} : M_i \rightarrow BO(n - m_i)$ the normal bundle of $f_i : M_i \looparrowright N$.

Definition 6.63. The *linking manifold* of the submanifolds $M_1, M_2 \subset \mathbb{R} \times N$ is the $(m_1 + m_2 - n)$ -dimensional submanifold of $M_1 \times M_2$

$$\begin{aligned} L(M_1, M_2, N) &= \{(x_1, x_2) \in M_1 \times M_2 \mid e_1(x_1) < e_2(x_2) \in \mathbb{R}, f_1(x_1) = f_2(x_2) \in N\}. \end{aligned}$$

□

Proposition 6.64. (i) *The linking manifold $L(M_1, M_2, N)$ is homeomorphic to the intersection in $\mathbb{R} \times N \times I$ of the tracks of isotopies unlinking M_1, M_2 .*

(ii) *The embedding*

$$g : L(M_1, M_2, N) \hookrightarrow \mathbb{R} \times \mathbb{R} \times N ;$$

$$(x_1, x_2) \mapsto (e_1(x_1), e_2(x_2), y) \quad (y = f_1(x_1) = f_2(x_2))$$

has normal bundle

$$\nu_g = ((\nu_{f_1} \oplus \epsilon) \times (\nu_{f_2} \oplus \epsilon))| : L(M_1, M_2, N) \rightarrow BO(2n - m_1 - m_2 + 2) .$$

Proof. (i) Assume that $e_1(M_1), e_2(M_2) \subset (0, 1)$, and define isotopies

$$h_1 : M_1 \times I \rightarrow \mathbb{R} \times N \times I ; (x, s) \mapsto (e_1(x) + s, f_1(x), s) ,$$

$$h_2 : M_2 \times I \rightarrow \mathbb{R} \times N \times I ; (y, t) \mapsto (e_2(y), f_2(y), t)$$

such that the submanifolds

$$h_1(M_1 \times \{1\}) , \quad h_2(M_2 \times \{1\}) \subset \mathbb{R} \times N \times \{1\}$$

isotopic to the submanifolds

$$M_1 = h_1(M_1 \times \{0\}) , \quad M_2 = h_2(M_2 \times \{0\}) \subset \mathbb{R} \times N \times \{0\}$$

are disjoint and unlinked, with

$$h_1(M_1 \times \{1\}) \subset \{(a, z, 1) \in \mathbb{R} \times N \times \{1\} \mid a > 1, z \in N\} ,$$

$$h_2(M_2 \times \{1\}) \subset \{(b, z, 1) \in \mathbb{R} \times N \times \{1\} \mid b < 1, z \in N\}$$

separated by $\{1\} \times N \times \{1\}$. The tracks of the isotopies are $(m_i + 1)$ -dimensional submanifolds

$$L_i = h_i(M_i \times I) \subset \mathbb{R} \times N \times I \quad (i = 1, 2)$$

and there is defined a homeomorphism

$$L(M_1, M_2, N) \rightarrow L_1 \cap L_2 ;$$

$$(x_1, x_2) \mapsto h_1(x_1, e_2(x_2) - e_1(x_1)) = h_2(x_2, e_2(x_2) - e_1(x_1)) .$$

(ii) The submanifolds $L_i \subset \mathbb{R} \times \mathbb{R} \times N$ ($i = 1, 2$) intersect transversely, with

$$\nu_{L_i \subset \mathbb{R} \times \mathbb{R} \times N} : L_i \longrightarrow M_i \xrightarrow{\nu_{f_i} \oplus \epsilon} BO(n - m_i + 1)$$

so that

$$\nu_g = ((\nu_{L_1 \subset \mathbb{R} \times \mathbb{R} \times N} \times (\nu_{L_2 \subset \mathbb{R} \times \mathbb{R} \times N}))|$$

$$= ((\nu_{f_1} \oplus \epsilon) \times (\nu_{f_2} \oplus \epsilon))| : L(M_1, M_2, N) \rightarrow BO(2n - m_1 - m_2 + 2) .$$

□

Let

$$F_i : \Sigma N^+ = (\mathbb{R} \times N)^\infty \rightarrow T(\nu_{e_i \times f_i}) = \Sigma T(\nu_{f_i})$$

be the Umkehr stable maps given by the Pontryagin-Thom construction. The embedding of the disjoint union

$$e_1 \times f_1 \sqcup e_2 \times f_2 : M_1 \sqcup M_2 \hookrightarrow \mathbb{R} \times N$$

has a compactification Umkehr stable map

$$F : \Sigma N^+ \rightarrow \Sigma T(\nu_{f_1}) \vee \Sigma T(\nu_{f_2}) ; (t, x) \mapsto \begin{cases} F_1(2t, x) & \text{if } 0 \leq t \leq 1/2 \\ F_2(2t - 1, x) & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

Proposition 6.65. (i) *The map of Boardman and Steer [4]*

$$\mu_2(F) : \Sigma^2 N^+ \rightarrow \Sigma T(\nu_{f_1}) \wedge \Sigma T(\nu_{f_2}) ; (s, t, x) \mapsto \begin{cases} (F(s, x), F(t, x)) & \text{if } s \leq t \\ * & \text{otherwise} \end{cases}$$

is the composite

$$\mu_2(F) = T(i)G : \Sigma^2 N^+ \xrightarrow{G} T(\nu_g) \xrightarrow{T(i)} \Sigma T(\nu_{f_1}) \wedge \Sigma T(\nu_{f_2})$$

of the Umkehr stable map for $g : L(M_1, M_2, N) \hookrightarrow \mathbb{R} \times \mathbb{R} \times N$

$$G : (\mathbb{R} \times \mathbb{R} \times N)^\infty = \Sigma^2 N^+ \rightarrow T(\nu_g)$$

and the inclusion of Thom spaces

$$T(i) : T(\nu_g) \rightarrow T((\nu_{f_1} \oplus \epsilon) \times (\nu_{f_2} \oplus \epsilon)) = \Sigma T(\nu_{f_1}) \wedge \Sigma T(\nu_{f_2})$$

induced by the inclusion

$$i : L(M_1, M_2, N) \hookrightarrow M_1 \times M_2 ; (x_1, x_2) \mapsto (x_1, x_2) .$$

(ii) *The image of the fundamental class $[N] \in H_n(N)$ under $\mu_2(F)$ is*

$$\mu_2(F)_*[N] = i_*[L] \in \widetilde{H}_n(T(\nu_{f_1}) \wedge T(\nu_{f_2})) = H_{m_1+m_2-n}(M_1 \times M_2)$$

with $[L] \in H_{m_1+m_2-n}(L)$ the fundamental class of $L = L(M_1, M_2, N)$. Viewing $i_*[L] \in H_{m_1+m_2-n}(M_1 \times M_2)$ as a chain map

$$i_*[L] : C(M_1)^{m_1-*} \rightarrow C(M_2)_{*-n+m_2}$$

(up to chain homotopy) there is defined a commutative diagram

$$\begin{array}{ccc}
H^{m_1-*}(M_1) & \xrightarrow{i_*[L]} & H_{*-n+m_2}(M_2) \cong H_{*+1}(\mathbb{R} \times N, \mathbb{R} \times N \setminus (e_2 \times f_2)(M_2)) \\
[M_1] \cap - \downarrow \cong & & \downarrow \\
H_*(M_1) & \xrightarrow{(e_1 \times f_1)_*} & H_*(\mathbb{R} \times N \setminus (e_2 \times f_2)(M_2)) .
\end{array}$$

□

Example 6.66. The linking manifold $L = L(M_1, M_2, S^n)$ of $M_1, M_2 \subset \mathbb{R} \times S^n$ with $m_1 + m_2 = n$ is 0-dimensional, and

$$\mu_2(F)_*[N] = i_*[L] \in H_0(M_1 \times M_2) = \mathbb{Z}$$

may be identified with

$$\begin{aligned}
\text{linking number}(M_1 \cup M_2 \subset \mathbb{R} \times S^n) &= (e_1 \times f_1)_*[M_1] \\
&\in H_{m_1}((\mathbb{R} \times S^n) \setminus (e_2 \times f_2)(M_2)) = H^{m_2}(M_2) = \mathbb{Z} .
\end{aligned}$$

□

Proposition 6.67. (Boardman and Steer [4]) *Let N be an n -dimensional manifold. The Hopf map*

$$H : [\Sigma N^+, S^{k+1}] \rightarrow [\Sigma^2 N^+, S^{2k+2}] ; \phi \mapsto h_{\mathbb{R}}(\phi)$$

sends $\phi : \Sigma N^+ \rightarrow S^{k+1}$ to the cobordism class of the framed submanifold

$$L(M_1, M_2, N)^{n-2k} \subset \mathbb{R} \times \mathbb{R} \times N$$

with

$$M_i = \phi^{-1}(x_i) \subset \mathbb{R} \times N \quad (i = 1, 2)$$

for distinct regular values $x_1, x_2 \in S^{k+1}$, regarding $[\Sigma^2 N^+, S^{2k+2}]$ as the bordism group of framed embeddings $L^{n-2k} \subset \mathbb{R} \times \mathbb{R} \times N$.

Proof. It may be assumed that the inclusions are framed embeddings $e_i \times f_i : M_i \hookrightarrow \mathbb{R} \times N$ with $f_i : M_i \looparrowright N$ immersions. By 6.65

$$H(\phi) : \Sigma^2 N^+ \xrightarrow{\mu_2(F)} \Sigma T(\nu_{f_1}) \wedge \Sigma T(\nu_{f_2}) = \Sigma^{2k+2}(M_1 \times M_2)^\infty$$

is such that

$$H(\phi)^{-1}(M_1 \times M_2) = L(M_1, M_2, N)^{n-2k} \subset \mathbb{R} \times \mathbb{R} \times N .$$

It follows that

$$h_{\mathbb{R}}(\phi) : \Sigma^2 N^+ \xrightarrow{H(\phi)} \Sigma^{2k+2}(M_1 \times M_2)^+ \rightarrow S^{2k+2}$$

is such that

$$h_{\mathbb{R}}(\phi)^{-1}(*) = L(M_1, M_2, N)^{n-2k} \subset \mathbb{R} \times \mathbb{R} \times N .$$

□

Example 6.68. (i) For $N = S^{2k}$ 6.67 gives the original invariant of Hopf [26]

$$\begin{aligned} H : \pi_{2k+1}(S^{k+1}) &\rightarrow \pi_{2k+2}(S^{2k+2}) = \mathbb{Z} ; \\ \phi &\mapsto H(\phi) = \text{linking number}((M_1)^k \cup (M_2)^k \subset S^{2k+1}) \end{aligned}$$

with $(M_i)^k = \phi^{-1}(x_i)$.

(ii) For $N = S^n$ 6.67 gives the generalized Hopf invariant

$$H : \pi_{n+1}(S^{k+1}) \rightarrow \pi_{n+2}(S^{2k+2}) ; \phi \mapsto H(\phi) .$$

of G.W. Whitehead [71] and Hilton [23], with

$$\begin{aligned} H(\phi) &= \text{framed cobordism class}(L(M_1, M_2, S^n)^{n-2k} \subset S^{n+2}) \in \pi_{n+2}(S^{2k+2}) \\ ((M_i)^{n-k} &= \phi^{-1}(x_i) \subset \mathbb{R} \times S^n \subset S^{n+1} \quad (i = 1, 2)) \end{aligned}$$

as in Kervaire [33].

□

Definition 6.69. Let $e \times f : M \hookrightarrow \mathbb{R} \times N$ be an embedding of an m -dimensional manifold M , with N an n -dimensional manifold and $f : M \looparrowright N$ an immersion. The *self-linking manifold* is the $(2m-n)$ -dimensional manifold

$$L(M, N) = \{(x, y) \in M \times M \mid e(x) < e(y) \in \mathbb{R}, f(x) = f(y) \in N\} .$$

□

Proposition 6.70. (i) *The ordered double point manifold*

$$D_2(f) = \{(x, y) \in M \times M \mid x \neq y \in M, f(x) = f(y) \in N\}$$

is the disjoint union of the self-linking manifold and its transpose

$$D_2(f) = L(M, N) \sqcup TL(M, N) ,$$

with

$$TL(M, N) = \{(x, y) \in M \times M \mid e(x) > e(y) \in \mathbb{R}, f(x) = f(y) \in N\} .$$

(ii) *The embedding*

$$g : L(M, N) \hookrightarrow \mathbb{R} \times \mathbb{R} \times N ; (x, y) \mapsto (e(x), e(y), f(x))$$

has normal bundle

$$\begin{aligned} \nu_g &= ((\nu_f \oplus \epsilon) \times (\nu_f \oplus \epsilon))| \\ &= ((\nu_f \times \nu_f) \oplus \epsilon^2)| : L(M, N) \rightarrow BO(2n - 2m + 2) \end{aligned}$$

with $\nu_f : M \rightarrow BO(n - m)$ the normal bundle of f .

(iii) *The immersion*

$$g_2 : L(M, N) \hookrightarrow N ; (x, y) \mapsto f(x)$$

has normal bundle

$$\nu_{g_2} = (\nu_f \times \nu_f)| : L(M, N) \rightarrow BO(2n - 2m) .$$

(iv) *The immersion $f : M \looparrowright N$ is an embedding if and only if $L(M, N) = \emptyset$.* \square

Let $F : \Sigma N^+ \rightarrow \Sigma T(\nu_f)$ be the Umkehr stable map of $e \times f$.

Proposition 6.71. (i) *The Hopf invariant map of Boardman and Steer [4]*

$$\lambda_2(F) = \mu_2(\nabla F) : \Sigma^2 N^+ \rightarrow \Sigma T(\nu_f) \wedge \Sigma T(\nu_f)$$

(identified with $h_{\mathbb{R}}(F)$ in 5.57) is the composite

$$h_{\mathbb{R}}(F) = \lambda_2(F) = T(i)G : \Sigma^2 N^+ \xrightarrow{G} T(\nu_g) \xrightarrow{T(i)} \Sigma T(\nu_f) \wedge \Sigma T(\nu_f)$$

of the Umkehr stable map of g

$$G : (\mathbb{R} \times \mathbb{R} \times N)^\infty = \Sigma^2 N^+ \rightarrow T(\nu_g)$$

and the inclusion of Thom spaces

$$T(i) : T(\nu_g) \rightarrow T((\nu_f \oplus \epsilon) \times (\nu_f \oplus \epsilon)) = \Sigma T(\nu_f) \wedge \Sigma T(\nu_f)$$

induced by the inclusion

$$i : L(M, N) \hookrightarrow M \times M ; (x_1, x_2) \mapsto (x_1, x_2) .$$

(ii) *The double point Hopf invariant (6.60) is the bordism class of the immersion of the self-linking manifold $g_2 : L(M, N) \looparrowright N^n$, which is the stable homotopy class of the ultraquadratic construction (5.52)*

$$\psi_{\mathbb{R}}(F) : N^+ \xrightarrow{G_2} T(\nu_{g_2}) \xrightarrow{T(i)} T(\nu_f) \wedge T(\nu_f)$$

with G_2 the Umkehr stable map of g_2 .

(iii) The image of the fundamental class $[N] \in H_n(N)$ under $h_{\mathbb{R}}(F)$ is

$$h_{\mathbb{R}}(F)_*[N] = i_*[L(M, N)] \in \tilde{H}_n(T(\nu_f) \wedge T(\nu_f)) = H_{2m-n}(M \times M)$$

with $[L(M, N)] \in H_{2m-n}(L(M, N))$ the fundamental class of $L(M, N)$.

Proof. (i) Apply 6.65 with

$$M_1 = M_2 = M, \quad f_1 = f_2 = f : M \rightarrow N, \quad e_1 = e + c, \quad e_2 = e$$

for some $c > 0$ so small that

$$c < e(y) - e(x) \quad ((x, y) \in L(M, N)),$$

in which case

$$e_1 \times f_1 : M_1 \hookrightarrow \mathbb{R} \times N, \quad e_2 \times f_2 : M_2 \hookrightarrow \mathbb{R} \times N$$

are disjoint embeddings, and the linking manifold is

$$L(M_1, M_2, N) = L(M, N).$$

The embedding of the disjoint union

$$e_1 \times f \sqcup e_2 \times f : M \sqcup M \hookrightarrow \mathbb{R} \times N$$

has compactification Umkehr stable map

$$\nabla F : \Sigma N^+ \rightarrow \Sigma T(\nu_f) \vee \Sigma T(\nu_f); (t, x) \mapsto \begin{cases} F(2t, x)_1 & \text{if } 0 \leq t \leq 1/2 \\ F(2t - 1, x)_2 & \text{if } 1/2 \leq t \leq 1 \end{cases}$$

with projections

$$\pi_i(\nabla F) \simeq F : \Sigma N^+ \rightarrow \Sigma T(\nu_f).$$

(ii) This is just the special case $j = 1$ of 6.61 (i).

(iii) Immediate from (ii). □

The fibration

$$V_{k+1,1} = S^k \xrightarrow{\tau_{S^k}} BO(k) \longrightarrow BO(k+1)$$

classifies the tangent bundle of S^k and the stable trivialization

$$\delta\tau_{S^k} : \tau_{S^k} \oplus \epsilon \cong \epsilon^{k+1}$$

determined by the embedding $S^k \hookrightarrow S^{k+1}$.

Lemma 6.72. *For any space M there is a natural one-one correspondence between the equivalence classes of pairs*

$$(k\text{-plane bundle } \xi \text{ over } M, \text{ stable isomorphism } \delta\xi : \xi \oplus \epsilon \cong \epsilon^{k+1})$$

and the homotopy classes of maps $\gamma : M \rightarrow S^k$.

Proof. A map $\gamma : M \rightarrow V_{k+1,1} = S^k$ classifies a k -plane bundle

$$\xi = \gamma^* \tau_{S^k} : M \rightarrow BO(k)$$

together with a stable isomorphism

$$\delta\xi = \gamma^* \delta\tau_{S^k} : \xi \oplus \epsilon \cong \epsilon^{k+1}.$$

Conversely, given $(\xi, \delta\xi)$ the composite of the non-zero section

$$M \rightarrow E(\nu_f \oplus \epsilon) \setminus M ; x \mapsto (x, (0, 1))$$

and the map

$$E(\xi \oplus \epsilon) \setminus M \xrightarrow[\cong]{\delta\xi} E(\epsilon^{k+1}) \setminus M = M \times (\mathbb{R}^{k+1} \setminus \{0\}) \rightarrow \mathbb{R}^{k+1} \setminus \{0\} \simeq S^k$$

is a map $\gamma : M \rightarrow S^k$ classifying ξ and $\delta\xi$.

□

The Hopf invariant of

$$T(\delta\xi) : T(\xi \oplus \epsilon) = \Sigma X^\infty \rightarrow T(\epsilon^{k+1}) = \Sigma^{k+1} M^+$$

can be regarded (as in 5.58) as an inequivariant map

$$h_{\mathbb{R}}(T(\delta\xi)) : \Sigma^2 X^\infty \rightarrow \Sigma T(\epsilon^k) \wedge \Sigma T(\epsilon^k),$$

which has the following geometric interpretation for a manifold M .

Definition 6.73. Let M be an m -dimensional manifold, together with a k -plane bundle $\xi : M \rightarrow BO(k)$ and a stable trivialization $\delta\xi : \xi \oplus \epsilon \cong \epsilon^{k+1}$. A *linking manifold* for $(\xi, \delta\xi)$ is a framed codimension k submanifold

$$L(M, \xi, \delta\xi)^{m-k} = \gamma^{-1}(*) \subset M$$

with $\gamma : M \rightarrow S^k$ a classifying map for $(\xi, \delta\xi)$, and $*$ $\in S^k$ a regular value of γ . \square

The homotopy class $\gamma \in [M, S^k]$ may be identified with the framed bordism class of $L(M, \xi, \delta\xi)^{m-k} \subset M$.

Proposition 6.74. (i) *A linking manifold for $(\xi, \delta\xi)$ is the linking manifold*

$$L(M, \xi, \delta\xi) = L(M_1, M_2, X)$$

for the disjoint embeddings

$$\begin{aligned} e_1 \times f_1 : M_1 &= M \hookrightarrow \mathbb{R} \times X ; x \mapsto (0, (0, x)) , \\ e_2 \times f_2 : M_2 &= M \hookrightarrow \mathbb{R} \times E(\xi) ; x \mapsto \delta\xi^{-1}(*, x) , \end{aligned}$$

with

$$\delta\xi^{-1} : E(\epsilon^{k+1}) = \mathbb{R}^{k+1} \times M \rightarrow E(\xi \oplus \epsilon) = \mathbb{R} \times E(\xi) .$$

(ii) *The bordism class of the immersion of the linking manifold $L(M, \xi, \delta\xi) \looparrowright E(\xi)$ is the stable homotopy class of the Hopf invariant map*

$$h_{\mathbb{R}}(T(\delta\xi)) : \Sigma^2 X^\infty \rightarrow \Sigma^{k+1} M^+ \wedge \Sigma^{k+1} M^+$$

with

$$T(\delta\xi) : T(\xi \oplus \epsilon) = \Sigma T(\xi) \rightarrow T(\epsilon^{k+1}) = \Sigma^{k+1} M^+ .$$

(iii) *The Hopf invariant of $T(\delta\xi)$ is given by*

$$\begin{aligned} h_{\mathbb{R}}(T(\delta\xi)) : \Sigma^2 T(\xi) \cong \Sigma^{k+2} M^+ &\xrightarrow{\Sigma^{k+2} \ell} \Sigma^{2k+2} L(M, \xi, \delta\xi)^\infty \\ &\xrightarrow{\Sigma^{2k+2} i} \Sigma^{2k+2} (M^+ \wedge M^+) \end{aligned}$$

with $\ell : M^+ \rightarrow \Sigma^k L(M, \xi, \delta\xi)^\infty$ a Pontryagin-Thom map for a linking manifold $L(M, \xi, \delta\xi) \subset M$, and

$$i : L(M, \xi, \delta\xi) \hookrightarrow M \times M ; x \mapsto (x, x) .$$

\square

Proposition 6.75. *Let $(e \times f : M^m \hookrightarrow \mathbb{R} \times N^n, f : M \looparrowright N)$ be an $(m, n, 1)$ -dimensional embedding-immersion pair with $e \times f$ framed, so that $\nu_f : M \rightarrow BO(n-m)$ has a stable trivialization*

$$\delta\nu_f : \nu_f \oplus \epsilon \cong \epsilon^{n-m+1} .$$

The Hopf invariant of the composite map

$$\phi : \Sigma N^+ \xrightarrow{F} \Sigma T(\nu_f) \xrightarrow{T(\delta\nu_f)} T(\epsilon^{n-m+1})$$

is the sum

$$h_{\mathbb{R}}(\phi) = (T(\delta\nu_f) \wedge T(\delta\nu_f))h_{\mathbb{R}}(F) + h_{\mathbb{R}}(T(\delta\nu_f))(\Sigma F) : \\ \Sigma^2 N^+ \rightarrow T(\epsilon^{n-m+1}) \wedge T(\epsilon^{n-m+1})$$

with

$$h_{\mathbb{R}}(F) : \Sigma^2 N^+ \rightarrow \Sigma T(\nu_f) \wedge \Sigma T(\nu_f)$$

the Pontryagin-Thom map for the self-linking manifold $L(M, N) \hookrightarrow \mathbb{R} \times \mathbb{R} \times N$ and

$$h_{\mathbb{R}}(T(\delta\nu_f)) : \Sigma^2 T(\nu_f) \cong \Sigma^{n-m+2} M^+ \\ \xrightarrow{\Sigma^{n-m+2}\ell} \Sigma^{2n-2m+2} L(M, \nu_f, \delta\nu_f)^\infty \xrightarrow{\Sigma^{2n-2m+2}i} \Sigma^{2n-2m+2} (M^+ \wedge M^+)$$

with $\ell : M^+ \rightarrow \Sigma^{n-m} L(M, \nu_f, \delta\nu_f)^+$ the Pontryagin-Thom map for the linking manifold $L(M, \nu_f, \delta\nu_f) \subset M$ and

$$i : L(M, \nu_f, \delta\nu_f) \hookrightarrow M \times M ; x \mapsto (x, x) .$$

□

Example 6.76. (i) Given an n -dimensional manifold N and a map

$$\phi : (\mathbb{R} \times N)^\infty = \Sigma N^+ \rightarrow S^{k+1}$$

apply the Pontryagin-Thom construction to obtain an $(n-k, n, 1)$ -dimensional embedding-immersion pair

$$(e \times f : M^{n-k} = \phi^{-1}(*) \hookrightarrow \mathbb{R} \times N , f : M \looparrowright N)$$

with $e \times f$ framed, so that the normal bundle $\nu_f : M \rightarrow BO(k)$ has a stable trivialization

$$\delta\nu_f : \nu_f \oplus \epsilon \cong \epsilon^{k+1} .$$

If $F : \Sigma N^+ \rightarrow \Sigma T(\nu_f)$ is the Umkehr stable map then

$$\phi : \Sigma N^+ \xrightarrow{F} \Sigma T(\nu_f) \cong \Sigma^{k+1} M^+ \rightarrow S^{k+1} .$$

By 6.54 the Hopf invariant map

$$h_{\mathbb{R}}(\phi) : \Sigma^2 N^+ \rightarrow S^{2k+2}$$

is such that

$$h_{\mathbb{R}}(\phi)^{-1}(\ast)^{n-k} = L(M, N) \cup L(M, \nu_f, \delta\nu_f) \subset \mathbb{R} \times \mathbb{R} \times N .$$

(ii) Setting $N = S^n$ in (i) shows that the Hopf map (6.68)

$$H : \pi_{n+1}(S^{k+1}) \rightarrow \pi_{n+2}(S^{2k+2}) ; \phi \mapsto h_{\mathbb{R}}(\phi)$$

sends the bordism class of the $(n-k, n, 1)$ -dimensional embedding-immersion pair

$$(e \times f : M^{n-k} = \phi^{-1}(\ast) \hookrightarrow \mathbb{R} \times S^n , f : M \looparrowright S^n)$$

with $e \times f$ framed to the bordism class of the framed submanifold

$$L(M, S^n)^{n-2k} \cup L(M, \nu_f, \delta\nu_f)^{n-2k} \subset S^{n+2} .$$

□

Example 6.77. The homotopy group $\pi_{m+k}(T(\tau_{S^k}))$ is the cobordism group of embeddings $f : M^m \hookrightarrow S^{m+k}$ which are framed in S^{m+k+1} . A map $\rho : S^{m+k} \rightarrow T(\tau_{S^k})$ which is transverse at the zero section $S^k \subset T(\tau_{S^k})$ determines an embedding

$$f : M^m = \rho^{-1}(S^k) \hookrightarrow S^{m+k}$$

which is framed in S^{m+k+1} , with a map $\gamma = \rho| : M \rightarrow S^k$ such that

$$\nu_f = \gamma^* \tau_{S^k} : M \rightarrow BO(k) ,$$

with a stable trivialization

$$\delta\nu_f = \gamma^* \delta\tau_{S^k} : \nu_f \oplus \epsilon \cong \epsilon^{k+1} .$$

The linking manifold construction (6.73) gives rise to the commutative square of Wood [76]

$$\begin{array}{ccc} \pi_{m+k}(T(\tau_{S^k})) & \xrightarrow{h} & \pi_{m+k}(S^{2k}) \\ \downarrow & & \downarrow E \\ \pi_{m+k+1}(S^{k+1}) & \xrightarrow{H} & \pi_{m+k+1}(S^{2k+1}) \end{array}$$

with

$$\begin{aligned} h : \pi_{m+k}(T(\tau_{S^k})) &\rightarrow \pi_{m+k}(S^{2k}) ; \rho \mapsto (L(M, \nu_f, \delta\nu_f)^{m-k} \subset S^{m+k}) , \\ \pi_{m+k}(T(\tau_{S^k})) &\rightarrow \pi_{m+k+1}(S^{k+1}) ; \rho \mapsto (M \subset S^{m+k+1}) . \end{aligned}$$

The Thom space of $\tau_{S^k} : S^k \rightarrow BO(k)$ is

$$T(\tau_{S^k}) = S^k \cup_{J(\tau_{S^k})} e^{2k}$$

with $J(\tau_{S^k}) : S^{2k-1} \rightarrow S^k$ given by the J -homomorphism

$$J : \pi_k(BO(k)) = \pi_{k-1}(O(k)) \rightarrow \pi_{2k-1}(S^k) ,$$

and h is induced by $T(\tau_{S^k}) \rightarrow S^{2k}$. The Hopf map H is given by

$$\begin{aligned} H : \pi_{m+k+1}(S^{k+1}) &\rightarrow \pi_{m+k+1}(S^{2k+1}) ; \\ (M, e, f) &\mapsto (L(M, S^{m+k})^{m-k} \subset S^{m+k+1}) + (L(M, \nu_f, \delta \nu_f)^{m-k} \subset S^{m+k+1}) , \end{aligned}$$

regarding $\pi_{m+k+1}(S^{k+1})$ as the cobordism group of $(m, m+k, 1)$ -dimensional embedding-immersion pairs $(e \times f : M^m \hookrightarrow \mathbb{R} \times S^{m+k}, f : M \looparrowright S^{m+k})$ such that $e \times f$ is framed. \square

Example 6.78. Let $m \leq 2k - 2$, so that

$$\pi_{m+k+1}(S^{2k+1}) = \pi_m(S^k)$$

by the Freudenthal suspension theorem; equivalently, every (framed) submanifold $L^{m-k} \subset S^{m+k+1}$ can be compressed to $L^{m-k} \subset S^m$.

(i) Regard $\pi_{m+k+1}(S^{k+1})$ as the cobordism group of $(m, m+k, 1)$ -dimensional embedding-immersion pairs

$$(e \times f : M^m \hookrightarrow \mathbb{R} \times S^{m+k} , f : M \looparrowright S^{m+k})$$

with $e \times f$ framed and f an embedding, so that

$$(L(M, S^{m+k})^{m-k} \subset S^{m+k+1}) = 0 \in \pi_{m+k+1}(S^{2k+1})$$

and

$$\begin{aligned} H : \pi_{m+k+1}(S^{k+1}) &\rightarrow \pi_{m+k+1}(S^{2k+1}) ; \\ (M, e, f) &\mapsto (L(M, \nu_f, \delta \nu_f)^{m-k} \subset S^{m+k+1}) \end{aligned}$$

is the ‘singularity Hopf invariant’ of Koschorke and Sanderson [37, p. 201].

(ii) Regard $\pi_{m+k+1}(S^{k+1})$ as the cobordism group of $(m, m+k, 1)$ -dimensional embedding-immersion pairs

$$(e \times f : M^m \hookrightarrow \mathbb{R} \times S^{m+k} , f : M \looparrowright S^{m+k})$$

with a framing $\nu_f \cong \epsilon^k$, so that

$$(L(M, \nu_f, \delta \nu_f)^{m-k} \subset S^{m+k+1}) = 0 \in \pi_{m+k+1}(S^{2k+1})$$

and

$$H : \pi_{m+k+1}(S^{k+1}) \rightarrow \pi_{m+k+1}(S^{2k+1}) ;$$

$$(M, e, f) \mapsto (L(M, S^{m+k})^{m-k} \subset S^{m+k+1})$$

is the ‘double point Hopf invariant’ of [37, p. 202] (and 6.62 (ii)). \square

6.8 Intersections and self-intersections for $M^m \rightarrow N^{2m}$

We now consider the application of the geometric Hopf invariant to the intersection and self-intersection properties of embeddings and immersions $f : M \rightarrow N$ of an m -dimensional manifold M^m in a $2m$ -dimensional manifold N^{2m} . It will be assumed that M and N are oriented, so that the normal bundle is an oriented m -plane bundle $\nu_f : M \rightarrow BSO(m)$.

Definition 6.79. (i) The *intersection pairing* of a $2m$ -dimensional manifold N is the $(-1)^m$ -symmetric cup product pairing

$$\lambda : H_m(N) \times H_m(N) \rightarrow \mathbb{Z} ; (a, b) \mapsto \langle a^* \cup b^*, [N] \rangle$$

with $a^*, b^* \in H^m(N)$ the Poincaré duals of $a, b \in H_m(N)$.

(ii) Let $(x_1, x_2) \in M_1 \times M_2$ be a transverse intersection point of embeddings

$$f_1 : (M_1)^m \hookrightarrow N^{2m} , \quad f_2 : (M_2)^m \hookrightarrow N^{2m}$$

so that there is defined an isomorphism of $2m$ -dimensional vector spaces

$$\tau_{M_1}(x_1) \oplus \tau_{M_2}(x_2) \cong \tau_N(y)$$

with

$$y = f_1(x_1) = f_2(x_2) \in N .$$

Assuming M_1, M_2, N are oriented the *intersection index* of (x_1, x_2) is

$$I(x_1, x_2) = \begin{cases} +1 & \text{if } \tau_{M_1}(x_1) \oplus \tau_{M_2}(x_2) \cong \tau_N(y) \text{ is orientation-preserving} \\ -1 & \text{otherwise} \end{cases}$$

where $y = f_1(x_1) = f_2(x_2) \in N$. \square

Proposition 6.80. (i) The *intersection pairing* $\lambda : H_m(N) \times H_m(N) \rightarrow \mathbb{Z}$ of a $2m$ -dimensional manifold N is such that

$$\lambda(a, b) = (-1)^m \lambda(b, a) \in \mathbb{Z} .$$

(ii) If $f_1 : (M_1)^m \hookrightarrow N^{2m}$, $f_2 : (M_2)^m \hookrightarrow N^{2m}$ are transverse embeddings of oriented manifolds

$$\lambda(f_1[M_1], f_2[M_2]) = \sum_{f_1(x_1)=f_2(x_2)} I(x_1, x_2) \in \mathbb{Z} . \quad \square$$

The homotopy exact sequence of the fibration

$$V_{\infty, \infty-m} \rightarrow BO(m) \rightarrow BO$$

is mapped by the J -homomorphism to the stable EHP sequence of S^m

$$\begin{array}{ccccccc} \pi_{m+1}(BO) & \longrightarrow & \pi_m(V_{\infty, \infty-m}) & \longrightarrow & \pi_m(BO(m)) & \longrightarrow & \pi_m(BO) \\ \downarrow J & & \downarrow \cong & & \downarrow J & & \downarrow J \\ \pi_m^S & \xrightarrow{H} & Q_{(-1)^m}(\mathbb{Z}) & \longrightarrow & \pi_{2m-1}(S^m) & \longrightarrow & \pi_{m-1}^S \end{array}$$

The isomorphism

$$\pi_m(V_{\infty, \infty-m}) \xrightarrow{\cong} \Omega_0(P(\mathbb{R}(\infty)), m\lambda) = Q_{(-1)^m}(\mathbb{Z}) ; (\xi, \delta\xi) \mapsto D_2(S^m, \xi, \delta\xi)/\mathbb{Z}_2$$

is defined by the double point Hopf invariant (6.60), and

$$\pi_{m+1}(BO, BO(m)) = Q_{(-1)^m}(\mathbb{Z}) \rightarrow \pi_m(BO(m)) ; 1 \mapsto \tau_{S^m} .$$

The Euler number defines a morphism

$$\chi : \pi_m(BSO(m)) \rightarrow \mathbb{Z} ; \xi \mapsto \chi(\xi)$$

such that

$$\begin{aligned} \pi_{m+1}(BO, BO(m)) &= Q_{(-1)^m}(\mathbb{Z}) \longrightarrow \pi_m(BSO(m)) \xrightarrow{\chi} \mathbb{Z} ; \\ 1 &\mapsto \chi(\tau_{S^m}) = 1 + (-1)^m . \end{aligned}$$

Proposition 6.81. For any embedding $f : M^m \hookrightarrow N^{2m}$

$$\lambda(f_*[M^m], f_*[M^m]) = \chi(\nu_f) \in \mathbb{Z} .$$

Proof. There exists an isotopic embedding $f' : M^m \hookrightarrow N^{2m}$ which intersects transversely with f in $\chi(\nu_f)$ points (counted algebraically), and

$$\lambda(f_*[M], f_*[M]) = \sum_{(x, x') \in D_2(f, f')} I(x, x') = \chi(\nu_f) \in \mathbb{Z}.$$

□

Example 6.82. If N^{4k} is a $(2k-1)$ -connected $4k$ -dimensional manifold and $k \geq 2$ then every element $x \in H_{2k}(N)$ is represented by an embedding $f : S^{2k} \hookrightarrow N$, with

$$\lambda(x, x) = \chi(\nu_f) \in \mathbb{Z}$$

as in Wall [67].

□

Recall the definition of the double point set (6.15) of a map $f : M \rightarrow N$

$$D_2(f) = \{(x, y) \in M \times M \mid x \neq y \in M, f(x) = f(y) \in N\}.$$

Transposition defines a free \mathbb{Z}_2 -action

$$T : D_2(f) \rightarrow D_2(f) ; (x, y) \mapsto (y, x).$$

For any ordered double point $(x, y) \in D_2(f)$ let $[x, y] \in D_2(f)/\mathbb{Z}_2$ be the unordered double point.

Definition 6.83. Let $f : M^m \looparrowright N^{2m}$ be an immersion, so that the double point set $D_2(f)$ is a finite 0-dimensional manifold with a free \mathbb{Z}_2 -action.

(i) The *self-intersection index* of an unordered double point $[x, y] \in D_2(f)/\mathbb{Z}_2$ is

$$I[x, y] = [I(x, y)] \in \mathbb{Z}/\{1 + (-1)^{m+1}\} = \begin{cases} \mathbb{Z} & \text{if } m \equiv 0 \pmod{2} \\ \mathbb{Z}_2 & \text{if } m \equiv 1 \pmod{2} \end{cases}$$

which does not depend on the choice of lift of $[x, y]$ to an ordered double point $(x, y) \in D_2(f)$, since $I(x, y) = (-1)^m I(y, x) \in \mathbb{Z}$.

(ii) The *geometric self-intersection number* of f is

$$\mu(f) = \sum_{[x, y] \in D_2(f)/\mathbb{Z}_2} I[x, y] \in Q_{(-)^m}(\mathbb{Z}) = \mathbb{Z}/\{1 + (-1)^{m+1}\}. \quad \square$$

Proposition 6.84. (Whitney [74], Wall [68, 5.3]) *The geometric self-intersection number $\mu(f)$ is a regular homotopy invariant of $f : M^m \looparrowright N^{2m}$ such that*

$$\lambda(f, f) = (1 + (-1)^m)\mu(f) + \chi(\nu_f) \in \mathbb{Z}$$

with $\mu(f) = 0 \in Q_{(-)^m}(\mathbb{Z})$ if (and for $\pi_1(N) = \{1\}$, $m \geq 2$ only if) f is regular homotopic to an embedding.

□

In particular, for even m

$$\mu(f) = (\lambda(f, f) - \chi(\nu_f))/2 \in \mathbb{Z}.$$

Definition 6.85. Let $(e \times f : M^m \hookrightarrow V \times N^{2m}, f : M \looparrowright N)$ be an $(m, 2m, j)$ -dimensional embedding-immersion pair, with Umkehr stable map

$$F : (V \times N)^\infty = V^\infty \wedge N^+ \rightarrow V^\infty \wedge T(\nu_f).$$

(i) The *homological self-intersection number*

$$\mu_V(f) = \psi_V(F)_*[N] \in \begin{cases} \mathbb{Z} & \text{if } j = 1 \\ \mathbb{Z}/\{1 + (-1)^{m+1}\} & \text{if } \dim V \geq 2 \end{cases}$$

is the image of the fundamental class $[N] \in H_{2m}(N)$ under the induced map

$$\begin{aligned} \psi_V(F)_* : H_{2m}(N) &\rightarrow \tilde{H}_{2m}(S(LV)^\infty \wedge_{\mathbb{Z}_2} (T(\nu_f) \wedge T(\nu_f))) \\ &= H_0(S(LV) \times_{\mathbb{Z}_2} (M \times M); \mathbb{Z}^w) \\ &= \begin{cases} \mathbb{Z} & \text{if } j = 1 \\ \mathbb{Z}/\{1 + (-1)^{m+1}\} & \text{if } j \geq 2. \end{cases} \end{aligned}$$

(ii) If $j = 1$ set $V = \mathbb{R}$, and use the self-linking manifold (6.69)

$$L(M, N) = \{(x, y) \in M \times M \mid e(x) < e(y) \in \mathbb{R}, f(x) = f(y) \in N\}$$

and the decomposition

$$D_2(f) = L(M, N) \cup TL(M, N)$$

to lift each unordered double point $(x, y) \in D_2(f)/\mathbb{Z}_2$ to an ordered double point $(x, y) \in L(M, N)$, ordered according to $e(x) < e(y)$. The *integral geometric self-intersection* of f is the self-linking number

$$\mu^{\mathbb{Z}}(f) = [L(M, N)] = \sum_{(x, y) \in L(M, N)} I(x, y) \in \mathbb{Z}.$$

For even m $\mu^{\mathbb{Z}}(f) = \mu(f)$. □

Proposition 6.86. (i) *If m is even or if $\dim(V) \geq 2$ the geometric self-intersection number is just the homological self-intersection number*

$$\mu(f) = \mu_V(f) \in Q_{(-1)^m}(\mathbb{Z}) = \mathbb{Z}/\{1 + (-1)^{m+1}\}.$$

(ii) If $\dim(V) = 1$ the integral geometric self-intersection number is just the homological self-intersection number

$$\mu^{\mathbb{Z}}(f) = \mu_{\mathbb{R}}(f) \in \mathbb{Z} .$$

Proof. Immediate from 6.54 and 6.61. \square

Example 6.87. We refer to Crabb and Ranicki [14] for an interpretation in terms of the geometric Hopf invariant of the Smale-Hirsch-Haefliger regular homotopy classification of immersions $f : M^m \looparrowright N^n$ in the metastable dimension range $3m < 2n - 1$ (when a generic f has no triple points). In particular, this applies to the case $n = 2m$ with $m \geq 2$. \square

6.9 The simply-connected surgery obstruction

We shall now apply the geometric Hopf invariant to construct the quadratic form on the homology kernel of a normal map $(g, c) : N^n \rightarrow X$ from an n -dimensional manifold N to an n -dimensional Poincaré complex X , i.e. a degree 1 map $g : N \rightarrow X$ with a bundle map $c : \nu_N \rightarrow \eta$ to a vector bundle η over X .

An n -dimensional geometric Poincaré complex M (e.g. a manifold) has a Spivak normal structure

$$(\nu_M : M \rightarrow BG(k), \rho_M : S^{n+k} \rightarrow T(\nu_M))$$

with

$$\nu_M : S^{k-1} \rightarrow \partial W \rightarrow W \simeq M$$

the spherical fibration defined by a regular neighbourhood $(W, \partial W)$ of an embedding $M \subset S^{n+k}$, and

$$\rho_M : S^{n+k} \rightarrow S^{n+k}/(S^{n+k} \setminus W) = W/\partial W = T(\nu_M) .$$

The composite of the Hurewicz map and the Thom isomorphism

$$\pi_{n+k}(T(\nu_M)) \rightarrow \tilde{H}_{n+k}(T(\nu_M)) \cong H_n(M)$$

sends ρ_M to the fundamental class $[M] \in H_n(M)$ (using twisted coefficients in the nonoriented case). The composite

$$\sigma_M = \Delta\rho_M : S^{n+k} \xrightarrow{\rho_M} T(\nu_M) \xrightarrow{\Delta} M^+ \wedge T(\nu_M)$$

is an S -duality map, with S -duality isomorphisms

$$\begin{aligned} \sigma_M : \{T(\nu_M); X\} &\xrightarrow{\cong} \{S^{n+k}; M^+ \wedge X\} ; F \mapsto F\sigma_M , \\ \sigma_M : \{M^+; X\} &\xrightarrow{\cong} \{S^{n+k}; X \wedge T(\nu_M)\} ; G \mapsto G\sigma_M . \end{aligned}$$

For any n -dimensional geometric Poincaré complexes M, N there is thus defined an S -duality isomorphism

$$\{T(\nu_M); T(\nu_N)\} \cong \{N^+; M^+\} .$$

As in Ranicki [50] given a degree 1 normal map $(f, b) : M \rightarrow N$ call any S -dual of $T(b) : T(\nu_M) \rightarrow T(\nu_N)$ a *geometric Umkehr map* for (f, b)

$$F = T(b)^* : V^\infty \wedge N^+ \rightarrow V^\infty \wedge M^+$$

inducing the Umkehr chain map

$$f^! : C(X) \simeq C(X)^{n-*} \xrightarrow{f^*} C(N)^{n-*} \simeq C(N) ,$$

such that

$$\begin{aligned} (1_V \wedge f)F &\simeq 1 : V^\infty \wedge N^+ \rightarrow V^\infty \wedge M^+ , \\ ff^! &\simeq 1 : C(N) \rightarrow C(N) . \end{aligned}$$

The homology and cohomology groups split as

$$H_*(N) = H_*(X) \oplus K_*(N) , \quad H^*(N) = H^*(X) \oplus K^*(N)$$

with

$$\begin{aligned} K_m(N) &= \ker(g_* : H_m(N) \rightarrow H_m(X)) , \\ K^m(N) &= \ker(g^! : H^m(N) \rightarrow H^m(X)) . \end{aligned}$$

The Poincaré duality isomorphisms $H^*(N) \cong H_{n-*}(N)$ restrict to Poincaré duality isomorphisms

$$K^*(N) \cong K_{n-*}(N) ,$$

If M is an n -dimensional manifold and the n -dimensional geometric Poincaré complex N is replaced by an $(n+k)$ -dimensional manifold with boundary $(X, \partial X)$ homotopy equivalent to $(N \times D^k, N \times S^{k-1})$ (for $k \geq \max(3, 5-n, n+1)$, applying the π - π Theorem of Wall [68, 3.3]¹ and the Whitney embedding theorem) it is possible to approximate (f, b) by a framed embedding $e : V \times M \hookrightarrow X \setminus \partial X$ with V a k -dimensional inner product space,

¹ which applies because the Spivak normal fibration of N has a vector bundle reduction

and the adjunction Umkehr map of e is a geometric Umkehr map for (f, b)

$$F : X/\partial X \simeq V^\infty \wedge N^+ \rightarrow X/(X - e(V \times M)) = V^\infty \wedge M^+ .$$

and there is defined an intersection pairing

$$\lambda : K_m(N) \times K_{n-m}(N) \rightarrow \mathbb{Z}$$

such that

$$\begin{aligned} \lambda(x, y) &= (-)^{m(n-m)} \lambda(y, x) \in \mathbb{Z} \quad (x \in K_m(N), y \in K_{n-m}(N)) , \\ \lambda([N] \cap a, [N] \cap b) &= \langle a \cup b, [N] \rangle \in \mathbb{Z} \quad (a \in K^{n-m}(N), b \in K^m(N)) . \end{aligned}$$

Let $(g, c) : N^{2m} \rightarrow X$ be a $2m$ -dimensional normal map, with $\eta : X \rightarrow BO(k)$ a k -plane bundle, for some $k \gg 1$. The bundle map $c : \nu_N \rightarrow \eta$ determines a bundle isomorphism

$$\tau_N \oplus \eta \cong \epsilon^{2m+k} .$$

The normal bundle of an immersion $f : S^m \looparrowright N$ is an m -plane bundle $\nu_f : S^m \rightarrow BO(m)$ such that

$$\nu_f \oplus \tau_{S^m} = f^* \tau_N : S^m \rightarrow BO(2m) ,$$

so that

$$\nu_f \oplus \tau_{S^m} \oplus (gf)^* \eta = f^* (\tau_N \oplus \eta) \cong \epsilon^{2m+k} .$$

A null-homotopy $gf \simeq * : S^m \rightarrow X$ thus determines a stable isomorphism

$$\delta \nu_f : \nu_f \oplus \epsilon^{m+k} \cong \epsilon^{2m+k} ,$$

as classified by an element

$$(\delta \nu_f, \nu_f) \in \pi_m(V_{2m+k, m+k}) = Q_{(-1)^m}(\mathbb{Z}) .$$

Proposition 6.88. *Let $(g, c) : N \rightarrow X$ be a $2m$ -dimensional normal map, with Umkehr stable map $G : \Sigma^j X^\infty \rightarrow \Sigma^j N^+$.*

(i) *The geometric Hopf invariant map*

$$h_{\mathbb{R}^j}(G) : X^\infty \looparrowright S(L\mathbb{R}^j)^\infty \wedge_{\mathbb{Z}_2} (N^+ \wedge N^+) = (S(L\mathbb{R}^j) \times_{\mathbb{Z}_2} (N \times N))^\infty$$

determines a quadratic refinement of the intersection pairing $\lambda : K_m(N) \times K_m(N) \rightarrow \mathbb{Z}$

$$\begin{aligned} \mu : K_m(N) &\rightarrow Q_{(-1)^m}(\mathbb{Z}) = \mathbb{Z}/\{1 + (-1)^{m+1}\} ; \\ x &\mapsto -(x \otimes x)h_{\mathbb{R}^j}(G)_*[N] = \begin{cases} \lambda(x, x)/2 & \text{if } m \equiv 0 \pmod{2} \\ \langle Sq_{xG}^{m+1}(\iota), [X] \rangle & \text{if } m \equiv 1 \pmod{2} \end{cases} \end{aligned}$$

such that

- (a) $\lambda(x, x) = (1 + (-1)^m)\mu(x) \in \mathbb{Z}$,
- (b) $\mu(ax) = a^2\mu(x) \in Q_{(-1)^m}(\mathbb{Z})$ ($a \in \mathbb{Z}$),
- (c) $\mu(x + y) = \mu(x) + \mu(y) + \lambda(x, y) \in Q_{(-1)^m}(\mathbb{Z})$,
- (d) if $x \in K_m(N)$ is represented by an immersion $f : S^m \looparrowright N$ with a null-homotopy $gf \simeq * : S^m \rightarrow X$ then

$$\begin{aligned} \lambda(x, x) &= (1 + (-1)^m)\mu(f) + \chi(\nu_f) \in \mathbb{Z} , \\ \mu(x) &= H(\phi) = \mu(f) + (\nu_f, \delta\nu_f) \in Q_{(-1)^m}(\mathbb{Z}) \text{ with} \\ H(\phi) &= h_{\mathbb{R}^k}(\phi)[N] \in \tilde{H}_{2m}((S^{k-1})^\infty \wedge_{\mathbb{Z}_2} (S^m \wedge S^m)) = Q_{(-1)^m}(\mathbb{Z}) \\ &\text{the Hopf invariant of the stable map} \end{aligned}$$

$$\phi : \Sigma^k N^+ \xrightarrow{F} \Sigma^k T(\nu_f) \xrightarrow{T(\delta\nu_f)} \Sigma^k T(\epsilon^m) \longrightarrow S^{m+k}$$

where F is the Umkehr stable map of f .

(ii) If g is m -connected then $K_m(N) = \pi_{m+1}(g)$ and every element $x \in K_m(N)$ is represented by an immersion $f : S^m \looparrowright N$ with a null-homotopy $gf \simeq * : S^m \rightarrow X$. There are two special cases:

- (a) For all m it is possible to choose a representative with $\nu_f = \epsilon^m$ and $(\nu_f, \delta\nu_f) = 0$.
- (b) If $m \geq 3$ and $\pi_1(N) = \{1\}$ it is possible to choose a representative with f an embedding, so that $\mu(f) = 0$.

(iii) If g is m -connected then $\mu(x) = 0$ if (and for $m \geq 3$, $\pi_1(N) = \{1\}$ only if) it is possible to kill $x \in K_m(N)$ by surgery on (g, c) , i.e. to represent x by a framed embedding $f : S^m \hookrightarrow N$ with a null-homotopy $gf \simeq * : S^m \rightarrow X$.

(iv) The simply-connected surgery obstruction of Browder [7]

$$\begin{aligned} \sigma_*(g, c) &= \begin{cases} \text{signature}(K_m(N), \lambda)/8 \\ \text{Arf}(K_m(N; \mathbb{Z}_2), \lambda, \mu) \end{cases} \\ &\in L_{2m}(\mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } m \equiv 0 \pmod{2} \\ \mathbb{Z}_2 & \text{if } m \equiv 1 \pmod{2} \end{cases} \end{aligned}$$

is such that $\sigma_*(g, c) = 0$ if (and for $\pi_1(X) = \{1\}$, $m \geq 3$ only if) (g, c) is normal bordant to a homotopy equivalence.

Remark 6.89. The quadratic refinement μ was constructed by Browder [7] using functional Steenrod squares, by Wall [68] using immersions (assuming (g, c) is m -connected), and by Ranicki [50] using the quadratic construction ψ_G . \square

We shall now consider the 1-dimensional case $V = \mathbb{R}$, such as arises in codimension 2 surgery.

The homotopy exact sequence of the fibration

$$S^m \rightarrow BO(m) \rightarrow BO(m+1)$$

is mapped by the J -homomorphism to the EHP sequence of S^m

$$\begin{array}{ccccccc} \pi_{m+1}(BO(m+1)) & \longrightarrow & \pi_m(S^m) & \longrightarrow & \pi_m(BO(m)) & \longrightarrow & \pi_m(BO(m+1)) \\ \downarrow J & & \downarrow \cong & & \downarrow J & & \downarrow J \\ \pi_{2m+1}(S^{m+1}) & \xrightarrow{H} & \mathbb{Z} & \xrightarrow{P} & \pi_{2m-1}(S^m) & \xrightarrow{E} & \pi_{2m}(S^{m+1}) \end{array}$$

where

$$HJ = \chi : \pi_{m+1}(BO(m+1)) \rightarrow \mathbb{Z}$$

sends an $(m+1)$ -plane bundle over S^{m+1} to its Euler number (only determined up to sign in general). The image of the Hopf invariant is

$$\text{im}(H : \pi_{2m+1}(S^{m+1}) \rightarrow \mathbb{Z}) = \begin{cases} 0 & \text{if } m \text{ is even} \\ \mathbb{Z} & \text{if } m = 1, 3, 7 \\ 2\mathbb{Z} & \text{otherwise (Adams [1])} \end{cases}.$$

The morphism

$$\pi_{m+1}(BO(m+1), BO(m)) = \mathbb{Z} \rightarrow \pi_m(BO(m)) ; 1 \mapsto \tau_{S^m}$$

is $\begin{cases} \text{injective} \\ 0 \end{cases}$ for $\begin{cases} m \neq 1, 3, 7 \\ m = 1, 3, 7 \end{cases}$ with the generator

$$(\delta\tau_{S^m}, \tau_{S^m}) = 1 \in \pi_{m+1}(BO(m+1), BO(m)) = \mathbb{Z}$$

represented by the $\begin{cases} \text{unique} \\ \text{exotic} \end{cases}$ stable trivialization $\delta\tau_{S^m} : \tau_{S^m} \oplus \epsilon \cong \epsilon^{m+1}$.

Definition 6.90. (Ranicki [51, p.818]) An n -dimensional normal map $(g, c) : N \rightarrow X$ is *ultranormal* if it is obtained from an $(n+1)$ -dimensional manifold W and a $\mathbb{Z}[\pi_1(X)]$ -homology equivalence $h : W \rightarrow X \times S^1$ by codimension 1 transversality

$$(g, c) = h| : N = h^{-1}(X \times \{*\}) \rightarrow X. \quad \square$$

Example 6.91. ([51, p.827]) Let k be a $(2m-1)$ -knot, i.e. a (locally flat) embedding $k : S^{2m-1} \hookrightarrow S^{2m+1}$. The knot complement

$$W = \text{cl.}(S^{2m+1} \setminus (k(S^{2m-1}) \times D^2))$$

is a $(2m+1)$ -dimensional manifold with boundary $\partial W = S^{2m-1} \times S^1$. Let $N^{2m} \subset S^{2m+1}$ be a Seifert surface k , with $\partial N = k(S^m)$. The inclusion $g : N \hookrightarrow D^{2m+2}$ defines an ultranormal map

$$(g, c) = h| : (N, \partial N) = h^{-1}((D^{2m+2}, k(S^{2m-1})) \times \{*\}) \rightarrow (D^{2m+2}, k(S^{2m-1}))$$

for a homology equivalence

$$h : (W, \partial W) \rightarrow (D^{2m+2}, k(S^{2m-1})) \times S^1$$

with both g and h identities on boundaries. \square

Remark 6.92. Every m -connected $2m$ -dimensional normal map can be realized as an ultranormal map ([51, 7.8]). \square

An ultranormal map $(g, c) : N \rightarrow X$ has an Umkehr stable map involving only a single suspension

$$G : \Sigma X^\infty = (X \times \mathbb{R})^\infty \xrightarrow{h^{-1}} \overline{W}^\infty \longrightarrow \Sigma N^+$$

with $\overline{W} = h^*(X \times \mathbb{R})$ the infinite cyclic cover of W , and $\overline{W}^\infty \rightarrow \Sigma N^+$ the Pontryagin-Thom map of an embedding $N \times \mathbb{R} \hookrightarrow \overline{W}$. We shall now relate the geometric Hopf map of G (5.58)

$$h_{\mathbb{R}}(G) : X^{\infty} \rightarrow N^+ \wedge N^+$$

in the case $\dim(N) = 2m$ to the Seifert form on

$$K_m(N) = \ker(g_* : H_m(N) \rightarrow H_m(X))$$

which was originally defined using linking numbers. In the first instance, we consider the (self-) intersection properties of $(m, 2m, 1)$ -dimensional embedding-immersion pairs $(e \times f : M^m \hookrightarrow \mathbb{R} \times N^{2m}, f : M \looparrowright N)$.

Definition 6.93. The *degree* of an m -plane bundle $\xi : M \rightarrow BO(m)$ and a stable isomorphism $\delta\xi : \xi \oplus \epsilon \cong \epsilon^{m+1}$ over an m -dimensional manifold M is

$$(\xi, \delta\xi)^{\mathbb{Z}} = \text{degree}(\gamma : M \rightarrow S^m) \in \mathbb{Z}$$

with $\gamma : M \rightarrow S^m$ a classifying map for $(\xi, \delta\xi)$ (6.72). \square

Proposition 6.94. (i) *The degree of $(\xi, \delta\xi)$ determines the Euler number of ξ by*

$$\chi(\xi) = (1 + (-1)^m)(\xi, \delta\xi)^{\mathbb{Z}} \in \mathbb{Z}.$$

(ii) *For any m -dimensional manifold M the degree defines a morphism*

$$[M, S^m] \rightarrow \mathbb{Z} ; \gamma \mapsto \text{degree}(\gamma : M \rightarrow S^m).$$

For connected M this is an isomorphism (by the Hopf-Whitney theorem) and the degree classifies pairs $(\xi, \delta\xi)$ as in 6.93.

(iii) *The degree is the linking number of the submanifolds*

$$\begin{aligned} i_0 : M &\hookrightarrow E(\epsilon^{m+1}) = M \times \mathbb{R}^{m+1} ; x \mapsto (x, 0), \\ i_1 : M &\hookrightarrow E(\epsilon^{m+1}) = M \times \mathbb{R}^{m+1} ; x \mapsto E(\delta\xi)(x, 0, 1), \end{aligned}$$

which is the degree of the composite

$$\gamma : M \xrightarrow{i_1} (M \times \mathbb{R}^{m+1}) \setminus i_0(M) = M \times (\mathbb{R}^{m+1} \setminus \{0\}) \longrightarrow \mathbb{R}^{m+1} \setminus \{0\} \simeq S^m,$$

or equivalently the algebraic number of elements in the 0-dimensional linking manifold $L(M, \xi, \delta\xi) = \gamma^{-1}()$ (6.73).* \square

Proposition 6.95. *Let $(e \times f : M^m \hookrightarrow \mathbb{R} \times N^{2m}, f : M \looparrowright N)$ be an $(m, 2m, 1)$ -dimensional embedding-immersion pair with $e \times f$ framed, so that $\nu_f : M \rightarrow BO(m)$ has a stable trivialization*

$$\delta\nu_f : \nu_f \oplus \epsilon \cong \epsilon^{m+1}.$$

Let $F : \Sigma N^+ \rightarrow \Sigma T(\nu_f)$ be an Umkehr stable map for f , so that the homotopy class of the Pontryagin-Thom map

$$\phi : \Sigma N^+ \xrightarrow{F} \Sigma T(\nu_f) \xrightarrow{T(\delta\nu_f)} T(\epsilon^{m+1}) \rightarrow S^{m+1}$$

is the framed cobordism class of

$$\phi^{-1}(*) = M \subset \mathbb{R} \times N .$$

(i) The Hopf invariant map $h_{\mathbb{R}}(\phi) : \Sigma^2 N^+ \rightarrow S^{2m+2}$ is the Pontryagin-Thom map for the framed 0-dimensional submanifold

$$h_{\mathbb{R}}(\phi)^{-1}(*) = L(M, N) \cup L(M, \nu_f, \delta\nu_f) \subset \mathbb{R} \times \mathbb{R} \times N ,$$

with degree

$$H(\phi) = \text{degree}(h_{\mathbb{R}}(\phi)) = \mu^{\mathbb{Z}}(f) + (\delta\nu_f, \nu_f)^{\mathbb{Z}} \in \mathbb{Z} .$$

(ii) If $f : M \looparrowright N$ is framed, and $\delta\nu_f$ is the stabilization of the framing $\nu_f \cong \epsilon^m$, then $L(M, \nu_f, \delta\nu_f) = \emptyset$ and

$$(\nu_f, \delta\nu_f)^{\mathbb{Z}} = 0 , \quad H(\phi) = \mu^{\mathbb{Z}}(f) \in \mathbb{Z} .$$

(iii) If $f : M \looparrowright N$ is an embedding then $L(M, N) = \emptyset$ and

$$\mu^{\mathbb{Z}}(f) = 0 , \quad H(\phi) = (\nu_f, \delta\nu_f)^{\mathbb{Z}} \in \mathbb{Z} .$$

Proof. (i) This is the special case $n = 2m$ of 6.75.

(ii)+(iii) Immediate from (i). □

Example 6.96. (i) For every immersion $f : S^m \looparrowright S^{2m}$ there exists a map $e : S^m \rightarrow \mathbb{R}$ such that $e \times f : S^m \rightarrow \mathbb{R} \times S^{2m}$ is a framed embedding, with the stable trivialization

$$\delta\nu_f : \nu_f \oplus \epsilon \cong \epsilon^{m+1}$$

such that the Pontryagin-Thom map

$$\phi : \Sigma(S^{2m})^{\infty} \xrightarrow{F} \Sigma T(\nu_f) \xrightarrow{T(\delta\nu_f)} T(\epsilon^{m+1}) \longrightarrow S^{m+1}$$

is null-homotopic, or equivalently such that $e \times f : S^m \hookrightarrow \mathbb{R} \times S^{2m}$ is framed null-cobordant (by $D^{m+1} \subset \mathbb{R} \times S^{2m}$ for $m \geq 2$, and by a Seifert surface $N^2 \subset \mathbb{R} \times S^2$ for $m = 1$). For this preferred choice of framing 6.95 (i) gives

$$H(\phi) = \mu^{\mathbb{Z}}(f) + (\delta\nu_f, \nu_f)^{\mathbb{Z}} = 0 \in \mathbb{Z} .$$

(ii) The Whitney immersion $f : S^m \looparrowright S^{2m}$ has

$$(\delta\nu_f, \nu_f)^{\mathbb{Z}} = -(\delta\tau_{S^m}, \tau_{S^m})^{\mathbb{Z}} = -1 \in \pi_{m+1}(BO(m+1), BO(m)) = \mathbb{Z}$$

with

$$\mu^{\mathbb{Z}}(f) = 1, \quad H(\phi) = \mu^{\mathbb{Z}}(f) + (\delta\nu_f, \nu_f)^{\mathbb{Z}} = 0 \in \mathbb{Z}.$$

In particular, $f : S^1 \looparrowright S^2$ is just the figure 8 immersion.

(iii) The integral geometric self-intersection of a framed immersion $f : M^m \looparrowright S^{2m}$ is identified by 6.95 (ii) with the Hopf invariant of the Pontryagin-Thom map $\phi : \Sigma(S^{2m})^\infty \rightarrow S^{m+1}$

$$\mu^{\mathbb{Z}}(f) = H(\phi) \in \text{im}(H : \pi_{2m+1}(S^{m+1}) \rightarrow \mathbb{Z}) = \begin{cases} 0 & \text{if } m \text{ is even} \\ \mathbb{Z} & \text{if } m = 1, 3, 7 \\ 2\mathbb{Z} & \text{otherwise.} \end{cases}$$

Thus the number of self-intersections is even, except in the Hopf invariant 1 dimensions $m = 1, 3, 7$. \square

Let $N^{2m} \subset S^{2m+1}$ be a codimension 1 framed submanifold, and let $i : \mathbb{R} \times N \hookrightarrow S^{2m+1}$ be the inclusion of a tubular neighbourhood. Use the disjoint embeddings

$$\begin{aligned} i_0 : N &\hookrightarrow S^{2m+1}; \quad x \mapsto i(0, x), \\ i_1 : N &\hookrightarrow S^{2m+1}; \quad x \mapsto i(1, x) \end{aligned}$$

to define a pairing

$$s : N \times N \rightarrow S^{2m}; \quad (x, y) \mapsto \frac{i_0(x) - i_1(y)}{\|i_0(x) - i_1(y)\|}.$$

The induced product in homology

$$\sigma : H_m(N) \times H_m(N) \rightarrow H_{2m}(S^{2m}) = \mathbb{Z}; \quad (x, y) \mapsto s_*(x \otimes y)$$

is such that for any embeddings $f, f' : S^m \hookrightarrow N$

$$\begin{aligned} \sigma(f_*[S^m], f'_*[S^m]) &= \text{linking number}(i_0 f \sqcup i_1 f' : S^m \sqcup S^m \hookrightarrow S^{2m+1}) \\ &= \text{degree}(S^m \xrightarrow{i_0 f} S^{2m+1} \setminus i_1 f'(S^m) \simeq S^m) \in \mathbb{Z}. \end{aligned}$$

For any embedding $f : S^m \hookrightarrow N$ (assumed unknotted in S^{2m+1} if $m = 1$) the composite $i_0 f : S^m \hookrightarrow S^{2m+1}$ is isotopic to the standard embedding $S^m \hookrightarrow S^{2m+1}$, so that the normal bundle $\nu_f : S^m \rightarrow BO(m)$ is equipped with a stable trivialization $\delta\nu_f : \nu_f \oplus \epsilon \cong \epsilon^{m+1}$. Define the disjoint embeddings

$$\begin{aligned} j_0 : S^m &\hookrightarrow E(\nu_f \oplus \epsilon) = E(\nu_f) \times \mathbb{R} ; x \mapsto (x, 0, 0) , \\ j_1 : S^m &\hookrightarrow E(\nu_f \oplus \epsilon) = E(\nu_f) \times \mathbb{R} ; x \mapsto (x, 0, 1) . \end{aligned}$$

It follows from the commutative diagram

$$\begin{array}{ccc} S^m & \xrightarrow{j_1} & E(\nu_f \oplus \epsilon) \setminus j_0(S^m) \cong E(\epsilon^{m+1}) \setminus E(\delta\nu_f)^{-1} j_0(S^m) \\ \downarrow i_1 f & & \downarrow \\ S^{2m+1} \setminus M & \longrightarrow & S^{2m+1} \setminus i_0 f(S^m) \end{array}$$

that

$$\begin{aligned} \sigma(f_*[S^m], f_*[S^m]) &= \text{linking number}(i_0 f(S^m) \cup i_1 f(S^m) \subset S^{2m+1}) \\ &= (\delta\nu_f, \nu_f)_{\mathbb{Z}} \in \pi_{m+1}(BO(m+1), BO(m)) = \mathbb{Z} . \end{aligned}$$

The above construction applies also to the embedding of a Seifert surface $N^{2m} \hookrightarrow S^{2m+1}$ for a $(2m-1)$ -knot $k : S^{2m-1} \hookrightarrow S^{2m+1}$, with

$$\partial N = S^{2m-1} \hookrightarrow S^{2m+1} .$$

If $x_1, x_2, \dots, x_\ell \in H_m(N)$ is a basis for the f.g. free \mathbb{Z} -module $H_m(N)/\text{torsion}$ then $(\sigma(x_i, x_j))$ is the Seifert matrix of k with respect to N .

Let N^{2m} be a $2m$ -dimensional manifold with an embedding $i : \mathbb{R} \times N \hookrightarrow S^{2m+1}$, as before. Let

$$(e \times f : S^m \hookrightarrow \mathbb{R} \times N , f : S^m \looparrowright N)$$

be an $(m, 2m, 1)$ -dimensional embedding-immersion pair, with Umkehr stable map $F : \Sigma N^+ \rightarrow \Sigma T(\nu_f)$ and integral self-intersection number

$$\mu^{\mathbb{Z}}(f) = \psi_{\mathbb{R}}(F)_*[N] \in \mathbb{Z} .$$

The composite $i(e \times f) : S^m \hookrightarrow S^{2m+1}$ is an embedding with normal bundle

$$\nu_{i(e \times f)} = \nu_f \oplus \epsilon : S^m \rightarrow BO(m+1) .$$

An isotopy of $i(e \times f)$ to the standard embedding $S^m \hookrightarrow S^{2m+1}$ determines a stable trivialization

$$\delta\nu_f : \nu_f \oplus \epsilon \cong \epsilon^{m+1}$$

which is classified by the Hopf invariant of the homotopy equivalence

$$T(\delta\nu_f) : \Sigma T(\nu_f) \simeq S^{2m+1} \vee S^{m+1} ,$$

that is

$$(\delta\nu_f, \nu_f)^{\mathbb{Z}} = \psi_{\mathbb{R}}(T(\delta\nu_f))[T(\nu_f)] \in \pi_{m+1}(BO(m+1), BO(m)) = \mathbb{Z}.$$

(For $m \neq 1, 3, 7$ this is independent of $e : S^m \rightarrow \mathbb{R}$, and is just the integer $d \in \mathbb{Z}$ such that $\nu_f = d\tau_{S^m} \in \ker(\pi_m(BO(m)) \rightarrow \pi_m(BO(m+1)))$.) Let $G : S^{2m+1} \rightarrow \Sigma N^+$ be the Pontryagin-Thom map for $i : \mathbb{R} \times N \hookrightarrow S^{2m+1}$. (If N is a Seifert surface of a knot $k : S^{2m-1} \hookrightarrow S^{2m+1}$ then G is the Umkehr stable map for the ultranormal normal map

$$(g, c) = \text{inclusion} : (N, \partial N) \rightarrow (D^{2m+2}, k(S^{2m-1}))$$

of 6.12). The composite

$$T(\delta\nu_f)FG : S^{2m+1} \xrightarrow{G} \Sigma N^+ \xrightarrow{F} \Sigma T(\nu_f) \xrightarrow{T(\delta\nu_f)} S^{2m+1} \vee S^{m+1}$$

is the Pontryagin-Thom map of the standard embedding $S^m \hookrightarrow S^{2m+1}$, so that

$$T(\delta\nu_f)FG \simeq \text{inclusion} : S^{2m+1} \rightarrow S^{2m+1} \vee S^{m+1}$$

and by the composition formula 5.49 (vi)

$$\begin{aligned} & \psi_{\mathbb{R}}(T(\delta\nu_f)FG)[S^{2m}] \\ &= (F \otimes F)_* \psi_{\mathbb{R}}(G)_*[S^{2m}] + \psi_{\mathbb{R}}(F)_*[N] + \psi_{\mathbb{R}}(T(\delta\nu_f))[T(\nu_f)] \\ &= 0 \in \mathbb{Z}. \end{aligned}$$

The evaluation of the ultraquadratic construction on G (5.52)

$$\psi_{\mathbb{R}}(G) : S^{2m} \rightarrow N^+ \wedge N^+$$

on $[S^{2m}]$ is a homology class

$$h'_{\mathbb{R}}(G)_*[S^{2m}] \in \tilde{H}_{2m}(N^+ \wedge N^+) = H_{2m}(N \times N)$$

such that

$$\sigma : H_m(N) \times H_m(N) \rightarrow \mathbb{Z} ; (x, y) \mapsto -\langle x^* \times y^*, h'_{\mathbb{R}}(G)_*[S^{2m}] \rangle$$

with $x^*, y^* \in H^m(N)$ the Poincaré duals of $x, y \in H_m(N)$. Thus

$$\begin{aligned} \sigma(f_*[S^m], f_*[S^m]) &= -(F \otimes F)_* h'_{\mathbb{R}}(G)_*[S^{2m}] \\ &= h'_{\mathbb{R}}(F)_*[N] + h'_{\mathbb{R}}(T(\delta\nu_f))[T(\nu_f)] \\ &= \mu^{\mathbb{Z}}(f) + (\delta\nu_f, \nu_f)^{\mathbb{Z}} \in \mathbb{Z} \end{aligned}$$

In particular, if $x = f_*[S^m] \in H_m(N)$ for an embedding $f : S^m \hookrightarrow N$ then

$$\mu^{\mathbb{Z}}(f) = 0, \quad \sigma(x, x) = (\delta\nu_f, \nu_f)^{\mathbb{Z}} \in \mathbb{Z}.$$

On the other hand, if $x = f_*[S^m] \in H_m(N)$ for a framed immersion $f : S^m \looparrowright N$ then

$$(\delta\nu_f, \nu_f)^{\mathbb{Z}} = 0 \quad , \quad \sigma(x, x) = \mu^{\mathbb{Z}}(f) \in \mathbb{Z} .$$

Now suppose that N is $(m-1)$ -connected. Every $x \in H_m(N)$ is represented by a framed immersion $f : S^m \looparrowright N$. If $m \geq 3$ then every $x \in H_m(N)$ is also represented by an embedding $f : S^m \hookrightarrow N$ (which will not in general be framed). An element $x \in H_m(N)$ is such that $\sigma(x, x) = 0$ if (and for $m \geq 3$ only if) it can be represented by a framed embedding $f : S^m \hookrightarrow N$, in which case x can be killed by ambient surgery to obtain an ambient cobordant submanifold

$$(N \hookrightarrow S^{2m+1}) \mapsto (N' \hookrightarrow S^{2m+1})$$

with

$$N' = \text{cl.}(N \setminus (f(S^m) \times D^m)) \cup D^{m+1} \times S^{m-1}$$

Remark 6.97. See Levine [40] for the classification in the case $m \geq 3$ of $(2m-1)$ -knots $k : S^{2m-1} \hookrightarrow S^{2m+1}$ which are simple, i.e. which admit an $(m-1)$ -connected Seifert surface $N^{2m} \hookrightarrow S^{2m+1}$. The isotopy classes of such knots are in one-one correspondence with the S -equivalence classes of Seifert matrices $\sigma = (\sigma_{ij})_{1 \leq i, j \leq \ell}$ with $\sigma_{ij} \in \mathbb{Z}$ and $(\sigma_{ij} + (-1)^m \sigma_{ji})$ invertible. The knot k corresponding to σ has an $(m-1)$ -connected Seifert surface $N^{2m} \hookrightarrow S^{2m+1}$ with a handle presentation involving ℓ m -handles

$$N = D^{2m} \cup \bigcup_{\ell} (D^m \times D^m) .$$

The attaching maps are embeddings $\phi_i : S^{m-1} \times D^m \hookrightarrow S^{2m-1}$ with

$$\begin{aligned} \text{linking number}(\phi_i(S^{m-1} \times \{0\}) \cup \phi_j(S^{m-1} \times \{0\}) \subset S^{2m-1}) \\ = \sigma_{ij} + (-1)^m \sigma_{ji} \in \mathbb{Z} \quad (1 \leq i, j \leq \ell, i \neq j) . \end{aligned}$$

The basis elements

$$x_i = (0, \dots, 0, 1, 0, \dots, 0) \in H_m(N) = \mathbb{Z}^{\ell} \quad (1 \leq i \leq \ell)$$

are represented by $(m, 2m, 1)$ -dimensional embedding-immersion pairs

$$(e_i \times f_i : S^m \hookrightarrow \mathbb{R} \times N^{2m} , f_i : S^m \looparrowright N)$$

with $e_i \times f_i$ framed and f_i an embedding, such that

$$(\delta\nu_{f_i}, \nu_{f_i}) = \sigma_{ii}(\delta\tau_{S^m}, \tau_{S^m}) \in \pi_{m+1}(BO(m+1), BO(m)) = \mathbb{Z} .$$

□

Proposition 6.98. *Let $N^n \subset M^{n+1}$ be a framed codimension 1 submanifold with complement P , so that*

$$M = N \times I \cup_{i_0, i_1} P$$

and there is defined a cofibration sequence

$$P^+ \longrightarrow M^+ \xrightarrow{F} \Sigma N^+ \xrightarrow{G} \Sigma P^+$$

with

$$\begin{aligned} F &= \text{projection} : M^+ \rightarrow M/P = N \times I / \partial(N \times I) = \Sigma N^+, \\ G &= \Sigma i_0 - \Sigma i_1 : \Sigma N^+ \rightarrow \Sigma P^+ \end{aligned}$$

where $i_0, i_1 : N \rightarrow P$ are the inclusions of the two boundary components. The spectral Hopf invariant of F , the Hopf invariant of G and i_0, i_1 are related by a stable homotopy commutative diagram

$$\begin{array}{ccc} \Sigma M^+ & \xrightarrow{sh_{\mathbb{R}}(F)} & \Sigma P^+ \wedge \Sigma P^+ \\ \Sigma F \downarrow & \nearrow -h_{\mathbb{R}}(G) & \uparrow G \wedge \Sigma i_1 \\ \Sigma^2 N^+ & \xrightarrow{\Sigma^2 \Delta_N} & \Sigma N^+ \wedge \Sigma N^+ \end{array}$$

Proof. By 5.63 we have the stable homotopy identity

$$sh_{\mathbb{R}}(F) + h_{\mathbb{R}}(G)\Sigma F = 0 : \Sigma M^+ \rightarrow \Sigma P^+ \wedge \Sigma P^+.$$

By the sum formula 5.33 (viii)

$$h_{\mathbb{R}}(G) = h_{\mathbb{R}}(\Sigma i_0) + h_{\mathbb{R}}(-\Sigma i_1) + (\Sigma i_0 \wedge -\Sigma i_1)\Sigma^2 \Delta_N : \Sigma^2 N^+ \rightarrow \Sigma P^+ \wedge \Sigma P^+.$$

By the suspension formula 5.33 (iii)

$$h_{\mathbb{R}}(\Sigma i_0) = 0 : \Sigma^2 N^+ \rightarrow \Sigma P^+ \wedge \Sigma P^+.$$

By Example 5.37 the Hopf invariant map of

$$-1 : S^1 \rightarrow S^1 ; t \mapsto 1 - t$$

is given by

$$h_{\mathbb{R}}(-1 : S^1 \rightarrow S^1) = 1 : S^2 \rightarrow S^2.$$

By the product formula 5.33 (vi)

$$\begin{aligned}
h_{\mathbb{R}}(-\Sigma i_1) &= h_{\mathbb{R}}(-1 \wedge i_1 : S^1 \wedge N^+ \rightarrow S^1 \wedge P^+) \\
&= (\Sigma i_1 \wedge \Sigma i_1) \Sigma^2 \Delta_N : \Sigma^2 N^+ \rightarrow \Sigma P^+ \wedge \Sigma P^+ .
\end{aligned}$$

Substitution in the expression for $h_{\mathbb{R}}(G)$ gives

$$\begin{aligned}
h_{\mathbb{R}}(G) &= ((\Sigma i_0 - \Sigma i_1) \wedge -\Sigma i_1) \Sigma^2 \Delta_N \\
&= -(G \wedge \Sigma i_1) \Sigma^2 M^+ : \Sigma^2 N^+ \rightarrow \Sigma P^+ \wedge \Sigma P^+ .
\end{aligned}$$

□

Remark 6.99. Let $k : S^{n-1} \subset S^{n+1}$ be an $(n-1)$ -knot, and let $N^n \subset S^{n+1}$ be a Seifert surface, so that $\partial N = k(S^{n-1})$. The knot complement is a codimension 0 submanifold

$$X = \text{cl.}(S^{n+1} \setminus (k(S^{n-1}) \times D^2)) \subset S^{n+1}$$

which can be cut along $N \subset X$ to obtain a codimension 0 submanifold

$$P = \text{cl.}(X \setminus (N \times I)) \subset X$$

such that

$$\begin{aligned}
\partial P &= N \cup_{\partial N} N , \quad X = (N \times I) \cup_{i_0, i_1} P , \\
S^{n+1} &= (N \times I \cup \partial N \times D^2) \cup_{i_0, i_1} P
\end{aligned}$$

with $i_0, i_1 : N \rightarrow P$ the two inclusions. As in 6.98 there is defined a cofibration sequence

$$P^+ \longrightarrow S^{n+1} \xrightarrow{F} \Sigma(N/\partial N) \xrightarrow{G} \Sigma P^+$$

such that up to homotopy

$$\Sigma i_0 - \Sigma i_1 : \Sigma N^+ \longrightarrow \Sigma(N/\partial N) \xrightarrow{G} \Sigma P^+ ,$$

and up to stable homotopy

$$sh_{\mathbb{R}}(F) = (G \wedge \Sigma i_1)(\Sigma^2 \Delta_{N/\partial N})(\Sigma F) : S^{n+2} \rightarrow \Sigma P^+ \wedge \Sigma P^+ .$$

The expression of the spectral Hopf invariant $sh_{\mathbb{R}}(F)$ of the unstable normal invariant F of $N^n \subset S^{n+1}$ in terms of the homotopy-theoretic Seifert form i_1 is a generalization of Theorem 3.1 of Richter [54]. □

Chapter 7

The π -equivariant geometric Hopf invariant

The stable homotopy constructions of §5 (in particular the geometric Hopf invariant) and the double point theorem of §6 are so natural that they have π -equivariant versions, for any group discrete π , inducing the corresponding chain level constructions of Ranicki [49, 50], with applications to non-simply-connected surgery obstruction theory.

7.1 π -equivariant S -duality

Definition 7.1. (i) Given pointed π -spaces X, Y define the *integral π -equivariant homotopy groups*

$$\{X; Y\}_{0, \pi} = \varinjlim_U [U^\infty \wedge X; U^\infty \wedge Y]_\pi$$

with U running over finite-dimensional inner product spaces.

(ii) Given pointed π -spaces X, Y, Z , a finite-dimensional inner product space V and a map

$$\sigma : V^\infty \rightarrow X \wedge_\pi Y$$

define the *slant products*

$$\begin{aligned} \sigma \backslash - & : \{X, Z\}_{0, \pi} \rightarrow \{V^\infty, Z \wedge_\pi Y\} ; \\ (f : U^\infty \wedge X \rightarrow U^\infty \wedge Z) & \mapsto ((f \wedge 1)(1 \wedge \sigma) : (U \oplus V) \rightarrow Z \wedge_\pi Y) , \\ \sigma \backslash - & : \{Y, Z\}_{0, \pi} \rightarrow \{V^\infty, X \wedge_\pi Z\} ; \\ (f : U^\infty \wedge Y \rightarrow U^\infty \wedge Z) & \mapsto ((f \wedge 1)(1 \wedge \sigma) : (U \oplus V) \rightarrow X \wedge_\pi Z) . \end{aligned}$$

(iii) A map $\sigma : V^\infty \rightarrow X \wedge_\pi Y$ is an *integral π -equivariant S -duality* map if the slant products $\sigma \backslash -$ in (ii) are isomorphisms for every π -space Z . \square

Proposition 7.2. (Ranicki [50, §3]) *For any semifree finite pointed CW π -complex X there exist a finite-dimensional inner product space V , a semifree finite pointed CW π -complex Y and an integral π -equivariant S -duality map $\sigma : V^\infty \rightarrow X \wedge_\pi Y$.* \square

Remark 7.3. The theory of [50] deals with the ‘integral’ stable π -equivariant homotopy groups

$$\{X; Y\}_{0, \pi} = \varinjlim_V [V^\infty \wedge X, V^\infty \wedge Y]_\pi$$

with the direct limit running over all the finite-dimensional inner product spaces V . The forgetful map for $\pi = \mathbb{Z}_2$

$$\{X; Y\}_{0, \mathbb{Z}_2} \rightarrow \{X; Y\}_{\mathbb{Z}_2}$$

is in general neither injective nor surjective, but by Adams [3] (cf. Proposition 4.32 (ii)) it is an isomorphism for finite CW \mathbb{Z}_2 -complexes X, Y with X semifree, and in this case the \mathbb{Z}_2 -equivariant S -duality theories of Wirthmüller [75] (cf. Proposition 4.85) and [50] coincide. \square

7.2 The π -equivariant constructions

Let X, Y be pointed π -spaces, and let V be an inner product space with trivial π -action. The geometric Hopf invariant (5.3) of a π -equivariant map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is a \mathbb{Z}_2 -equivariant map

$$h_V(F) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y$$

which is also π -equivariant, with

$$\pi \times Y \wedge Y \rightarrow Y \wedge Y ; (g, (y_1, y_2)) \mapsto (gy_1, gy_2) .$$

The ‘ π -equivariant geometric Hopf invariant of F ’ is the induced map of the π -quotients

$$h_V(F)/\pi : \Sigma S(LV)^+ \wedge V^\infty \wedge X/\pi \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge_\pi Y$$

with

$$\begin{aligned} Y \wedge_\pi Y &= Y \wedge Y / \{(y_1, y_2) \sim (gy_1, gy_2) \mid g \in \pi, y_1, y_2 \in Y\} , \\ T : Y \wedge_\pi Y &\rightarrow Y \wedge_\pi Y ; [y_1, y_2] \mapsto [y_2, y_1] . \end{aligned}$$

Proposition 7.4. (i) *For any pointed space X , any pointed π -space Y and any inner product space V there is defined a long exact sequence of abelian groups/pointed sets*

$$\begin{aligned} \dots \longrightarrow [\Sigma X, Y \wedge_\pi Y]_{\mathbb{Z}_2} &\xrightarrow{s_{LV}^*} [\Sigma S(LV)^+ \wedge X, Y \wedge_\pi Y]_{\mathbb{Z}_2} \xrightarrow{\alpha_{LV}^*} \\ &[LV^\infty \wedge X, Y \wedge_\pi Y]_{\mathbb{Z}_2} \xrightarrow{0_{LV}^*} [X, Y \wedge_\pi Y]_{\mathbb{Z}_2} \end{aligned}$$

(ii) *For any pointed space X , any pointed π -space Y and any inner product spaces V, W there is defined a split short exact sequence of abelian groups*

$$\begin{aligned} 0 \rightarrow [\Sigma S(LV)^+ \wedge W^\infty \wedge X, LV^\infty \wedge W^\infty \wedge (Y \wedge_\pi Y)]_{\mathbb{Z}_2} &\xrightarrow{\alpha_{LV}^*} \\ [LV^\infty \wedge W^\infty \wedge X, LV^\infty \wedge W^\infty \wedge (Y \wedge_\pi Y)]_{\mathbb{Z}_2} &\xrightarrow{\rho} [W^\infty \wedge X, W^\infty \wedge Y/\pi] \rightarrow 0 \end{aligned}$$

with ρ defined by the fixed points of the \mathbb{Z}_2 -action. The map ρ is split by

$$\begin{aligned} \sigma : [W^\infty \wedge X, W^\infty \wedge Y/\pi] &\rightarrow [LV^\infty \wedge W^\infty \wedge X, LV^\infty \wedge W^\infty \wedge (Y \wedge_\pi Y)]_{\mathbb{Z}_2} ; \\ F &\mapsto \sigma(F) = 1_{LV^\infty \wedge W^\infty} \wedge \Delta F \quad (\sigma(F)(v, w, x) = (v, u, y) \text{ if } F(w, x) = (u, y)) , \end{aligned}$$

with

$$\Delta : Y/\pi = (Y \wedge_\pi Y)^{\mathbb{Z}_2} \hookrightarrow Y \wedge_\pi Y ; [y] \mapsto [y, y] .$$

For any \mathbb{Z}_2 -equivariant map

$$G : LV^\infty \wedge W^\infty \wedge X \rightarrow LV^\infty \wedge W^\infty \wedge (Y \wedge_\pi Y)$$

the map of fixed points

$$F = \rho(G) : W^\infty \wedge X \rightarrow W^\infty \wedge Y/\pi$$

is such that $(1 \wedge \Delta)\sigma(F)$ and G agree on $0^+ \wedge W^\infty \wedge X$, with the relative difference \mathbb{Z}_2 -equivariant map

$$\delta((1 \wedge \Delta)\sigma(F), G) : \Sigma S(LV)^+ \wedge W^\infty \wedge X \rightarrow LV^\infty \wedge W^\infty \wedge (Y \wedge_\pi Y)$$

such that

$$G - (1 \wedge \Delta)\sigma(F) = \alpha_{LV}^* \delta((1 \wedge \Delta)\sigma(F), G) \in \text{im}(\alpha_{LV}^*) = \ker(\rho) .$$

Proof. This is just a special case of Proposition 4.11, with Y replaced by $Y \wedge_\pi Y$, and $Y^{\mathbb{Z}_2}$ replaced by $(Y \wedge_\pi Y)^{\mathbb{Z}_2} = Y/\pi$. \square

Proposition 7.5. (i) *For any inner product spaces U, V , any pointed spaces X and any pointed π -space Y there is defined a commutative braid of exact sequences of stable homotopy groups*

$$\begin{array}{ccccc}
 & & \alpha_{LV} & & \\
 & & \curvearrowright & & \\
 A_1 & & \{X; S(LV)^+ \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} & \xrightarrow{s_{LV}} & \{S(LU)^+ \wedge X; Y \wedge_\pi Y\}_{\mathbb{Z}_2} \\
 & \nearrow & & \nwarrow s_{LU}^* & \\
 & A_2 & & & A_3 \\
 & \searrow & & \nearrow 0_{LV}^* & \\
 A_4 & & \{LU^\infty \wedge X; Y \wedge_\pi Y\}_{\mathbb{Z}_2} & \xrightarrow{0_{LU}} & \{X; V^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} \\
 & \nwarrow \alpha_{LU}^* & & \nearrow & \\
 & & \curvearrowright & &
 \end{array}$$

with

$$\begin{aligned}
 A_1 &= \{\Sigma X; LV^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2}, \quad A_2 = \{\Sigma S(LU \oplus LV)^+ \wedge X; LV^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2}, \\
 A_3 &= \{S(LU \oplus LV)^+ \wedge X; V^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2}, \quad A_4 = \{\Sigma S(LU)^+ \wedge X; Y \wedge_\pi Y\}_{\mathbb{Z}_2}.
 \end{aligned}$$

(ii) *For any inner product space $V \subseteq \mathbb{R}(\infty)$*

$$\{X; LV^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} = \{X; Y/\pi\} \oplus \{X; S(L\mathbb{R}(\infty))/S(LV) \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2},$$

with a split short exact sequence

$$\begin{aligned}
 0 \rightarrow \{X; S(L\mathbb{R}(\infty))/S(LV) \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} &\xrightarrow{\delta} \{X; LV^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} \\
 &\xrightarrow{\rho} \{X; Y/\pi\} \rightarrow 0
 \end{aligned}$$

where δ is induced by the \mathbb{Z}_2 -equivariant connecting map

$$\delta : S(L\mathbb{R}(\infty))/S(LV) = \varinjlim_k S(L\mathbb{R}^k)/S(LV) \rightarrow sS(LV) = LV^\infty,$$

ρ is defined by the fixed points of the \mathbb{Z}_2 -action and ρ is split by

$$\sigma : \{X; Y/\pi\} \rightarrow \{X; LV^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2}; \quad F \mapsto (0 \wedge \Delta_{Y/\pi})F.$$

In particular, for $V = \mathbb{R}(\infty)$ the fixed point map is an isomorphism

$$\rho : \{X; L\mathbb{R}(\infty)^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} \xrightarrow{\cong} \{X; Y/\pi\} ,$$

and the case $V = \{0\}$ gives

$$\{X; Y \wedge_\pi Y\}_{\mathbb{Z}_2} = \{X; Y/\pi\} \oplus \{X; S(L\mathbb{R}(\infty))^+ \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} .$$

(iii) For any inner product space U there is defined a long exact sequence

$$\begin{aligned} \dots &\longrightarrow \{\Sigma S(LU \oplus L\mathbb{R}(\infty))^+ \wedge X; L\mathbb{R}(\infty)^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} \\ &\longrightarrow \{LU^\infty \wedge X; Y \wedge_\pi Y\}_{\mathbb{Z}_2} \xrightarrow{\rho} \{X; Y/\pi\} \\ &\longrightarrow \{S(LU \oplus L\mathbb{R}(\infty))^+ \wedge X; L\mathbb{R}(\infty)^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} \longrightarrow \dots \end{aligned}$$

with ρ defined by the fixed points of the \mathbb{Z}_2 -action, and

$$\begin{aligned} &\{S(LU \oplus L\mathbb{R}(\infty))^+ \wedge X; L\mathbb{R}(\infty)^\infty \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} \\ &= \{LU^\infty \wedge X; \Sigma S(LU \oplus L\mathbb{R}(\infty))^+ \wedge Y \wedge_\pi Y\}_{\mathbb{Z}_2} . \end{aligned}$$

(iv) For any pointed π -space X the morphism

$$\begin{aligned} 0_{LV} &: \{X/\pi; X \wedge_\pi X\}_{\mathbb{Z}_2} \\ &\rightarrow \{X/\pi; LV^\infty \wedge X \wedge_\pi X\}_{\mathbb{Z}_2} = \{V^\infty \wedge X/\pi; V^\infty \wedge LV^\infty \wedge X \wedge_\pi X\}_{\mathbb{Z}_2} \end{aligned}$$

sends $\Delta_{X/\pi}$ to $0_{LV} \wedge \Delta_{X/\pi} = (\kappa_V^{-1} \wedge 1)\Delta_{V^\infty \wedge X/\pi}$.

Proof. This is a \mathbb{Z}_2 -equivariant version of the braid of Proposition 3.3; for $\pi = \{1\}$ this is just Proposition 4.41.

□

The definitions of the various constructions of §5 translate verbatim into their π -equivariant analogues, with matching properties. We shall only state the definitions here.

Definition 7.6. The π -equivariant symmetric construction $\phi_V(X)$ is defined for a pointed π -space X and an inner product space V to be the $\pi \times \mathbb{Z}_2$ -equivariant map

$$\phi_V(X) = {}_{sLV} \wedge \Delta_X : S(LV)^+ \wedge X \rightarrow X \wedge X ; (v, [x]) \mapsto [x, x] .$$

□

Passing to the π -quotients there is defined a \mathbb{Z}_2 -equivariant map

$$\phi_V(X) = s_{LV} \wedge \Delta_X : S(LV)^+ \wedge X/\pi \rightarrow X \wedge_\pi X .$$

Definition 7.7. The π -equivariant *geometric Hopf invariant* of a π -equivariant map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is the \mathbb{Z}_2 -equivariant map given by the relative difference of the \mathbb{Z}_2 -equivariant maps

$$p = (1 \wedge \Delta_Y)(1 \wedge F) , \quad q = (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge \Delta_X)$$

with

$$\begin{aligned} h_V(F) &= \delta(p, q) : \Sigma S(LV)^+ \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge Y \wedge Y ; \\ (t, u, v, x) &\mapsto \begin{cases} p([1 - 2t, u], v, x) & \text{if } 0 \leq t \leq 1/2 \\ q([2t - 1, u], v, x) & \text{if } 1/2 \leq t \leq 1 \end{cases} \\ &(t \in I, u \in S(LV), v \in V, x \in X) . \end{aligned}$$

□

Definition 7.8. The π -equivariant *stable geometric Hopf invariant* of a π -equivariant map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is the stable relative difference of p and q (3.6), the $\pi \times \mathbb{Z}_2$ -equivariant map

$$h'_V(F) = \delta'(p, q) : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge S(LV)^+ \wedge Y \wedge Y$$

which we shall regard as a stable \mathbb{Z}_2 -equivariant map

$$h'_V(F) : X/\pi \rightarrow S(LV)^+ \wedge Y \wedge_\pi Y .$$

□

Definition 7.9. The π -equivariant *quadratic construction* on a π -equivariant map $F : V^\infty \wedge X \rightarrow V^\infty \wedge Y$ is the stable π -equivariant map

$$\psi_V(F) : X \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)$$

given by the image of the \mathbb{Z}_2 -equivariant stable geometric Hopf invariant (7.8)

$$h'_V(F) = \delta'(p, q) : LV^\infty \wedge V^\infty \wedge X \rightarrow LV^\infty \wedge V^\infty \wedge S(LV)^+ \wedge Y \wedge Y$$

under the isomorphism given by Proposition 4.32

$$\{X; S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge Y)\}_\pi \cong \{X; S(LV)^+ \wedge Y \wedge Y\}_{\pi \times \mathbb{Z}_2} .$$

□

Passing to the π -quotients gives

$$\psi_V : [V^\infty \wedge X, V^\infty \wedge Y]_\pi \rightarrow \{X/\pi; S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge_\pi Y)\} \cong \{X; S(LV)^+ \wedge (Y \wedge_\pi Y)\}_{\mathbb{Z}_2}.$$

Definition 7.10. The π -equivariant *ultraquadratic construction* on a π -equivariant map $F : \Sigma X \rightarrow \Sigma Y$ is the quadratic construction for the special case $V = \mathbb{R}$

$$\widehat{\psi}(F) = \psi_{\mathbb{R}}(F) : X/\pi \rightarrow S(LV)^+ \wedge_{\mathbb{Z}_2} (Y \wedge_\pi Y) = Y \wedge_\pi Y,$$

identifying $S(LV) = S^0 = \{1, -1\}$ with \mathbb{Z}_2 acting by permutation. \square

Definition 7.11. The π -equivariant *spectral Hopf invariant map* of a π -equivariant map $F : X \rightarrow V^\infty \wedge Y$ is the $\pi \times \mathbb{Z}_2$ -equivariant map given by the relative difference (1.3)

$$\begin{aligned} sh_V(F) &= \delta((G \wedge G)\delta\phi_V(V^\infty \wedge Y)(1 \wedge F), \phi_V(\mathcal{C}(F))(1 \wedge GF)) : \\ &\quad \Sigma S(LV)^+ \wedge X \rightarrow \mathcal{C}(F) \wedge \mathcal{C}(F) \end{aligned}$$

with $G : V^\infty \wedge Y \rightarrow \mathcal{C}(F)$ the inclusion in the mapping cone and

$$1 \wedge GF : CS(LV)^+ \wedge X = S(LV)^+ \wedge CX \rightarrow S(LV)^+ \wedge \mathcal{C}(F)$$

the π -equivariant null-homotopy of $1 \wedge GF : S(LV)^+ \wedge X \rightarrow S(LV)^+ \wedge \mathcal{C}(F)$ determined by the inclusion $GF : CX \rightarrow \mathcal{C}(F) = (V^\infty \wedge Y) \cup_F CX$. \square

Here is the π -equivariant version of the Double Point Theorem 6.19.

Let $e = (g, f) : V \times X \hookrightarrow V \times Y$ be an embedding of a map $f : X \rightarrow Y$. Given a regular cover $p : \tilde{Y} \rightarrow Y$ with group of covering translations π let

$$\tilde{X} = f^* \tilde{Y} = \{(x, \tilde{y}) \in X \times \tilde{Y} \mid f(x) = p(\tilde{y}) \in Y\}$$

be the pullback cover of X , so that there are defined π -equivariant lifts of f, e

$$\begin{aligned} \tilde{f} : \tilde{X} &\rightarrow \tilde{Y} ; (x, \tilde{y}) \mapsto \tilde{y}, \\ \tilde{e} = (\tilde{g}, \tilde{f}) : V \times \tilde{X} &\hookrightarrow V \times \tilde{Y} ; (v, (x, \tilde{y})) \mapsto (g(v, x), \tilde{y}) \end{aligned}$$

and hence a π -equivariant Umkehr map

$$\tilde{F} : V^\infty \wedge \tilde{Y}^\infty \rightarrow V^\infty \wedge \tilde{X}^\infty.$$

Proposition 7.12. The π -equivariant *geometric Hopf invariant* $h_V(\tilde{F})$ is given up to natural $\pi \times \mathbb{Z}_2$ -equivariant homotopy by the composite

$$\begin{aligned} h_V(\tilde{F})/\pi &= (1 \wedge A)h_V(\tilde{F})_Y : \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty \\ &\rightarrow LV^\infty \wedge V^\infty \wedge (\tilde{X} \times_{\tilde{Y}} \tilde{X})^\infty \rightarrow LV^\infty \wedge V^\infty \wedge (\tilde{X} \times_\pi \tilde{X})^\infty \end{aligned}$$

with

$$\begin{aligned} h_V(\tilde{F})/\pi &= (1 \wedge i)\tilde{F}_1 + (1 \wedge \Delta_X)\delta(\tilde{F}_2, \tilde{F}_3) : \\ \Sigma S(LV)^+ \wedge V^\infty \wedge Y^\infty &\rightarrow \\ LV^\infty \wedge V^\infty \wedge (X \times_Y X)^\infty &= LV^\infty \wedge V^\infty \wedge (D_2(f)^\infty \vee X^\infty) , \\ i &= \text{inclusion} : D_2(f) \rightarrow X \times_Y X ; (x_1, x_2) \mapsto (x_1, x_2) , \\ \Delta_X &= \text{diagonal} : X \rightarrow X \times_Y X ; x \mapsto (x, x) , \\ A &= \text{assembly} = \Delta_X \sqcup i : X \times_Y X = X \sqcup D_2(f) \rightarrow X \times X , \\ \tilde{F}_1, \tilde{F}_2, \tilde{F}_3 &\text{ as in 6.18.} \end{aligned}$$

7.3 Non-simply-connected surgery obstruction theory

The surgery obstruction group $L_n(A)$ was defined by Wall [30] for any ring with involution A to be the Witt group of $(-1)^k$ -quadratic forms over A for $n = 2k$, and the Witt group of automorphisms of $(-1)^k$ -quadratic forms for $n = 2k + 1$. The surgery obstruction $\sigma_*(f, b) \in L_n(\mathbb{Z}[\pi_1(X)])$ of an n -dimensional normal map $(f, b) : M \rightarrow X$ was constructed by first making (f, b) $[n/2]$ -connected by surgery below the middle dimension, and then using the middle-dimensional quadratic $\pi_1(X)$ -valued self-intersection data on the kernel homology $\mathbb{Z}[\pi_1(X)]$ -modules

$$K_i(M) = \ker(\tilde{f}_* : H_i(\tilde{M}) \rightarrow H_i(\tilde{X}))$$

with $\tilde{f} : \tilde{M} \rightarrow \tilde{X}$ a $\pi_1(X)$ -equivariant lift of f to the universal cover \tilde{X} of X and the pullback cover $\tilde{M} = f^*\tilde{X}$ of M . The algebraic L -group $L_n(A)$ was interpreted in Ranicki [50] as the cobordism group of n -dimensional quadratic Poincaré complexes over A . The surgery obstruction $\sigma_*(f, b)$ was represented in [50] by the cobordism class of an n -dimensional quadratic Poincaré complex (C, ψ) over $\mathbb{Z}[\pi_1(X)]$, with $H_*(C) = K_*(M)$ the kernel $\mathbb{Z}[\pi_1(X)]$ -modules, and ψ obtained from a $\pi_1(X)$ -equivariant stable Umkehr map $F : \Sigma^\infty \tilde{X}^+ \rightarrow \Sigma^\infty \tilde{M}^+$ S -dual to the induced map of $\pi_1(X)$ -equivariant Thom spaces $T(\tilde{b}) : T(\nu_{\tilde{M}}) \rightarrow T(\nu_{\tilde{X}})$ by a chain level quadratic construction capturing the self-intersection data, namely the ‘ $\pi_1(X)$ -equivariant quadratic construction’ ψ . The chain level ψ generalized the functional Steenrod squares used by Browder [7] in the homotopy-theoretic construction of the simply-connected surgery obstruction. In [50] ψ was generalized to $\pi_1(X)$ -equivariant

spectral quadratic and ultraquadratic constructions. In particular, we obtain the special cases required for non-simply-connected surgery obstruction theory:

- (a) The $\pi_1(N)$ -equivariant homotopy-theoretic expression for the double points of an immersion $f : M^m \looparrowright N^{2m}$

$$\mu(f) \in \mathbb{Z}[\pi_1(N)] / \{x - (-)^m \bar{x} \mid x \in \mathbb{Z}[\pi_1(N)]\}$$

of Wall [68, §5].

- (b) The $\pi_1(X)$ -equivariant homotopy-theoretic expression for the quadratic structure

$$\psi_b \in Q_n(\mathcal{C}(f^!))$$

on the $\mathbb{Z}[\pi_1(X)]$ -module chain complex kernel of an n -dimensional normal map $(f, b) : M \rightarrow X$ of Ranicki [50].

It is also possible to obtain the π -equivariant versions of the quadratic construction from the fibrewise Umkehr maps, as in Appendix A below.

Appendix A

The fibrewise Umkehr map

Given a normal map $(f, b) : M \rightarrow X$ we construct a fibrewise stable Umkehr map $F : X \sqcup_X X \rightarrow M \sqcup_X X$ over X , using the method of Crabb and James [13, p.261] and Crabb and Ranicki [14].

Let B be a reasonable space, such as a finite CW complex. A *fibrewise space* over B is a space X together with a map $p_X : X \rightarrow B$. A *fibrewise map* $f : X \rightarrow Y$ is a map such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow p_X & \swarrow p_Y \\ & B & \end{array}$$

commutes. For fibrewise pointed spaces X, Y let $[X, Y]_B$ denote the set of (fibrewise) homotopy classes of fibrewise maps $f : X \rightarrow Y$.

The *product* of fibrewise spaces X, Y is the fibrewise space

$$\begin{aligned} X \times_B Y &= \{(x, y) \in X \times Y \mid p_X(x) = p_Y(y) \in B\} \\ &= \bigcup_{b \in B} (p_X)^{-1}(b) \times (p_Y)^{-1}(b) \\ &= (p_X \times p_Y)^{-1}(\Delta_B) \subset X \times Y \end{aligned}$$

with $\Delta_B \subset B \times B$ the diagonal subspace.

Let $q_B : \tilde{B} \rightarrow B$ be the projection of a regular cover with group of covering translations π . For a fibrewise space X over B the pullback cover of X

$$\begin{aligned} \tilde{X} &= (p_X)^*(\tilde{B}) = X \times_B \tilde{B} \\ &= \{(x, \tilde{b}) \in X \times \tilde{B} \mid p_X(x) = q_B(\tilde{b}) \in B\} \end{aligned}$$

has projection

$$\tilde{X} \rightarrow X ; (x, \tilde{b}) \mapsto x \text{ } (p_X(x) = q_B(\tilde{b}) \in B)$$

with a π -equivariant lift of p_X

$$\tilde{p}_X : \tilde{X} \rightarrow \tilde{B} ; (x, \tilde{b}) \mapsto \tilde{b} .$$

For a fibrewise product $X \times_B Y$ the pullback cover of $X \times_B Y \rightarrow B$

$$\begin{aligned} \widetilde{X \times_B Y} &= \tilde{X} \times_{\tilde{B}} \tilde{Y} \\ &= \{(x, y, \tilde{b}) \in X \times Y \times \tilde{B} \mid p_X(x) = p_Y(y) = q_B(\tilde{b}) \in B\} \end{aligned}$$

can be regarded as the π -equivariant subspace $\tilde{X} \times_{\tilde{B}} \tilde{Y} \subset \tilde{X} \times \tilde{Y}$ consisting of all the $((x, \tilde{b}), (y, \tilde{c})) \in \tilde{X} \times \tilde{Y}$ with $\tilde{b} = \tilde{c} \in \tilde{B}$. Passing to the quotients by the π -actions we obtain the injective *assembly map*

$$A : X \times_B Y \rightarrow \tilde{X} \times_{\pi} \tilde{Y} ; (x, y) \mapsto [(x, \tilde{b}), (y, \tilde{b})]$$

with $\tilde{b} \in q_B^{-1}(p_X(x)) = q_B^{-1}(p_Y(y)) \subseteq \tilde{B}$. In particular, for $X = Y = B$ the assembly is just the diagonal map

$$A = \Delta : B \times_B B = B \rightarrow \tilde{B} \times_{\pi} \tilde{B} ; b \mapsto [\tilde{b}, \tilde{b}] .$$

The *coproduct* of fibrewise spaces X, Y is the disjoint union

$$X \sqcup_B Y = X \sqcup Y$$

with the topology characterized by the property that fibrewise maps $X \sqcup Y \rightarrow Z$ are in one-one correspondence with the fibrewise maps $X \rightarrow Z, Y \rightarrow Z$. The *quotient* fibrewise space of a closed subspace $A \subseteq X$ is

$$X/_B A = (X \sqcup_B B) / \sim$$

with \sim the equivalence relation generated by $x \in A, b \in B$ with $p_X(x) = b$.

A fibrewise space X is *pointed* if the projection $p_X : X \rightarrow B$ is equipped with a section $s_X : B \rightarrow X$. The *smash product* of pointed fibrewise spaces X, Y is the pointed fibrewise space

$$X \wedge_B Y = X \times_B Y /_B ((X \times_B s_Y(B)) \cup (s_X(B) \times_B Y)) .$$

In particular, for any spaces P, Q over B

$$(P \sqcup_B B) \wedge_B (Q \sqcup_B B) = (P \times_B Q) \sqcup_B B .$$

A fibrewise map $f : X \rightarrow Y$ over B induces a π -equivariant map

$$\tilde{f} : \tilde{X} \rightarrow \tilde{Y} ; (x, \tilde{b}) \rightarrow (f(x), \tilde{b}) .$$

Given a covering projection $q_B : \tilde{B} \rightarrow B$ with group of covering projections π and a pointed fibrewise space X over B define the pointed π -space \tilde{X}/\tilde{B} , with the action of π free away from the base point. Define the *assembly map* from the fibrewise homotopy set to the π -equivariant homotopy set

$$A : [X, Y]_B \rightarrow [\tilde{X}/\tilde{B}, \tilde{Y}/\tilde{B}]^\pi ; f \mapsto \tilde{f} .$$

The *suspension* of X is the pointed fibrewise space

$$\Sigma_B X = (B \times S^1) \wedge_B X .$$

The (disk, sphere)-bundle of a (finite-dimensional real) vector bundle ξ over a space M over B is a pair of spaces $(D(\xi), S(\xi))$ over B . The *fibrewise Thom space* of ξ is the fibrewise space over B is

$$T_B(\xi) = D(\xi)/_B S(\xi) .$$

(The fibrewise Thom space is denoted by M^ξ in [13] and [14]). For each $x \in B$ there is a base point $*_x \in T_B(\xi)$, so that $B \subset T_B(\xi)$, with

$$T_B(\xi)/B = D(\xi)/S(\xi) = T(\xi)$$

the ordinary Thom space of ξ . The effect on the fibrewise Thom space of adding the trivial line bundle ϵ to ξ is given by

$$T_B(\xi \oplus \epsilon) = \Sigma_B T_B(\xi) .$$

In particular

$$T_B(0) = M \sqcup_B B , T_B(\epsilon^k) = \Sigma_B^k(M \sqcup_B B) \quad (k \geq 0) .$$

If $q_B : \tilde{B} \rightarrow B$ is a covering projection with group of covering translations π the pullback $\tilde{\xi} = q_B^*(\xi)$ is a vector bundle over \tilde{B} such that

$$T_B(\tilde{\xi}) = T(\tilde{\xi})/\tilde{B}$$

is a semifree π -space with quotient

$$T_B(\tilde{\xi})/\pi = T_B(\xi) .$$

A *fibrewise stable map* $F : X \rightarrow Y$ of pointed fibrewise spaces is an equivalence class of fibrewise maps

$$F : \Sigma_B^k X \rightarrow \Sigma_B^k Y \quad (k \geq 0) .$$

The abelian group of fibrewise stable maps $X \rightarrow Y$ is written

$$\omega_B\{X; Y\} = \lim_{k \rightarrow \infty} [\Sigma_B^k X; \Sigma_B^k Y]_B .$$

The fibrewise geometric Hopf map is a function

$$\omega_B\{X; Y\} \rightarrow \omega_B\{X; S(\infty)^+ \wedge_{\mathbb{Z}_2} (Y \wedge_B Y)\} ; F \mapsto h(F)$$

(See [13, pp. 168–169, 306–308] for details.)

For any map $f : M \rightarrow N$ write the path space fibration as

$$f' : M' = \{(x, \alpha) \in M \times N^{[0,1]} \mid \alpha(0) = f(x)\} \rightarrow N ; (x, \alpha) \mapsto \alpha(1) .$$

The projection

$$p : M' \rightarrow M ; (x, \alpha) \mapsto x$$

is a homotopy equivalence such that $fp \simeq f'$.

Suppose given an embedding $f : M \hookrightarrow N \setminus \partial N$ of a closed m -dimensional manifold M in the interior of an n -dimensional manifold N with boundary ∂N (which may be empty). Let $\nu_f : M \rightarrow BO(n-m)$ be the normal bundle, and let $D(\nu_f) \hookrightarrow N$ be a tubular neighbourhood. Applying the standard Pontryagin-Thom construction gives a non-fibrewise Umkehr map

$$F : N/\partial N \rightarrow T(\nu_f) ; y \mapsto \begin{cases} y & \text{if } y \in D(\nu_f) \\ * & \text{if } y \in N \setminus D(\nu_f) \end{cases}$$

which we shall now extend to a fibrewise Umkehr map over N

$$F_N : N \cup_{\partial N} N \rightarrow T_N(p^*\nu_f)$$

constructed as follows. For $y \in D(\nu_f)$ in the fibre of $D(\nu_f)$ over $x \in M$, let $\alpha : [0, 1] \rightarrow N$ be the linear path from $f(x)$ to y in the fibre $D(\nu_f(x))$ (in N), and let $z \in T_N(p^*\nu_f)$ be the point over y given by y in the fibre of $p^*\nu_f$ over $(x, \alpha) \in M'$. Define a map

$$N \rightarrow T_N(p^*\nu_f) ; y \mapsto \begin{cases} z & \text{if } y \in D(\nu_f) \\ * & \text{if } y \in N \setminus D(\nu_f) . \end{cases}$$

Note that $\alpha(1) = y$, so that this is fibrewise over N . Extend this map on N to $N \cup_{\partial N} N$ by mapping the other (basepoint) copy of N to the basepoint in each fibre of $T_N(p^*\nu_f)$, to obtain the fibrewise Umkehr map $F_N : N \cup_{\partial N} N \rightarrow T_N(p^*\nu_f)$. Passing to the quotient by N recovers the non-fibrewise Umkehr map

$$F : N/\partial N = (N \cup_{\partial N} N)/N \rightarrow T_N(p^*\nu_f)/N = T(p^*\nu_f) \simeq T(\nu_f) .$$

Given a regular cover $q_N : \tilde{N} \rightarrow N$ with group of covering translations π (e.g. the universal cover for connected N , with $\pi = \pi_1(N)$) it is possible to lift f to a π -equivariant embedding

$$\tilde{f} : \tilde{M} = f^* \tilde{N} = \{(x, y) \in M \times \tilde{N} \mid f(x) = q_N(y) \in N\} \hookrightarrow \tilde{N} ; (x, y) \mapsto y$$

and the standard Pontryagin-Thom construction gives a π -equivariant Umkehr map

$$\tilde{F} : \tilde{N}/\partial\tilde{N} \rightarrow T(\nu_{\tilde{f}}) = T(q_M^* \nu_f)$$

with q_M the projection

$$q_M : \tilde{M} \rightarrow M ; (x, y) \mapsto x .$$

The assembly of the fibrewise Umkehr map F_N determines is π -equivariant Umkehr map \tilde{F}

$$A : [N \cup_{\partial N} N, T_N(p^* \nu_f)]_N \rightarrow [\tilde{N}/\partial\tilde{N}, T(q_M^* \nu_f)]^\pi ; F_N \mapsto \tilde{F} .$$

More precisely, lift the fibrewise map $F_N : N \cup_{\partial N} N \rightarrow T_N(p^* \nu_f)$ from N to \tilde{N} to get a π -equivariant map over \tilde{N}

$$\tilde{F} : \tilde{N} \cup_{\partial\tilde{N}} \tilde{N} \rightarrow T_{\tilde{N}}(\tilde{p}^* \nu_f) ,$$

where \tilde{p} is obtained by lifting p :

$$\begin{aligned} \tilde{p} : \tilde{M}' &= \{(x, \alpha, y) \in M \times N^{[0,1]} \times \tilde{N} \mid \alpha(0) = f(x), \alpha(1) = q_N(y)\} \\ &\rightarrow M ; (x, \alpha, y) \mapsto x . \end{aligned}$$

For $(x, \alpha, y) \in \tilde{M}'$, we have a unique path $\tilde{\alpha} : [0, 1] \rightarrow \tilde{N}$ lifting α with $\tilde{\alpha}(1) = y$. Then $\tilde{\alpha}(0) \in \tilde{N}$ projects to $f(x) \in N$. Hence $(x, \tilde{\alpha}(0)) \in \tilde{M}$. This allows us to define a map

$$r : \tilde{M}' \rightarrow \tilde{M} ; (x, \alpha, y) \mapsto (x, \tilde{\alpha}(0))$$

such that $\tilde{p} = q_M \circ r$. We obtain the π -equivariant map \tilde{g} by collapsing the basepoints \tilde{N}

$$\tilde{g} : \tilde{N}/\partial\tilde{N} = (\tilde{N} \cup_{\partial\tilde{N}} \tilde{N})/(\tilde{N} \times \{*\}) \rightarrow T_{\tilde{N}}(\tilde{p}^* \nu_f)/(\tilde{N} \times \{*\}) = T(q_M^* \nu_f) .$$

In particular, if $g = F_N : N \cup_{\partial N} N \rightarrow T_N(p^* \nu_f)$ then $\tilde{g} = \tilde{F} : \tilde{N}/\partial\tilde{N} \rightarrow T(q_M^* \nu_f)$. If $f : M \hookrightarrow N$ is a framed embedding, with $\nu_f = \epsilon^{n-m}$ the trivial $(n-m)$ -plane bundle, the fibrewise Umkehr is a map

$$F_N : N \cup_{\partial N} N \rightarrow T_N(p^* \nu_f) = \Sigma_N^{n-m}(M' \sqcup_N N)$$

and the π -equivariant Umkehr is a map

$$\tilde{F} : \tilde{N}/\widetilde{\partial N} \rightarrow T(q_M^* \nu_f) = \Sigma^{n-m}(\tilde{M}^+) .$$

More generally, suppose given an immersion $f : M \looparrowright N \setminus \partial N$ of a closed m -dimensional manifold in the interior of an n -dimensional manifold with boundary, with normal bundle $\nu_f : M \rightarrow BO(n-m)$. For sufficiently large $j \geq 0$ there exists a map $e : M \rightarrow \mathbb{R}^j$ such that

$$g = e \times f : M \rightarrow \mathbb{R}^j \times N ; x \mapsto (e(x), f(x))$$

is an embedding which is regular homotopic to the composite

$$M \looparrowright N \hookrightarrow \mathbb{R}^j \times N$$

of f and the embedding

$$N \hookrightarrow \mathbb{R}^j \times N ; x \mapsto (0, x)$$

with trivial normal bundle $\epsilon^j : N \rightarrow BO(j)$. The normal bundle of g is

$$\nu_g = \nu_f \oplus \epsilon^j : M \rightarrow BO(n-m+j) .$$

For a regular cover $q_N : \tilde{N} \rightarrow N$ with group of covering translations π g lifts to a π -equivariant embedding $\tilde{g} : M \hookrightarrow \mathbb{R}^j \times \tilde{N}$, and there is a π -equivariant Umkehr map

$$\tilde{F} : \Sigma^j(\tilde{N}/\widetilde{\partial N}) \rightarrow T(\nu_{\tilde{g}}) = \Sigma^j T(\nu_{\tilde{f}}) = \Sigma^j T(q_M^* \nu_f) .$$

The fibrewise Umkehr of g is

$$G_{\mathbb{R}^j \times N} : \mathbb{R}^j \times (N \cup_{\partial N} N) \rightarrow T_{\mathbb{R}^j \times N}((p')^*(\nu_{e \times f}))$$

with $p' : M'' \rightarrow M$ the projection of

$$M'' = \{(x, \alpha) \in M \times (\mathbb{R}^j \times N)^{[0,1]} \mid \alpha(0) = (e(x), f(x))\} .$$

It is possible to regard $G_{\mathbb{R}^j \times N}$ as a fibrewise stable map over N

$$F_N = G_{\mathbb{R}^j \times N} : \Sigma_N^j(N \cup_{\partial N} N) \rightarrow \Sigma_N^j T_N(p^* \nu_f) ,$$

representing an element

$$F_N \in \omega_N\{N \cup_{\partial N} N; T_N(p^* \nu_f)\}$$

such that

$$\omega_N\{N \cup_{\partial N} N; T_N(p^* \nu_f)\} \rightarrow \{\tilde{N}^+; T(\nu_{\tilde{f}})\}^\pi ; F_N \mapsto \tilde{F} .$$

The stable homotopy class of the fibrewise Umkehr map F depends only on the regular homotopy class of the immersion f .

Finally, suppose given a normal map $(f, b) : M \rightarrow X$ from an m -dimensional manifold M to an m -dimensional geometric Poincaré complex X . By the π - π theorem of Wall [68, Chapter 3], for sufficiently large $j \geq 0$ $(X \times D^j, X \times S^{j-1})$ is homotopy equivalent to an $(m+j)$ -dimensional manifold with boundary $(N, \partial N)$ and (f, b) can be approximated by a framed embedding $M \hookrightarrow N \setminus \partial N$. The fibrewise Umkehr map over N

$$N \cup_{\partial N} N \rightarrow \Sigma_N^j(M \sqcup_N N)$$

determines a fibrewise stable Umkehr map over X

$$F_X : \Sigma_X^j(X \sqcup_X X) \rightarrow \Sigma_X^j(M \sqcup_X X)$$

representing an element $F_X \in \omega_X\{X \sqcup_X X; M \sqcup_X X\}$. The fibrewise S -duality of Crabb and James [13, p.267] extends to Poincaré complexes, with $X \sqcup_X X$ the fibrewise dual over X of the fibrewise Thom space $T_X(\nu_X)$ of the Spivak normal fibration ν_X , and F_X is the fibrewise S -dual of $T_X(b) : T_X(\nu_M) \rightarrow T_X(\nu_X)$

$$F_X = T_X(b) \in \omega_X\{X \sqcup_X X; M \sqcup_X X\} = \omega_X\{T_X(\nu_M); T_X(\nu_X)\}.$$

The fibrewise Hopf invariant is a function

$$\begin{aligned} h(F_X) &\in \omega_X(X \sqcup_X X, S(\infty)^+ \wedge_{\mathbb{Z}_2} ((M \sqcup_X X) \wedge_X (M \sqcup_X X))) \\ &= \omega_X(X \sqcup_X X, (S(\infty) \times_{\mathbb{Z}_2} (M \times_X M)) \sqcup_X X). \end{aligned}$$

If \tilde{X} is the universal cover of X and $\tilde{M} = f^*\tilde{X}$ is the pullback cover of M then F_X induces the $\pi_1(X)$ -equivariant S -dual $\tilde{F} : \Sigma^j(\tilde{X}^+) \rightarrow \Sigma^j(\tilde{M}^+)$ of Ranicki [50], and the fibrewise Hopf invariant induces the $\pi_1(X)$ -equivariant Hopf invariant (inducing the quadratic construction on the chain level)

$$h(\tilde{F}) \in \{X^+, (S(\infty) \times_{\mathbb{Z}_2} (\tilde{M} \times_{\pi_1(X)} \tilde{M}))^+\}.$$

Appendix B

Notes on \mathbb{Z}_2 -equivariant bordism

B.1 Introduction

Let X be a (metrisable) \mathbb{Z}_2 -ANR, and let ξ and η be real \mathbb{Z}_2 -vector bundles over X . (We restrict the topology of X for the sake of precision, but this is not really necessary as we are always dealing with the system of maps from a compact \mathbb{Z}_2 -ENR to X .) The fibre dimensions of ξ and η are also allowed to vary over the components of X .

B.2 Framed bordism

Definition B.1. Let ζ_0 and ζ_1 be \mathbb{Z}_2 -vector bundles over a compact \mathbb{Z}_2 -ENR W . We define a *restricted stable isomorphism* $\zeta_0 \cong \zeta_1$ to be an equivalence class of \mathbb{Z}_2 -vector bundle isomorphisms $\zeta_0 \oplus \mathbb{R}^i \cong \zeta_1 \oplus \mathbb{R}^i$, the equivalence being generated by homotopy and stabilization with respect to i . It will sometimes be convenient to think of a restricted stable isomorphism as given by an isomorphism $\zeta_0 \oplus V_0 \cong \zeta_1 \oplus V_0$ for some finite-dimensional \mathbb{R} -vector space V_0 on which \mathbb{Z}_2 acts trivially.

We recall that a *stable isomorphism* from ζ_0 to ζ_1 is an equivalence class of vector bundle isomorphisms $\zeta_0 \oplus V \cong \zeta_1 \oplus V$, where V is any finite dimensional real \mathbb{Z}_2 -module. A restricted stable isomorphism thus determines a stable isomorphism. The restricted stable automorphisms of a \mathbb{Z}_2 -vector bundle correspond to elements of $KO^{-1}(W/\mathbb{Z}_2)$, while the stable automorphisms correspond to elements of $KO_{\mathbb{Z}_2}^{-1}(W)$. In general, different restricted

stable isomorphisms may give the same stable isomorphism, but not if the action of \mathbb{Z}_2 on W is free.

Lemma B.2. *Suppose that W is a compact free \mathbb{Z}_2 -ENR. Then every stable isomorphism $\zeta_0 \cong \zeta_1$ arises from a unique restricted stable isomorphism.*

Proof. For an isomorphism $\zeta_0 \oplus V \cong \zeta_1 \oplus V$, where V is a \mathbb{Z}_2 -module corresponds to an isomorphism over W/\mathbb{Z}_2 and $W \times_{\mathbb{Z}_2} V$ is a direct summand of a trivial bundle.

Definition B.3. We define the *restricted bordism group* $\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta)$ to be the bordism group of closed \mathbb{Z}_2 -manifolds M with a \mathbb{Z}_2 -map $f : M \rightarrow X$ and a restricted stable isomorphism $a : \tau M \oplus f^* \xi \cong f^* \eta$, specified by a vector bundle isomorphism $\tau M \oplus f^* \xi \oplus V_0 \cong f^* \eta \oplus V_0$ for some vector space V_0 with trivial \mathbb{Z}_2 -action. Bordisms are required to be restricted, too, that is, (M, f, a) is null-bordant if there is a manifold W with boundary $\partial W = M$, a map $g : W \rightarrow X$ extending f and a restricted stable isomorphism $b : \tau W \oplus g^* \xi \cong \mathbb{R} \oplus g^* \eta$ extending a . The negative of the class (M, f, a) is given by (M, f, a') where $a' : \tau M \oplus f^* \xi \oplus V_0 \oplus \mathbb{R} \cong f^* \eta \oplus V_0 \oplus \mathbb{R}$ is $a \oplus (-1)$.

Note that the manifold M may have components of different dimension. Here we do not use the notation ‘ $\xi - \eta$ ’ because the group does not, in general, depend only on the virtual \mathbb{Z}_2 -bundle.

Up to homotopy we may assume that any finite-dimensional real \mathbb{Z}_2 -vector bundle is equipped with an equivariant positive-definite inner product. We write $O_X(\xi, \eta)$ for the Stiefel bundle with fibre at $x \in X$ the Stiefel manifold $O(\xi_x, \eta_x)$ of linear isometries $\xi_x \hookrightarrow \eta_x$. It is a \mathbb{Z}_2 -space.

Definition B.4. Let $F_X^{\text{res}}(\xi, \eta)$ be the infinite Stiefel bundle

$$\varinjlim_{k \geq 0} \varinjlim_{i \geq 0} O_X(\xi \oplus \mathbb{R}^i, \eta \oplus \mathbb{R}^i \oplus \mathbb{R}^k)$$

over X , where the maps

$$O_X(\xi \oplus \mathbb{R}^i, \eta \oplus \mathbb{R}^i \oplus \mathbb{R}^k) \rightarrow O_X(\xi \oplus \mathbb{R}^{i+1}, \eta \oplus \mathbb{R}^{i+1} \oplus \mathbb{R}^k)$$

are given by the direct sum with $1 : \mathbb{R} \rightarrow \mathbb{R}$ in the decomposition $\mathbb{R}^{i+1} = \mathbb{R}^i \oplus \mathbb{R}$ and the maps

$$O_X(\xi \oplus \mathbb{R}^i, \eta \oplus \mathbb{R}^i \oplus \mathbb{R}^k) \rightarrow O_X(\xi \oplus \mathbb{R}^i, \eta \oplus \mathbb{R}^i \oplus \mathbb{R}^{k+1})$$

are given by including \mathbb{R}^k in \mathbb{R}^{k+1} . The group \mathbb{Z}_2 acts on $F_X^{\text{res}}(\xi, \eta)$.

Remark B.5. The projection $F_X^{\text{res}}(\xi, \eta) \rightarrow X$ is a non-equivariant homotopy equivalence. The fibre is contractible, because of the limit over the inclusions $\mathbb{R}^k \hookrightarrow \mathbb{R}^{k+1}$.

Remark B.6. Suppose that $h : X' \rightarrow X$ is a \mathbb{Z}_2 -map. Let ξ' and η' be the pullbacks of ξ and η to X' . Then $F_{X'}^{\text{res}}(\xi', \eta')$ is the pullback of $F_X^{\text{res}}(\xi, \eta)$.

Definition B.7. A Pontryagin-Thom map

$$\varphi : \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \rightarrow \omega_0^{\mathbb{Z}_2}(F_X^{\text{res}}(\xi, \eta); \xi - \eta).$$

is defined as follows. From $f : M \rightarrow X$ and a restricted stable isomorphism $a : \tau M \oplus f^*\xi \cong f^*\eta$ given by a vector bundle isomorphism $\tau M \oplus f^*\xi \oplus \mathbb{R}^i \cong f^*\eta \oplus \mathbb{R}^i$ for some i , including $f^*\xi \oplus \mathbb{R}^i$ in $\tau M \oplus f^*\xi \oplus \mathbb{R}^i$ we get a lift of f to a map $\tilde{f} : M \rightarrow \text{O}_X(\xi \oplus \mathbb{R}^i, \eta \oplus \mathbb{R}^i) \rightarrow F_X^{\text{res}}(\xi, \eta)$. The standard Pontryagin-Thom construction applied to \tilde{f} gives a class

$$\varphi(M, f, a) \in \omega_0^G(F_X^{\text{res}}(\xi, \eta); \xi - \eta).$$

We must check that this class is 0 if (M, f, a) is the boundary of (W, g, b) with a restricted stable isomorphism $b : \tau W \oplus g^*\xi \cong \mathbb{R} \oplus g^*\eta$ extending a on $M = \partial W$. From a bundle isomorphism $\tau W \oplus g^*\xi \oplus \mathbb{R}^i \cong \mathbb{R} \oplus g^*\eta \oplus \mathbb{R}^i$ we get a lift of g to $\tilde{g} : W \rightarrow \text{O}_X(\xi \oplus \mathbb{R}^i, \mathbb{R} \oplus \eta \oplus \mathbb{R}^i) \rightarrow F_X^{\text{res}}(\xi, \eta)$. Hence the Pontryagin-Thom construction on an associated \tilde{f} gives 0. (The check explains the definition of F_X^{res} .)

The main result of this section is

Theorem B.8. *The Pontryagin-Thom map is an isomorphism*

$$\varphi : \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \rightarrow \omega_0^{\mathbb{Z}_2}(F_X^{\text{res}}(\xi, \eta); \xi - \eta).$$

It will be proved by comparing a Conner-Floyd sequence with the localization exact sequence in stable homotopy.

Definition B.9. We define the *free bordism group* $\Omega_0^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta)$ to be the bordism group of closed free \mathbb{Z}_2 -manifolds M with a \mathbb{Z}_2 -map $f : M \rightarrow X$ and a stable \mathbb{Z}_2 -isomorphism $a : \tau M \oplus f^*\xi \cong f^*\eta$. (Bounding manifolds W are also required to have a free action of \mathbb{Z}_2 .)

Definition B.10. We have a free Pontryagin-Thom map

$$\varphi^{\text{free}} : \Omega_0^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta) \rightarrow \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times X; \xi - \eta)$$

given by the classical construction using the product $M \rightarrow E\mathbb{Z}_2 \times X$ of the classifying map of the free involution with f .

Proposition B.11. *Write $\bar{X} = E\mathbb{Z}_2 \times_{\mathbb{Z}_2} X$ and let $\bar{\xi}$ and $\bar{\eta}$ be the induced bundles over \bar{X} . Then we have isomorphisms*

$$\begin{array}{ccc} \Omega_0(\bar{X}; \bar{\xi}, \bar{\eta}) & \cong & \Omega_0^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta) \\ \downarrow & & \downarrow \varphi^{\text{free}} \\ \omega_0(\bar{X}; \bar{\xi} - \bar{\eta}) & \cong & \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times X; \xi - \eta) \end{array}$$

through which φ^{free} corresponds to the classical Pontryagin-Thom map.

Proof. This reduces, by naturality with respect to inclusions, to showing that, for a free \mathbb{Z}_2 -manifold M , the equivariant fundamental class $[M] \in \omega_0^{\mathbb{Z}_2}(M; -\tau M)$ corresponds to $[\bar{M}] \in \omega_0(\bar{M}; -\tau \bar{M})$ under the isomorphism: $\omega_0(\bar{M}; -\tau \bar{M}) \cong \omega_0^{\mathbb{Z}_2}(M; -\tau M) \cong \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times M; -\tau M)$. But the isomorphism is given by the equivariant Umkehr map $\pi^! : \omega_0^{\mathbb{Z}_2}(\bar{M}; -\tau \bar{M}) \rightarrow \omega_0^{\mathbb{Z}_2}(M; -\tau M)$ for the projection $\pi : M \rightarrow \bar{M}$, which takes the fundamental class of \bar{M} with the trivial involution to the fundamental class of M .

Corollary B.12. *The free Pontryagin-Thom map φ^{free} is an isomorphism.* \square

Definition B.13. Using the fact that stable isomorphisms over free \mathbb{Z}_2 -spaces correspond to restricted stable isomorphisms we get a homomorphism

$$\gamma : \Omega_0^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta) \rightarrow \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta).$$

Lemma B.14. *We have a commutative diagram*

$$\begin{array}{ccc} \Omega_0^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta) & \xrightarrow{\gamma} & \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \\ \cong \downarrow & & \downarrow \varphi \\ \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times F_X^{\text{res}}(\xi, \eta); \xi - \eta) & \rightarrow & \omega_0^{\mathbb{Z}_2}(F_X^{\text{res}}(\xi, \eta); \xi - \eta). \end{array}$$

Proof. We use the non-equivariant homotopy equivalence $F_X^{\text{res}}(\xi, \eta) \rightarrow X$ to get an isomorphism

$$\omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times F_X^{\text{res}}(\xi, \eta); \xi - \eta) \rightarrow \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times X; \xi - \eta).$$

The assertion is then clear from the definition of the Pontryagin-Thom maps.

Lemma B.15. *Let $\xi_{\mathbb{Z}_2}$ be the orthogonal complement of $\xi^{\mathbb{Z}_2}$ in $\xi|X^{\mathbb{Z}_2}$, and similarly for $\eta_{\mathbb{Z}_2}$. The fixed subspace $F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}$ is homotopy equivalent to the space of \mathbb{Z}_2 -monomorphisms $O_{X^{\mathbb{Z}_2}}^{\mathbb{Z}_2}(\xi_{\mathbb{Z}_2}, \eta_{\mathbb{Z}_2})$.*

Proof. The fixed subspace is

$$\left(\lim_{\substack{\rightarrow \\ k}} \lim_{\substack{\rightarrow \\ i}} O_{X^{\mathbb{Z}_2}}(\xi^{\mathbb{Z}_2} \oplus \mathbb{R}^i, \eta^{\mathbb{Z}_2} \oplus \mathbb{R}^i \oplus \mathbb{R}^k) \right) \times_{X^{\mathbb{Z}_2}} O_{X^{\mathbb{Z}_2}}(\xi_{\mathbb{Z}_2}, \eta_{\mathbb{Z}_2}).$$

Definition B.16. A *fixed point homomorphism*

$$\rho : \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \rightarrow \Omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2}, \eta^{\mathbb{Z}_2})$$

is defined as follows. Given a \mathbb{Z}_2 -manifold M a map $f : M \rightarrow X$ and a restricted stable isomorphism $a : \tau M \oplus f^* \xi \cong f^* \eta$, we form the fixed submanifold $M^{\mathbb{Z}_2}$. We have a map $f^{\mathbb{Z}_2} : M^{\mathbb{Z}_2} \rightarrow X^{\mathbb{Z}_2}$ and a stable isomorphism $\tau M^{\mathbb{Z}_2} \oplus (f^{\mathbb{Z}_2})^* \xi^{\mathbb{Z}_2} \cong (f^{\mathbb{Z}_2})^* \eta^{\mathbb{Z}_2}$. The group \mathbb{Z}_2 acts on $\xi_{\mathbb{Z}_2}$ and $\eta_{\mathbb{Z}_2}$ as ± 1 . The normal bundle ν of $M^{\mathbb{Z}_2} \hookrightarrow M$ is a \mathbb{Z}_2 -bundle, determined by an isomorphism $\nu \oplus (f^{\mathbb{Z}_2})^* \xi_{\mathbb{Z}_2} \cong (f^{\mathbb{Z}_2})^* \eta_{\mathbb{Z}_2}$. This gives us a \mathbb{Z}_2 -map $X^{\mathbb{Z}_2} \rightarrow O_{X^{\mathbb{Z}_2}}(\xi_{\mathbb{Z}_2}, \eta_{\mathbb{Z}_2})$, that is, a map $X^{\mathbb{Z}_2} \rightarrow O_{X^{\mathbb{Z}_2}}^{\mathbb{Z}_2}(\xi_{\mathbb{Z}_2}, \eta_{\mathbb{Z}_2})$. The manifold $M^{\mathbb{Z}_2}$ with the extra data represents the image of (M, f, a) under ρ .

Lemma B.17. *We have a commutative diagram*

$$\begin{array}{ccc} \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) & \xrightarrow{\rho} & \Omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2}, \eta^{\mathbb{Z}_2}) \\ \varphi \downarrow & & \downarrow \cong \\ \omega_0^{\mathbb{Z}_2}(F_X^{\text{res}}(\xi, \eta); \xi - \eta) & \xrightarrow{\rho} & \omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2}) \end{array}$$

Proof. The right-hand map is the non-equivariant Pontryagin-Thom homomorphism, so an isomorphism. Commutativity follows directly from the construction of the Pontryagin-Thom maps.

Definition B.18. We introduce groups $\Omega_*^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta)$ for $* \in \mathbb{Z}$ so that

$$\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \mathbb{R}^m \oplus \xi, \mathbb{R}^n \oplus \eta) = \Omega_{n-m}^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta)$$

for $m, n \geq 0$. In particular,

$$\Omega_{-1}^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) = \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \mathbb{R} \oplus \xi, \oplus \eta).$$

Definition B.19. A *boundary map*

$$\partial : \Omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2}, \eta^{\mathbb{Z}_2}) \rightarrow \Omega_{-1}^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta)$$

is constructed as follows. Consider a manifold N with a map $g : N \rightarrow X^{\mathbb{Z}_2}$, a stable isomorphism $\tau N \oplus g^* \xi^{\mathbb{Z}_2} \cong g^* \eta^{\mathbb{Z}_2}$, given by a bundle isomorphism $\tau N \oplus g^* \xi^{\mathbb{Z}_2} \oplus V_0 \cong g^* \eta^{\mathbb{Z}_2} \oplus V_0$, and a bundle \mathbb{Z}_2 -monomorphism $g^* \xi_{\mathbb{Z}_2} \hookrightarrow g^* \eta_{\mathbb{Z}_2}$ with complementary bundle ν . Form the free \mathbb{Z}_2 -manifold $S(\nu)$. It is equipped with a \mathbb{Z}_2 -map $h : S(\nu) \rightarrow N \rightarrow X^{\mathbb{Z}_2} \rightarrow X$. We have a \mathbb{Z}_2 -isomorphism $\mathbb{R} \oplus \tau S(\nu) \cong \tau N \oplus \nu$, so $\mathbb{R} \oplus \tau S(\nu) \oplus g^* \xi \oplus V_0 \cong g^* \eta \oplus V_0$. This extends to a \mathbb{Z}_2 -isomorphism $\mathbb{R} \oplus \tau(S(\nu)) \oplus h^* \xi \oplus V_0 \cong h^* \eta \oplus V_0$.

Lemma B.20. *We have a commutative diagram*

$$\begin{array}{ccc} \Omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2}, \eta^{\mathbb{Z}_2}) & \xrightarrow{\partial} & \Omega_{-1}^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta) \\ \varphi \downarrow & & \downarrow \cong \\ \omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2}) & \xrightarrow{\partial} & \omega_{-1}^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times F_X^{\text{res}}(\xi, \eta); \xi - \eta) \\ \downarrow & & \downarrow \cong \\ \omega_0(X^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2}) & \xrightarrow{\partial} & \omega_{-1}^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times X; \xi - \eta) \end{array}$$

Proof. The two lower maps ∂ are the boundary maps in the localization exact sequence. The right-hand maps are defined as before using the non-equivariant equivalence from $F_X^{\text{res}}(\xi, \eta)$ to X .

We look at an element of $\Omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2}, \eta^{\mathbb{Z}_2})$ represented by a manifold N with a map $g : N \rightarrow X^{\mathbb{Z}_2}$, a stable isomorphism $\tau N \oplus g^* \xi^{\mathbb{Z}_2} \cong g^* \eta^{\mathbb{Z}_2}$, given by a bundle isomorphism $\tau N \oplus g^* \xi^{\mathbb{Z}_2} \oplus V_0 \cong g^* \eta^{\mathbb{Z}_2} \oplus V_0$, and a bundle \mathbb{Z}_2 -monomorphism $g^* \xi_{\mathbb{Z}_2} \hookrightarrow g^* \eta_{\mathbb{Z}_2}$ with complementary bundle ν . By the obvious naturality under inclusions we may assume that $X = N = X^{\mathbb{Z}_2}$, g is the identity map, $\xi = 0$ and $\eta = \tau N \oplus \nu$, so that $\eta^{\mathbb{Z}_2} = \tau N$ and $\eta_{\mathbb{Z}_2} = \nu$.

The image of the element in $\omega_0(X^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2}) = \omega_0(N; -\tau N)$ is the fundamental class $[N]$ of N .

Its image in $\Omega_{-1}^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta) = \Omega_{-1}^{\mathbb{Z}_2, \text{free}}(N; 0, \tau N \oplus \nu)$ is represented by the sphere bundle $S(\nu) \rightarrow N$ with $\mathbb{R} \oplus \tau S(\nu) = \tau N \oplus \nu$. This maps by the Pontryagin-Thom construction to the \mathbb{Z}_2 -fundamental class $[S(\nu)] \in \omega_0^{\mathbb{Z}_2}(S(\nu); -\tau S(\nu)) = \omega_{-1}^{\mathbb{Z}_2}(S(\nu); -\tau N - \nu)$ and then by the classifying map $S(\nu) \rightarrow E\mathbb{Z}_2 \times N$ to the group $\omega_{-1}^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times N; -\tau N - \nu)$.

From the construction of the localization exact sequence we have a commutative square

$$\begin{array}{ccc} \omega_0^{\mathbb{Z}_2}(N; -\tau N) & \xrightarrow{\partial} & \omega_{-1}^{\mathbb{Z}_2}(S(\nu); -\tau N - \nu) \\ \rho \downarrow & & \downarrow \\ \Omega_0(N; -\tau N) & \xrightarrow{\partial} & \omega_{-1}^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times N; -\tau N - \nu) \end{array}$$

The assertion now follows from the fact that the equivariant fundamental class $[N]$ in the top line is mapped by ∂ to the fundamental class $[S(\nu)]$, because ∂ in the exact sequence of the pair $(D(\nu), S(\nu))$ is the Umkehr homomorphism of the projection $S(\nu) \rightarrow N$.

Proposition B.21. *The Conner-Floyd sequence:*

$$\dots \xrightarrow{\partial} \Omega_0^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta) \xrightarrow{\gamma} \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \xrightarrow{\rho} \Omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2}, \eta^{\mathbb{Z}_2}) \xrightarrow{\partial} \dots$$

is exact.

Proof. We use the standard surgery arguments.

It is clear that $\rho \circ \gamma = 0$. We show that $\ker \rho \subseteq \text{im } \gamma$. Suppose that a class represented by $f : M \rightarrow X$ and $a : \tau M \oplus f^* \xi \cong f^* \eta$ lies in the kernel of ρ . We have a manifold W with $\partial W = X^{\mathbb{Z}_2}$, a map $g : W \rightarrow X^{\mathbb{Z}_2}$ and a stable isomorphism $b : \tau W \oplus g^* \xi^{\mathbb{Z}_2} \cong \mathbb{R} \oplus g^* \eta^{\mathbb{Z}_2}$ extending a . Further, we have a vector bundle $\hat{\nu}$ over W extending the normal bundle ν over $M^{\mathbb{Z}_2}$ and an isomorphism $\hat{\nu} \oplus g^* \xi_{\mathbb{Z}_2} \cong g^* \eta_{\mathbb{Z}_2}$.

Take an equivariant tubular neighbourhood $D(\nu) \hookrightarrow M$ of $M^{\mathbb{Z}_2}$ in M . We construct a free \mathbb{Z}_2 -manifold $M' = (M - B(\nu)) \cup_{S(\nu)} S(\hat{\nu})$ cobordant to M . From f and g we get a map $f' : M' \rightarrow X$. From a and b we get a restricted stable isomorphism $a' : \tau M' \oplus (f')^* \xi \cong (f')^* \eta$. Notice that $\mathbb{R} \oplus \tau S(\hat{\nu}) = \tau W \oplus \hat{\nu}$, so that $\mathbb{R} \oplus \tau S(\hat{\nu}) \oplus g^* \xi^{\mathbb{Z}_2} \oplus g^* \xi_{\mathbb{Z}_2} \cong \tau W \oplus \hat{\nu} \oplus g^* \xi^{\mathbb{Z}_2} \oplus g^* \xi_{\mathbb{Z}_2} \cong \mathbb{R} \oplus g^* \eta^{\mathbb{Z}_2} \oplus g^* \eta_{\mathbb{Z}_2}$. This gives a lift to $\Omega_0^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta)$ and shows that $\ker \rho \subseteq \text{im } \gamma$.

The composition $\gamma \circ \partial$ is zero, because $S(\nu)$ is the boundary of $D(\nu)$. To see that $\ker \gamma \subseteq \text{im } \partial$, suppose that a manifold S represents a class in $\Omega_{-1}^{\mathbb{Z}_2, \text{free}}(X; \xi, \eta)$ that is the boundary of a manifold M . Then the fixed-point construction applied to M produces an element of $\Omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2}, \eta^{\mathbb{Z}_2})$ mapping to S under ∂ . (Notice that there are no fixed points on the boundary S of M .)

This same construction with $S = \emptyset$ shows that $\partial \circ \rho = 0$. We must check finally that $\ker \partial \subseteq \text{im } \rho$. Given N representing a class in $\ker \partial$, so that $S(\nu)$ is the boundary of a manifold M_0 , we form the closed manifold $M = M_0 \cup D(\nu)$ by gluing along $\partial M_0 = S(\nu) = \partial D(\nu)$. Then M (with the associated structure) gives an element of $\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta)$ lifting the class of N . (In more detail, we have $g : N \rightarrow X^{\mathbb{Z}_2}$, $\tau N \oplus g^* \xi^{\mathbb{Z}_2} \oplus V_0 \cong g^* \eta^{\mathbb{Z}_2} \oplus V_0$ and $\nu \oplus g^* \xi_{\mathbb{Z}_2} \cong g^* \eta_{\mathbb{Z}_2}$. We have $h_0 : M_0 \rightarrow X$ and $\tau M_0 \oplus h_0^* \xi \oplus V_0 \cong h_0^* \eta \oplus V_0$, for V_0 of sufficiently high dimension. We get a map $f : M \rightarrow X$ from h on $S(\nu)$ and h_0 on M_0 and a bundle isomorphism $\tau M \oplus f^* \xi \oplus V_0 \cong f^* \eta \oplus V_0$.)

Proof (Proof of the main theorem). This is proved by comparing the Conner-Floyd exact sequences with the localization exact sequence:

$$\begin{aligned} \cdots \xrightarrow{\partial} \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times F_X^{\text{res}}(\xi, \eta); \xi - \eta) &\xrightarrow{\gamma} \omega_0^{\mathbb{Z}_2}(F_X^{\text{res}}(\xi, \eta); \xi - \eta) \\ &\xrightarrow{\rho} \omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2}) \xrightarrow{\partial} \cdots \end{aligned}$$

using the lemmas to check commutativity of the various squares.

We consider some cases in which the Pontryagin-Thom homomorphism

$$\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \rightarrow \omega_0^{\mathbb{Z}_2}(X; \xi - \eta)$$

is an isomorphism.

Remark B.22. Suppose that either (i) $X^{\mathbb{Z}_2}$ is empty or (ii) $X^{\mathbb{Z}_2} \neq \emptyset$ but $\dim \xi_{\mathbb{Z}_2} > \dim \eta_{\mathbb{Z}_2}$. Then the fixed subspaces $F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}$ and $F_X(\xi, \eta)^{\mathbb{Z}_2}$ are empty and

$$\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) = \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times X; \xi - \eta).$$

It is clear that, for any manifold M representing a bordism class, we must have $M^{\mathbb{Z}_2} = \emptyset$, so that the action of \mathbb{Z}_2 is free.

Proposition B.23. *Suppose that $X^{\mathbb{Z}_2}$ is non-empty but $\xi_{\mathbb{Z}_2} = 0$. Then*

$$\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \cong \omega_0^{\mathbb{Z}_2}(X; \xi - \eta)$$

Proof. Since $O_{X^{\mathbb{Z}_2}}^{\mathbb{Z}_2}(\xi_{\mathbb{Z}_2}, \eta_{\mathbb{Z}_2}) = X^{\mathbb{Z}_2}$, the projection $F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2} \rightarrow X^{\mathbb{Z}_2}$ is a homotopy equivalence.

Proposition B.24. *Assuming that $X^{\mathbb{Z}_2}$ is non-empty, suppose that $\dim \eta_{\mathbb{Z}_2} - \dim \xi_{\mathbb{Z}_2} > \dim \eta^{\mathbb{Z}_2} - \dim \xi^{\mathbb{Z}_2} + 1$. Then we have an isomorphism*

$$\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \rightarrow \omega_0^{\mathbb{Z}_2}(X; \xi - \eta).$$

Proof. Suppose that $\dim \eta_{\mathbb{Z}_2} - \dim \xi_{\mathbb{Z}_2} = r \geq 1$. Then the fibre of the projection $F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2} \rightarrow X^{\mathbb{Z}_2}$, homotopy equivalent to the Stiefel manifold $O(\mathbb{R}^k, \mathbb{R}^{k+r})$ for some k , has $\pi_i = 0$ for $i < r$. So

$$\omega_0(F_X^{\text{res}}(\xi, \eta)^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2}) \rightarrow \omega_0(X^{\mathbb{Z}_2}; \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2})$$

is an isomorphism if $\dim \eta^{\mathbb{Z}_2} - \dim \xi^{\mathbb{Z}_2} < r$.

The stable homotopy group $\omega_0^{\mathbb{Z}_2}(X; \xi - \eta)$ can be represented as a direct limit of restricted bordism groups.

Suppose that ζ is a real \mathbb{Z}_2 -vector bundle over X . We define a homomorphism

$$\Sigma_\zeta : \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \rightarrow \Omega_0^{\mathbb{Z}_2, \text{res}}(S(\mathbb{R} \oplus \zeta); \xi, \eta \oplus \zeta)$$

by sending the class of a manifold M , $f : M \rightarrow X$ and a restricted stable isomorphism $a : \tau M \oplus f^*\xi \rightarrow f^*\eta$ to the manifold $S = S(\mathbb{R} \oplus f^*\zeta)$ (with a choice of smooth structure on $f^*\zeta$ – different choices will give isomorphic smooth vector bundles) with the map $S \rightarrow M \rightarrow X$ given by the projection composed with f and the restricted stable isomorphism

$$\tau S \oplus \mathbb{R} \oplus f^*\xi \cong \tau M \oplus f^*\zeta \oplus \mathbb{R} \cong f^*\eta \oplus f^*\zeta \oplus \mathbb{R}.$$

Lemma B.25. *There is a commutative diagram*

$$\begin{array}{ccc} \Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) & \xrightarrow{\Sigma_\zeta} & \Omega_0^{\mathbb{Z}_2, \text{res}}(S(\mathbb{R} \oplus \zeta); \xi, \eta \oplus \zeta) \\ \downarrow & & \downarrow \\ \omega_0^{\mathbb{Z}_2}(X; \xi - \eta) & \xrightarrow{\sigma_\zeta} & \omega_0^{\mathbb{Z}_2}(X; \xi - \eta) \oplus \omega_0^{\mathbb{Z}_2}(X; \xi - \eta - \zeta), \end{array}$$

in which the vertical maps are Pontryagin-Thom homomorphisms and σ_ζ is the inclusion of the first factor.

Proof. The homomorphism σ_ζ is the boundary ∂ in the exact sequence of the pair $(D(\mathbb{R} \oplus \zeta), S(\mathbb{R} \oplus \zeta))$. The group $\omega_0^{\mathbb{Z}_2}(S(\mathbb{R} \oplus \zeta); \xi - \eta - \zeta)$ is split by the section $(-1, 0)$ of $X \rightarrow S(\mathbb{R} \oplus \zeta)$.

If $\zeta = L^k$ and k is sufficiently large, the map

$$\Omega_0^{\mathbb{Z}_2, \text{res}}(S(\mathbb{R} \oplus L^k); \xi, \eta \oplus L^k) \rightarrow \omega_0^{\mathbb{Z}_2}(X; \xi - \eta) \oplus \omega_0^{\mathbb{Z}_2}(X; \xi - \eta - L^k)$$

is an isomorphism. By introducing relative bordism groups we could represent $\omega_0^{\mathbb{Z}_2}(X; \xi - \eta)$ exactly as the restricted bordism group of the pair $(S(\mathbb{R} \oplus L^k), X)$.

Proposition B.26. *Suppose that $\xi_{\mathbb{Z}_2} = \eta_{\mathbb{Z}_2}$. Then we have a direct sum decomposition*

$$\Omega_0^{\mathbb{Z}_2, \text{res}}(X; \xi, \eta) \cong \omega_0^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times X; \xi - \eta) \oplus \omega_0(\mathcal{O}_{X^{\mathbb{Z}_2}}(\xi_{\mathbb{Z}_2}); \xi^{\mathbb{Z}_2} - \eta^{\mathbb{Z}_2}).$$

Proof. The stable homotopy localization sequence splits.

When ξ and η are trivial, this reduces to

$$\Omega_m^{\mathbb{Z}_2, \text{res}}(X; L^k, L^k) \cong \omega_m(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} X) \oplus \omega_m(X^{\mathbb{Z}_2} \times O(k)).$$

In particular,

$$\Omega_m^{\mathbb{Z}_2, \text{res}}(X; 0, 0) \cong \omega_m(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} X) \oplus \omega_m(X^{\mathbb{Z}_2}).$$

Indeed, for any representing manifold M , the fixed submanifold $M^{\mathbb{Z}_2}$ has dimension m and M is the disjoint union of $M^{\mathbb{Z}_2}$ and a closed free \mathbb{Z}_2 -manifold.

Definition B.27. We defined the (unrestricted) *bordism group* $\Omega_0^{\mathbb{Z}_2}(X; \xi, \eta)$ to be the bordism group of closed \mathbb{Z}_2 -manifolds M with a \mathbb{Z}_2 -map $f : M \rightarrow X$ and a stable isomorphism $a : \tau M \oplus f^* \xi \cong f^* \eta$, specified by a vector bundle isomorphism $\tau M \oplus f^* \xi \oplus V \cong f^* \eta \oplus V$ for some \mathbb{Z}_2 -module V .

Thus

$$\Omega_0^{\mathbb{Z}_2}(X; \xi, \eta) = \varinjlim_j \Omega^{\mathbb{Z}_2, \text{res}}(X; L^j \oplus \xi, L^j \oplus \eta).$$

Definition B.28. Let $F_X(\xi, \eta)$ be the \mathbb{Z}_2 -bundle

$$\varinjlim_{j \geq 0} F_X^{\text{res}}(L^j \oplus \xi, L^j \oplus \eta).$$

We have an equivalence between $F_X(\xi, \eta)$ and $F_X(V \oplus \xi, V \oplus \eta)$ for any \mathbb{Z}_2 -module V . Thus $F_X(\xi, \eta)$ depends only on the virtual bundle $\xi - \eta$.

Our results on the restricted bordism groups now describe $\Omega_0^{\mathbb{Z}_2}(X; \xi, \eta)$ as a stable homotopy group.

Corollary B.29. *The Pontryagin-Thom map is an isomorphism*

$$\varphi : \Omega_0^{\mathbb{Z}_2}(X; \xi, \eta) \rightarrow \omega_0^{\mathbb{Z}_2}(F_X(\xi, \eta); \xi - \eta).$$

Corollary B.30. *The fixed subspace $F_X(\xi, \eta)^{\mathbb{Z}_2}$ is homotopy equivalent to the infinite Stiefel bundle*

$$\varinjlim_{j \geq 0} O_{X^{\mathbb{Z}_2}}^{\mathbb{Z}_2}(L^j \oplus \xi_{\mathbb{Z}_2}, L^j \oplus \eta_{\mathbb{Z}_2}).$$

It depends only on the virtual bundle $\xi_{\mathbb{Z}_2} - \eta_{\mathbb{Z}_2}$.

Remark B.31. Suppose that $\xi_{\mathbb{Z}_2} - \eta_{\mathbb{Z}_2} = 0$. Then $F_X(\xi, \eta)^{\mathbb{Z}_2}$ is homotopy equivalent to $X^{\mathbb{Z}_2} \times O(\infty)$.

Proposition B.32. *The bordism group $\Omega_m^{\mathbb{Z}_2}(X)$, that is, $\Omega_0^{\mathbb{Z}_2}(X; 0, \mathbb{R}^m)$, splits as a direct sum*

$$\Omega_m^{\mathbb{Z}_2}(X) \cong \omega_m(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} X) \oplus \omega_m(X^{\mathbb{Z}_2} \times O(\infty)).$$

Proof. This follows from the preceding Remark and the splitting in stable homotopy.

Remark B.33. Suppose that $\eta = X \times U$ is trivial. Then a *normal map* given by a map $f : M \rightarrow X$, an embedding $M \subseteq U \oplus V$, for some \mathbb{Z}_2 -module V , and an equivariant vector bundle isomorphism $\nu_M \rightarrow f^*\xi \oplus V$ determines an element of $\Omega_0^{\mathbb{Z}_2}(X; \xi, W)$. For it determines an isomorphism $\tau M \oplus f^*\xi \oplus V \rightarrow U \oplus V$.

B.3 Unoriented bordism

We write the infinite Grassmannian of m -dimensional real subspaces as

$$G_m(\infty \oplus \infty L) = \varinjlim_{i,j \geq 0} G_m(\mathbb{R}^i \oplus L\mathbb{R}^j),$$

and let η_m be the canonical m -dimensional vector bundle. It is a \mathbb{Z}_2 -space with fixed subspace

$$G_m(\infty \oplus \infty L)^{\mathbb{Z}_2} = \bigsqcup_{r+s=m} G_r(\infty) \times G_s(\infty L)^{\mathbb{Z}_2},$$

and η_m restricts to the (r, s) -component as the sum of $(\eta_m)_+ = \eta_r \oplus 0$ and $(\eta_m)_- = 0 \oplus \eta_s$. Here

$$G_r(\infty) = \varinjlim_{i \geq 0} G_r(\mathbb{R}^i) = BO(r) \quad \text{and} \quad G_s(\infty L) = \varinjlim_{j \geq 0} G_s(L\mathbb{R}^j).$$

We have maps

$$\iota_m : G_m(\infty \oplus \infty L) \rightarrow G_{m+1}(\infty \oplus \infty L)$$

mapping an m -dimensional subspace $E \subseteq \mathbb{R}^i \oplus L\mathbb{R}^j$ to the $(m+1)$ -dimensional subspace $\mathbb{R} \oplus E$ of $\mathbb{R}^{i+1} \oplus L\mathbb{R}^j$. So $\iota_m^* \eta_{m+1} = \mathbb{R} \oplus \eta_m$.

The infinite Grassmannian $G_m(\infty \oplus \infty L)$ is a classifying space for \mathbb{Z}_2 -vector bundles of dimension m . To be precise, given an m -dimensional \mathbb{Z}_2 -bundle ζ over a compact \mathbb{Z}_2 -ENR W , there is a map $g : W \rightarrow G_m(\infty \oplus \infty L)$ and a bundle isomorphism $a : \zeta \rightarrow g^* \eta_m$ and any two such (g, a) are homotopic. (A pair (g, a) will be given by a bundle monomorphism $\zeta \hookrightarrow W \times V$ for some \mathbb{Z}_2 -module V .)

Definition B.34. We define $\mathcal{N}_m^{\mathbb{Z}_2}(X)$ be the bordism group of closed \mathbb{Z}_2 -manifolds M of dimension m with a \mathbb{Z}_2 -map $f : M \rightarrow X$.

Definition B.35. We construct a homomorphism

$$\mathcal{N}_m^{\mathbb{Z}_2}(X) \rightarrow \varinjlim_{n \geq m} \Omega_0^{\mathbb{Z}_2, \text{res}}(X \times G_n(\infty \oplus \infty L); \mathbb{R}^{n-m}, \eta_n),$$

the direct limit being formed with respect to the maps ι_n .

Consider a \mathbb{Z}_2 -map $f : M \rightarrow X$. We have a \mathbb{Z}_2 -map $g : M \rightarrow G_m(\infty \oplus \infty L)$ and bundle isomorphism $a : \tau M \rightarrow g^* \eta_m$ classifying the tangent bundle. The pair $(f \times g, a)$ defines an element of $\Omega_0^{\mathbb{Z}_2, \text{res}}(X \times G_m(\infty \oplus \infty L); 0, \eta_m)$ which depends only on $f : M \rightarrow X$. For a homotopy from (g, a) to (g', a') gives a cobordism $h : W = M \times [0, 1] \rightarrow G_m(\infty \oplus \infty L)$ and isomorphism $b : \tau W \cong \mathbb{R} \oplus h^* \eta_m$ restricting on ∂W to $(g, a) \sqcup (g', a')$.

This homomorphism is easily seen to be an isomorphism. For a class on the right is given by an m -manifold M , a map $f : M \rightarrow X$, a map $g_n : M \rightarrow G_n(\infty \oplus \infty L)$ and a restricted stable isomorphism $\tau M \oplus \mathbb{R}^{n-m} \cong g_n^* \eta_n$. The restricted stable isomorphism can be represented by a bundle isomorphism $\tau M \oplus \mathbb{R}^{n-m} \oplus \mathbb{R}^i \cong g_n^* \eta_n \oplus \mathbb{R}^i$ for some $i \geq 0$. Mapping from $G_n(\infty \oplus \infty L)$ to $G_{n+i}(\infty \oplus \infty L)$ we get a map $g_{n+i} : M \rightarrow G_{n+i}(\infty \oplus \infty L)$ and a bundle isomorphism $a_{n+i} : \tau M \oplus \mathbb{R}^{n-m+i} \cong g_{n+i}^* \eta_{n+i}$. This will be homotopic to the stabilization of any classifying pair (g, a) for M . Thus we obtain a homotopy-theoretic description of $\mathcal{N}_*^{\mathbb{Z}_2}$:

Theorem B.36. *There is a natural isomorphism*

$$\mathcal{N}_m^{\mathbb{Z}_2}(X) \cong \varinjlim_{n \geq m} \omega_m^{\mathbb{Z}_2}(X \times G_n(\infty \oplus \infty L); \mathbb{R}^n - \eta_n).$$

Proof. The group $\omega_m^{\mathbb{Z}_2}(X \times G_n(\infty \oplus \infty L); \mathbb{R}^n - \eta_n)$ is to be understood as the direct limit

$$\varinjlim_V \omega_m^{\mathbb{Z}_2}(X \times G_n(V); \mathbb{R}^n - \eta_n)$$

over \mathbb{Z}_2 -modules V and

$$\omega_m^{\mathbb{Z}_2}(X \times G_n(V); \mathbb{R}^n - \eta_n) = \omega_m^{\mathbb{Z}_2}(X \times G_n(V); \mathbb{R}^n \oplus \eta_n^\perp - V).$$

Or, better, we can take $V = \mathbb{R}^n \oplus V'$ and then

$$\omega_m^{\mathbb{Z}_2}(X \times G_n(\mathbb{R}^n \oplus V'); \mathbb{R}^n - \eta_n) = \omega_m^{\mathbb{Z}_2}(X \times G_n(\mathbb{R}^n \oplus V'); \eta_n^\perp - V').$$

Notice that, for $n \geq m$,

$$\Omega_m^{\mathbb{Z}_2, \text{res}}(X \times G_n(\mathbb{R}^n \oplus V'); \mathbb{R}^n, \eta_n) \rightarrow \omega_m^{\mathbb{Z}_2}(X \times G_n(\mathbb{R}^n \oplus V'); \mathbb{R}^n - \eta_n)$$

is an isomorphism.

Definition B.37. Write $\mathcal{N}_m^{\mathbb{Z}_2, \text{free}}(X)$ for the bordism group of maps $f : M \rightarrow X$ from a free \mathbb{Z}_2 -manifold M .

Proposition B.38. *The classical Conner-Floyd sequence*

$$\cdots \rightarrow \mathcal{N}_m^{\mathbb{Z}_2, \text{free}}(X) \rightarrow \mathcal{N}_m^{\mathbb{Z}_2}(X) \rightarrow \bigoplus_{s=0}^m \mathcal{N}_{m-s}(X^{\mathbb{Z}_2} \times G_s(\infty)) \rightarrow \cdots$$

corresponds to the stable homotopy localization sequence.

Proof. Recall that the non-equivariant bordism of an ANR Y is given by

$$\mathcal{N}_m(Y) = \varinjlim_n \omega_m(Y \times G_n(\infty); \mathbb{R}^n - \eta_n).$$

The identification on fixed points is made by the isomorphism

$$\begin{aligned} & \varinjlim_n \omega_m(X^{\mathbb{Z}_2} \times G_n(\infty \oplus \infty L)^{\mathbb{Z}_2}; \mathbb{R}^n - \eta_n^{\mathbb{Z}_2}) \\ &= \varinjlim_n \bigoplus_{0 \leq s \leq n} \omega_{m-s}((X^{\mathbb{Z}_2} \times G_{n-s}(\infty) \times G_s(\infty L)); \mathbb{R}^{n-s} - \eta_{n-s}) \\ &= \bigoplus_{s=0}^m \mathcal{N}_{m-s}(X^{\mathbb{Z}_2} \times G_s(\infty L)). \end{aligned}$$

The free term $\mathcal{N}_m^{\mathbb{Z}_2, \text{free}}(X)$ is isomorphic to $\mathcal{N}_m^{\mathbb{Z}_2}(E\mathbb{Z}_2 \times X)$, which in turn is isomorphic to $\mathcal{N}_m(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} X)$.

Remark B.39. We can think of $\mathcal{N}_m(Y)$ as the direct limit

$$\varinjlim_{V, V'} \omega_m(Y \times G_n(V \oplus V'); V - \eta_n)$$

over finite dimensional \mathbb{R} -vector spaces V and V' of dimension n and n' respectively. The Grassmann manifold $G_n(V \oplus V')$ is included in $G_{k+n}(U \oplus V \oplus U' \oplus V')$, where U and U' are vector spaces of dimension k and k' , by taking the direct sum of an n -dimensional subspace of $V \oplus V'$ with U to get an $(k+n)$ -dimensional subspace of $U \oplus V \oplus V' \subseteq U \oplus V \oplus U' \oplus V'$. In the bordism interpretation, this description classifies the stable tangent bundle. For the more conventional interpretation involving the stable normal bundle we can identify these groups with

$$\varinjlim_{V, V'} \omega_m(Y \times G_{n'}(V \oplus V'); \eta_{n'} - V').$$

The orthogonal complement identifies $G_n(V \oplus V')$ with $G_{n'}(V \oplus V')$ and then $\eta_n \oplus \eta_{n'} = V \oplus V'$. The corresponding inclusion of $G_{n'}(V \oplus V')$ as a subspace of $G_{k'+n'}(U \oplus V \oplus U' \oplus V')$ is given by the direct sum with U' .

Definition B.40. We have expressed $\mathcal{N}_m^{\mathbb{Z}_2}(X)$ as the direct limit

$$\varinjlim_n \varinjlim_{V'} \omega_m^{\mathbb{Z}_2}(X \times G_n(\mathbb{R}^n \oplus V'); \mathbb{R}^n - \eta_n),$$

where V' is a \mathbb{Z}_2 -module and $G_n(\mathbb{R}^n \oplus V')$ is included in $G_n(\mathbb{R}^n \oplus U' \oplus V')$ through the inclusion $\mathbb{R}^n \oplus V' \hookrightarrow \mathbb{R}^n \oplus U' \oplus V'$ and $G_n(\mathbb{R}^n \oplus V')$ is included in $G_{n+1}(\mathbb{R}^{n+1} \oplus V')$ by the direct sum with $\mathbb{R} \subseteq \mathbb{R} \oplus \mathbb{R}^n = \mathbb{R}^{n+1}$.

We now define

$$N_m^{\mathbb{Z}_2}(X) = \varinjlim_{V, V'} \omega_m^{\mathbb{Z}_2}(X \times G_n(V \oplus V'); V - \eta_n),$$

where V and V' are \mathbb{Z}_2 -modules of dimension n and n' respectively and $G_n(V \oplus V')$ is included in $G_{k+n}(U \oplus V \oplus U' \oplus V')$, for \mathbb{Z}_2 -modules U and U' of dimension k and k' , by the direct sum with U .

Remark B.41. In the complementary interpretation we have

$$N_m^{\mathbb{Z}_2}(X) = \varinjlim_{V, V'} \omega_m^{\mathbb{Z}_2}(X \times G_{n'}(V \oplus V'); \eta_{n'} - V'),$$

where the inclusion of $G_{n'}(V \oplus V')$ into $G_{k'+n'}(U \oplus V \oplus U' \oplus V')$ is given by the direct sum with U' .

For the non-equivariant theory we shall write $N_m(Y) = \mathcal{N}_m(Y)$ and then have the following N_* -theory Conner-Floyd or localization sequence.

Proposition B.42. *The Conner-Floyd/localization sequence for $N_*^{\mathbb{Z}_2}$ is*

$$\cdots \rightarrow N_m(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} X) \rightarrow N_m^{\mathbb{Z}_2}(X) \rightarrow \bigoplus_{s=-\infty}^m N_{m-s}(X^{\mathbb{Z}_2} \times BO(\infty)) \rightarrow \cdots$$

The standard definition of $N_m(Y)$ is as the direct limit

$$\varinjlim_{V, V'} [(\mathbb{R}^m \oplus V')^+; Y_+ \wedge G_{n'}(V \oplus V')^{\eta_{n'}}]$$

over vector spaces V and V' . But the stabilization

$$[(\mathbb{R}^m \oplus V')^+; Y_+ \wedge G_{n'}(V \oplus V')^{\eta_{n'}}] \rightarrow \omega_m(Y \times G_{n'}(V \oplus V'); \eta_{n'} - V')$$

is an isomorphism if $k + n' < 2n' - 1$. We have a similar interpretation in the \mathbb{Z}_2 -equivariant theory, for rather formal reasons.

Lemma B.43. *There is an isomorphism*

$$\alpha : \varinjlim_{V, V'} [(\mathbb{R}^m \oplus V')^+; X_+ \wedge G_{n'}(V \oplus V')^{\eta_{n'}}]^{\mathbb{Z}_2} \rightarrow N_m^{\mathbb{Z}_2}(X).$$

Proof. We have

$$N_m^{\mathbb{Z}_2}(X) = \varinjlim_{V, V', W} [(\mathbb{R}^m \oplus V' \oplus U')^+; X_+ \wedge G_{n'}(V \oplus V')^{\eta_{n'} \oplus U'}]^{\mathbb{Z}_2}$$

over \mathbb{Z}_2 -modules U' . The stabilization map α is defined by including the $U' = 0$ terms. We get a map

$$\beta : N_m^{\mathbb{Z}_2}(X) \rightarrow \varinjlim_{V, V', W} [(\mathbb{R}^m \oplus V' \oplus U')^+; X_+ \wedge G_{n'+k'}(V \oplus V' \oplus U')^{\eta_{n'+k'}}]^{\mathbb{Z}_2},$$

where $k' = \dim U'$, by embedding $G_{n'}(V \oplus V')$ into $G_{n'+k'}(V \oplus V' \oplus U')$ by taking the direct sum with U' so that $\eta_{n'+k'}$ restricts to $\eta_{n'} \oplus U'$.

The composition $\beta \circ \alpha$ is clearly the identity.

The composition $\alpha \circ \beta$ takes

$$[(\mathbb{R}^m \oplus V' \oplus U')^+; X_+ \wedge G_{n'}(V \oplus V')^{\eta_{n'} \oplus U'}]^{\mathbb{Z}_2}$$

to

$$[(\mathbb{R}^m \oplus V' \oplus U')^+; X_+ \wedge G_{n'+k'}(V \oplus V' \oplus U')^{\eta_{n'+k'}}]^{\mathbb{Z}_2}$$

to

$$\omega_m^{\mathbb{Z}_2}(X \times G_{n'+k'}(V \oplus (V' \oplus U'))); \eta_{n'+k'} - (V' \oplus U')),$$

which factors through

$$\omega_m^{\mathbb{Z}_2}(X \times G_{n'}(V \oplus V'); \eta_{n'} - V').$$

B.4 Finite groups

We conclude with some comments on the general case. The basic definitions and the statements of the theorems extend in an obvious way from the group \mathbb{Z}_2 to any finite group G . The proofs, while technically more complicated, follow the same pattern.

The main result expresses the restricted equivariant bordism groups as stable homotopy groups.

Theorem B.44. *There is a Pontryagin-Thom isomorphism*

$$\varphi : \Omega_o^{G, \text{res}}(X; \xi, \eta) \rightarrow \omega_0^G(F_X^{\text{res}}(\xi, \eta); \xi - \eta).$$

The theorem is proved by an induction over orbit types. Choose a filtration $\mathcal{F}_0 = \{1\} \subseteq \mathcal{F}_1 \subseteq \cdots \subseteq \mathcal{F}_r = \mathcal{F}$ of the family \mathcal{F} of all subgroups of G by subfamilies (closed under conjugation and taking subgroups) such that \mathcal{F}_i and \mathcal{F}_{i+1} differ by a single conjugacy class of subgroups. One then shows that

$$\varphi : \Omega_o^{G, \text{res}}(E\mathcal{F}_i \times X; \xi, \eta) \rightarrow \omega_0^G(E\mathcal{F}_i \times F_X^{\text{res}}(\xi, \eta); \xi - \eta)$$

is an isomorphism for $i = 0, \dots, k$ by comparing Conner-Floyd and localization exact sequences.

The G -equivariant unoriented bordism groups are expressed as limits of stable homotopy groups of Thom spaces.

Theorem B.45. *There is a natural isomorphism*

$$\mathcal{N}_m^G(X) \cong \varinjlim_n \varinjlim_{V'} \omega_m^G(X \times G_n(\mathbb{R}^n \oplus V'); \mathbb{R}^n - \eta_n)$$

over G -modules V' .

The related theory $N_*^G(X)$ may then be defined in the tangential formulation as the limit

$$N_m^G(X) = \varinjlim_{V'} \varinjlim_{V'} \omega_m^G(X \times G_n(V \oplus V'); V - \eta_n)$$

over G -modules V and V' . In normal form the groups $N_*^G(X)$ can be expressed as

$$\varinjlim_V \varinjlim_{V'} \omega_m^G(X \times G_{n'}(V \oplus V'); \eta_{n'} - V'),$$

and homotopy theoretically as

$$\varinjlim_V \varinjlim_{V'} [(\mathbb{R}^m \oplus V')^+; X_+ \wedge G_{n'}(V \oplus V')^{\eta_{n'}}]^G.$$

Appendix C

Reprint of

**“The geometric Hopf invariant and
double points”**

**Journal of Fixed Point Theory and
Applications 7, 325–350 (2010)**

(Including the correction of a misprint in
Proposition 3.1)

The geometric Hopf invariant and double points

Michael Crabb and Andrew Ranicki

Abstract. The geometric Hopf invariant of a stable map F is a stable $\mathbb{Z}/2$ -equivariant map $h(F)$ such that the stable $\mathbb{Z}/2$ -equivariant homotopy class of $h(F)$ is the primary obstruction to F being homotopic to an unstable map. In this paper we express the geometric Hopf invariant of the Umkehr map F of an immersion $f : M^m \looparrowright N^n$ in terms of the double point set of f . We interpret the Smale-Hirsch-Haefliger regular homotopy classification of immersions f in the metastable dimension range $3m < 2n - 1$ (when a generic f has no triple points) in terms of the geometric Hopf invariant.

Mathematics Subject Classification (2010). Primary 55Q25; Secondary 57R42.

Keywords. Geometric Hopf invariant, immersion, double point.

Introduction

The original Hopf invariant $H(F) \in \mathbb{Z}$ of a map $F : S^3 \rightarrow S^2$ was interpreted by Steenrod as the evaluation of the cup product in the mapping cone $C(F)$. The mod 2 Hopf invariant $H_2(F) \in \mathbb{Z}/2$ of a map $F : S^j \rightarrow S^k$ was then defined using the functional Steenrod squares of F . The geometric Hopf invariant of a stable map $F : \Sigma^\infty X \rightarrow \Sigma^\infty Y$ is the stable $\mathbb{Z}/2$ -equivariant map

$$h(F) = (F \wedge F)\Delta_X - \Delta_Y F : \Sigma^\infty X \rightarrow \Sigma^\infty(Y \wedge Y)$$

measuring the failure of F to preserve the diagonal maps of X and Y , with $\mathbb{Z}/2$ acting by the identity on X and by the transposition $T : (y_1, y_2) \mapsto (y_2, y_1)$ on $Y \wedge Y$. Thus $h(F)$ is a homotopy-theoretic generalization of the functional Steenrod squares. The stable homotopy class of $h(F)$ is the primary obstruction to F being homotopic to an unstable map.

Given an immersion $f : M^m \looparrowright N^n$ with normal bundle $\nu(f)$ we express the geometric Hopf invariant $h(F)$ of the Umkehr map $F : \Sigma^\infty N^+ \rightarrow \Sigma^\infty M^{\nu(f)}$ in terms of the double point set of f , where $M^{\nu(f)}$ denotes the

Thom space. There are many antecedents for this expression! The stable homotopy class of $h(F)$ depends only on the regular homotopy class of f . If f is regular homotopic to an embedding then $h(F)$ is stably null-homotopic. We interpret the Smale-Hirsch-Haefliger regular homotopy classification of immersions f in the metastable dimension range $3m < 2n - 1$ (when a generic f has no triple points) in terms of the geometric Hopf invariant.

In [6] we shall provide a considerably more detailed exposition of the geometric Hopf invariant $h(F)$ and its applications to manifolds. This will include the $\pi_1(N)$ -equivariant geometric Hopf invariant $\tilde{h}(F)$ needed for a homotopy-theoretic treatment of the double point invariant $\mu(f)$ of Wall [23] for a generic immersion $f : M^m \looparrowright N^{2m}$ which plays such an important rôle in non-simply-connected surgery theory, with $M = S^m$. When both M and N are connected and oriented and f induces the trivial map $\pi_1(M) \rightarrow \pi_1(N)$, $\mu(f)$ is an element of the group

$$\mathbb{Z}[\pi_1(N)] / \langle g - (-)^m g^{-1} \mid g \in \pi_1(N) \rangle$$

and $\mu(f) = 0$ if (and, for $m > 2$, only if) f is regular homotopic to an embedding, by the Whitney trick for removing double points. In [6] $\tilde{h}(F)$ will be shown to induce the quadratic construction ψ_F of [19] on the chain level.

The present paper is set out as follows. Section 1 describes briefly the construction of the geometric Hopf invariant and its fibrewise generalization. The double point theorem is stated and proved in Section 2. In Section 3, building on work of Dax [7], Hatcher and Quinn [10], Salomonsen [20] and Li, Liu and Zhang [18], we relate the geometric Hopf invariant in a stable range to Haefliger's obstruction to the existence of a regular homotopy from an immersion to an embedding. The papers of Boardman and Steer [2] and Koschorke and Sanderson [17] are also relevant. The variation of the geometric Hopf invariant of an immersion under a (not necessarily regular) homotopy is computed in Section 4 in terms of the Smale-Hirsch-Haefliger classification. In Section 5 we use Whitney's figure-of-eight immersion [24] to construct, in a metastable range, immersions close to a given embedding. Prerequisites for that section, on the differential-topological classification of vector bundle monomorphisms, are given in an Appendix.

We shall write the one-point compactification of a locally compact Hausdorff topological space X as X^+ . A subscript '+' will be used for the adjunction of a disjoint basepoint to a space. If X is compact $X^+ = X_+$. For a Euclidean vector bundle ξ over a general space X , we write $D(\xi)$ for the closed unit disc bundle, $S(\xi)$ for the sphere bundle and $B(\xi)$ for open unit disc bundle. The Thom space of ξ is written as X^ξ . To simplify notation, we shall sometimes write Y^ξ , rather than $Y^{p^*\xi}$, for the Thom space of the pullback $p^*\xi$ by a map $p : Y \rightarrow X$, if the map p is clear from the context. Similarly, we sometimes write the trivial vector bundle over X with fibre the vector space V as just V , instead of $X \times V$,

Methods of fibrewise homotopy theory will be used extensively. Fibrewise constructions, such as the one-point compactification, the Thom space,

or the smash product, over a base B will be indicated by attaching a subscript ‘ B ’ to the relevant symbol. We follow the notation for (fibrewise) stable homotopy adopted in [4]. Consider fibrewise pointed spaces $X \rightarrow B$ and $Y \rightarrow B$ over an ENR (= Euclidean neighbourhood retract) base B . If B is compact and A is a closed sub-ENR, we write

$$\omega_B^0\{X; Y\} \quad \text{and} \quad \omega_{(B,A)}^0\{X; Y\},$$

respectively, for the abelian group of stable fibrewise maps $X \rightarrow Y$ over B and the relative group defined in terms of homotopy classes of maps that are zero over the subspace A . (See, for example, [5, Part II, Section 3].) We also need to consider fibrewise maps with compact supports. When B is not necessarily compact we write

$${}_c\omega_B^0\{X; Y\}$$

for the group of fibrewise stable maps that are zero outside a compact subspace of B . The ω^0 -theories are extended, using the fibrewise suspension Σ_B over B , to ω^i -cohomology theories indexed by $i \in \mathbb{Z}$. When $Y \rightarrow B$ is a trivial bundle $B \times S^i \rightarrow B$, there are natural identifications of the fibrewise groups with the reduced stable cohomotopy of an appropriate pointed space:

$$\omega_B^0\{X; B \times S^i\} = \tilde{\omega}^i(X/B), \quad \text{and} \quad \omega_{(B,A)}^0\{X; B \times S^i\} = \tilde{\omega}^i(X/(X_A \cup B)),$$

where $X_A \rightarrow A$ denotes the restriction of $X \rightarrow B$.

We shall also need $\mathbb{Z}/2$ -equivariant stable homotopy theory, which we indicate by a prefix as ${}^{\mathbb{Z}/2}\omega^*$. So, for example, the equivariant stable cohomotopy of a point, ${}^{\mathbb{Z}/2}\omega^0(*)$, is the direct limit over all finite-dimensional \mathbb{R} -vector spaces V and W of the homotopy classes of pointed $\mathbb{Z}/2$ -maps $(V \oplus LW)^+ \rightarrow (V \oplus LW)^+$. Here, and throughout the paper, we write L for the non-trivial 1-dimensional representation \mathbb{R} of $\mathbb{Z}/2$ with the involution -1 , and, for a finite-dimensional real vector space W , abbreviate the tensor product $L \otimes W$ to LW .

We thank Mark Grant for pointing out the relevance of [10].

1. A review of the geometric Hopf invariant

Let X and Y be pointed topological spaces. It is convenient to assume that X is a compact ENR and that Y is an ANR (= absolute neighbourhood retract).

We shall be working both with the unreduced suspension of an unpointed space A

$$sA = [0, 1] \times A / \sim, \quad (0, a) \sim (0, a'), \quad (1, a) \sim (1, a') \quad (a, a' \in A),$$

and with the reduced suspension of a pointed space $(B, * \in B)$

$$\Sigma B = [0, 1] \times B / \sim, \quad (0, b) \sim (1, b') \sim (t, *) \quad (b, b' \in B, t \in [0, 1]).$$

Let V be a finite dimensional Euclidean space, and let

$$S(V) = \{u \in V \mid \|u\| = 1\}$$

be the unit sphere in V with respect to an inner product $\| \cdot \|$ on V . For $t \in [0, 1]$, $u \in S(V)$ we write

$$[t, u] = \frac{tu}{1-t} \in V^+$$

with $[0, u] = 0$, $[1, u] = + \in V^+$. The maps

$$\begin{aligned} sS(V) &\rightarrow V^+ ; (t, u) \mapsto [t, u] , \\ \Sigma S(V)_+ &\rightarrow V^+ / \{0, +\} ; (t, u) \mapsto [t, u] \end{aligned}$$

are homeomorphisms.

Let the generator $T \in \mathbb{Z}/2$ act on $X \wedge X$ by transposition

$$T : X \wedge X \rightarrow X \wedge X ; (x, y) \mapsto (y, x).$$

The diagonal map

$$\Delta_X : X \rightarrow X \wedge X ; x \mapsto (x, x)$$

is $\mathbb{Z}/2$ -equivariant. The diagonal map Δ_{V^+} extends to a $\mathbb{Z}/2$ -equivariant homeomorphism

$$\kappa_V : LV^+ \wedge V^+ \rightarrow V^+ \wedge V^+ ; (u, v) \mapsto (u + v, -u + v).$$

The $\mathbb{Z}/2$ -action $u \mapsto -u$ on LV has fixed point $\{0\}$; the $\mathbb{Z}/2$ -action on the unit sphere $S(LV)$ is free.

The geometric Hopf invariant of a map $F : V^+ \wedge X \rightarrow V^+ \wedge Y$ measures the difference $(F \wedge F)\Delta_X - \Delta_Y F$, given that $(F \wedge F)\Delta_{V^+ \wedge X} = \Delta_{V^+ \wedge Y} F$. The diagram of $\mathbb{Z}/2$ -equivariant maps

$$\begin{array}{ccc} LV^+ \wedge V^+ \wedge X & \xrightarrow{1 \wedge \Delta_X} & LV^+ \wedge V^+ \wedge X \wedge X \\ \downarrow 1 \wedge F & & \downarrow G \\ LV^+ \wedge V^+ \wedge Y & \xrightarrow{1 \wedge \Delta_Y} & LV^+ \wedge V^+ \wedge Y \wedge Y \end{array}$$

does not commute in general, with G defined by

$$\begin{aligned} G &= (\kappa_V^{-1} \wedge 1)(F \wedge F)(\kappa_V \wedge 1) : \\ &LV^+ \wedge V^+ \wedge X \wedge X \rightarrow LV^+ \wedge V^+ \wedge Y \wedge Y ; \\ &(u, v, x_1, x_2) \mapsto ((w_1 - w_2)/2, (w_1 + w_2)/2, y_1, y_2) \\ &(F(u + v, x_1) = (w_1, y_1), F(-u + v, x_2) = (w_2, y_2)). \end{aligned}$$

However, the $\mathbb{Z}/2$ -equivariant maps defined by

$$\begin{aligned} p &= G(1 \wedge \Delta_X) : LV^+ \wedge V^+ \wedge X \rightarrow LV^+ \wedge V^+ \wedge Y \wedge Y ; \\ &(u, v, x) \mapsto ((w_1 - w_2)/2, (w_1 + w_2)/2, y_1, y_2) \\ &(F(u + v, x) = (w_1, y_1), F(-u + v, x) = (w_2, y_2)), \\ q &= (1 \wedge \Delta_Y)(1 \wedge F) : LV^+ \wedge V^+ \wedge X \rightarrow LV^+ \wedge V^+ \wedge Y \wedge Y ; \\ &(u, v, x) \mapsto (u, w, y, y) \quad (F(v, x) = (w, y)) \end{aligned}$$

agree on $0^+ \wedge V^+ \wedge X = V^+ \wedge X \subseteq LV^+ \wedge V^+ \wedge X$, with

$$|p| = |q| = (\kappa_V^{-1} \wedge 1) \Delta_{V^+ \wedge Y} F = (\kappa_V^{-1} \wedge 1) (F \wedge F) (\kappa_V \wedge 1) \Delta_{V^+ \wedge X} : \\ V^+ \wedge X \rightarrow LV^+ \wedge V^+ \wedge Y \wedge Y.$$

Definition 1.1. ([5, pp. 306–308]) The *geometric Hopf invariant* of a pointed map $F : V^+ \wedge X \rightarrow V^+ \wedge Y$ is the $\mathbb{Z}/2$ -equivariant map given by the relative difference of the $\mathbb{Z}/2$ -equivariant maps p, q

$$h_V(F) = \delta(p, q) : \Sigma S(LV)_+ \wedge V^+ \wedge X \rightarrow LV^+ \wedge V^+ \wedge Y \wedge Y ; \\ (t, u, v, x) \mapsto \begin{cases} q([1 - 2t, u], v, x) & \text{if } 0 \leq t \leq 1/2 \\ p([2t - 1, u], v, x) & \text{if } 1/2 \leq t \leq 1 \end{cases} \\ (t \in [0, 1], u \in S(LV), v \in V, x \in X).$$

We are primarily interested in the stable $\mathbb{Z}/2$ -equivariant class of $h_V(F)$, which depends only on the homotopy class of F , and which we write simply as

$$h_V(F) \in {}^{\mathbb{Z}/2}\omega^0\{\Sigma S(LV)_+ \wedge X; LV^+ \wedge (Y \wedge Y)\}.$$

Using duality, we can rewrite this stable homotopy group in different ways and thus obtain two other versions of the geometric Hopf invariant as follows. Smashing $h_V(F)$ with the identity on the sphere $S(LV)_+$ and composing with the duality map

$$LV^+ \rightarrow \Sigma S(LV)_+ \rightarrow S(LV)_+ \wedge \Sigma S(LV)_+$$

we get a map

$$V^+ \wedge LV^+ \wedge X \rightarrow V^+ \wedge LV^+ \wedge S(LV)_+ \wedge (Y \wedge Y)$$

and a second version of the geometric Hopf invariant as an element

$$h'_V(F) \in {}^{\mathbb{Z}/2}\omega^0\{X; S(LV)_+ \wedge (Y \wedge Y)\}.$$

Remark 1.2. The $\mathbb{Z}/2$ -equivariant cofibration sequence

$$S(LV)_+ \rightarrow S^0 = \{0\}^+ \rightarrow LV^+$$

induces an exact sequence of stable $\mathbb{Z}/2$ -equivariant homotopy groups

$${}^{\mathbb{Z}/2}\omega^0\{X; S(LV)_+ \wedge (Y \wedge Y)\} \rightarrow {}^{\mathbb{Z}/2}\omega^0\{X; Y \wedge Y\} \rightarrow {}^{\mathbb{Z}/2}\omega^0\{X; LV^+ \wedge Y \wedge Y\}.$$

The stable class $h'_V(F)$ has image $(F \wedge F) \Delta_X - \Delta_Y F$ in ${}^{\mathbb{Z}/2}\omega^0\{X; Y \wedge Y\}$.

We may also use the Adams isomorphism [1, Theorem 5.3]

$${}^{\mathbb{Z}/2}\omega^0\{X; S(LV)_+ \wedge (Y \wedge Y)\} \cong \omega^0\{X; S(LV)_+ \wedge_{\mathbb{Z}/2} (Y \wedge Y)\}$$

to regard $h'_V(F)$ as a non-equivariant class

$$h''_V(F) \in \omega^0\{X; S(LV)_+ \wedge_{\mathbb{Z}/2} (Y \wedge Y)\}.$$

Remark 1.3. Instead of applying the Adams isomorphism, we can use fibre-wise techniques. The unstable map $h_V(F)$ lifts to an equivariant fibrewise pointed map over $S(LV)$:

$$S(LV) \times (\Sigma V^+ \wedge X) \rightarrow S(LV) \times (V^+ \wedge LV^+ \wedge (Y \wedge Y)),$$

and we may divide by the free $\mathbb{Z}/2$ -action to get a fibrewise pointed map over the real projective space $P(V) = S(LV)/\mathbb{Z}/2$:

$$P(V) \times (\Sigma V^+ \wedge X) \rightarrow (V \oplus (H \otimes V))_{P(V)}^+ \wedge_{P(V)} (S(LV) \times_{\mathbb{Z}/2} (Y \wedge Y)),$$

where $H = S(LV) \times_{\mathbb{Z}/2} L$ is the Hopf line bundle over $P(V)$ and $\{-\}_{P(V)}^+$ denotes fibrewise one-point compactification over $P(V)$. This fibrewise map represents a stable class in the group

$$\omega_{P(V)}^{-1} \{P(V) \times X; (H \otimes V)_{P(V)}^+ \wedge_{P(V)} (S(LV) \times_{\mathbb{Z}/2} (Y \wedge Y))\},$$

which may be identified by fibrewise Poincaré-Atiyah duality (since $\mathbb{R} \oplus \tau P(V) = H \otimes V$) with

$$\omega^0 \{X; S(LV)_+ \wedge_{\mathbb{Z}/2} (Y \wedge Y)\}.$$

(For the duality theorem, see, for example, [5, Part II, Section 12].)

Remark 1.4. By [3, pp. 60–61] the limit over all finite-dimensional V of the exact sequences in Remark 1.2 is a split exact sequence

$$\omega^0 \{X; (E\mathbb{Z}/2)_+ \wedge_{\mathbb{Z}/2} (Y \wedge Y)\} \xrightarrow{\delta} \mathbb{Z}/2 \omega^0 \{X; Y \wedge Y\} \xrightarrow{\rho} \omega^0 \{X; Y\}$$

with the fixed point map ρ split by

$$\omega^0 \{X; Y\} \rightarrow \mathbb{Z}/2 \omega^0 \{X; Y \wedge Y\}; F \mapsto \Delta_Y F.$$

The stable $\mathbb{Z}/2$ -equivariant homotopy class

$$h(F) = (F \wedge F) \Delta_X - \Delta_Y F \in \mathbb{Z}/2 \omega^0 \{X; Y \wedge Y\}$$

is the image under δ of

$$h'(F) = \varinjlim_V h'_V(F) \in \omega^0 \{X; (E\mathbb{Z}/2)_+ \wedge_{\mathbb{Z}/2} (Y \wedge Y)\}.$$

We shall need some fundamental properties of the geometric Hopf invariant. First, it is an obstruction to desuspension. The Hopf invariant of a suspension is zero, but there is a more precise result for the geometric Hopf invariant.

Given a pointed space $(A, * \in A)$ and $a \neq * \in A$ let $a_1, a_2 \in A \vee A$ denote the two distinct copies of a , and write

$$A \vee_a A = (A \vee A) / (a_1 \sim a_2).$$

Lemma 1.5. (i) *The $\mathbb{Z}/2$ -equivariant maps*

$$i_V : S(LV) \rightarrow \Sigma S(LV)_+; v \mapsto (1/2, v),$$

$$j_V : \Sigma S(LV)_+ \rightarrow LV^+ \vee_0 LV^+; (t, u) \mapsto \begin{cases} [1 - 2t, u]_1 & \text{if } 0 \leq t \leq 1/2 \\ [2t - 1, u]_2 & \text{if } 1/2 \leq t \leq 1 \end{cases},$$

$$k_V : LV^+ \vee_0 LV^+ \rightarrow LV^+; v_1 \mapsto v, v_2 \mapsto v$$

fit into a $\mathbb{Z}/2$ -equivariant cofibration

$$S(LV) \xrightarrow{i_V} \Sigma S(LV)_+ \xrightarrow{j_V} LV^+ \vee_0 LV^+ \xrightarrow{k_V} LV^+$$

with the composite $k_V \circ j_V : \Sigma S(LV)_+ \rightarrow LV^+$ a $\mathbb{Z}/2$ -equivariantly null-homotopic map.

(ii) The geometric Hopf invariant of $F : V^+ \wedge X \rightarrow V^+ \wedge Y$ is the composite

$$\begin{aligned} h_V(F) &= (q \vee_0 p)(j_V \wedge 1_{V^+ \wedge X}) : \\ \Sigma S(LV)_+ \wedge V^+ \wedge X &\rightarrow (LV^+ \vee_0 LV^+) \wedge V^+ \wedge X \rightarrow LV^+ \wedge V^+ \wedge Y \wedge Y \end{aligned}$$

(iii) Suppose that F is the suspension $1_{V^+} \wedge F_0$ of a map $F_0 : X \rightarrow Y$, so that

$$p = q = 1 \wedge (\Delta_Y \circ F_0) : LV^+ \wedge V^+ \wedge X \rightarrow LV^+ \wedge V^+ \wedge Y \wedge Y.$$

Then

$$h_V(F) = 1 \wedge (k_V \circ j_V) \wedge (\Delta_Y \circ F_0) \simeq *$$

is $\mathbb{Z}/2$ -equivariantly null-homotopic.

The second property, giving a formula for the Hopf invariant of a composition, is suggested by the stable form described in Remark 1.2.

Proposition 1.6. (Composition formula). *Let X, Y and Z be pointed spaces, and suppose that $F : V^+ \wedge X \rightarrow V^+ \wedge Y$ and $G : V^+ \wedge Y \rightarrow V^+ \wedge Z$ are pointed maps. Then*

$$h_V(G \circ F) = h_V(G)[F] + [G \wedge G]h_V(F) \in \mathbb{Z}/2\omega^0\{\Sigma S(LV)_+ \wedge X; LV^+ \wedge (Z \wedge Z)\},$$

where

$$[F] \in \mathbb{Z}/2\omega^0\{X; Y\} \quad \text{and} \quad [G \wedge G] \in \mathbb{Z}/2\omega^0\{Y \wedge Y; Z \wedge Z\}$$

are the stable classes determined by F (with the trivial action of $\mathbb{Z}/2$) and $G \wedge G$.

Lastly, the Hopf invariant satisfies the following sum formula.

Proposition 1.7. (Sum formula). *Let F_+, F_- be maps $V^+ \wedge X \rightarrow V^+ \wedge Y$. Suppose that $v \in S(V)$. Choose a tubular neighbourhood of $\{v, -v\}$ in V and let $\nabla : V^+ \rightarrow V^+ \wedge V^+$ be the associated Pontryagin-Thom map. Then the Hopf invariant of the sum $F = (F_+ \wedge F_-) \circ (\nabla \wedge 1_X)$ is given by*

$$h_V(F) = h_V(F_+) + h_V(F_-) + \iota[(F_+ \wedge F_-) \circ \Delta_X]$$

where the induction homomorphism ι is the composition of the isomorphism

$$\omega^0\{X; Y \wedge Y\} \cong \mathbb{Z}/2\omega^0\{\Sigma S(Lv)_+ \wedge X; (Lv)^+ \wedge (Y \wedge Y)\}$$

and the map

$$\mathbb{Z}/2\omega^0\{\Sigma S(Lv)_+ \wedge X; (Lv)^+ \wedge (Y \wedge Y)\} \rightarrow \mathbb{Z}/2\omega^0\{\Sigma S(LV)_+ \wedge X; LV^+ \wedge (Y \wedge Y)\}$$

induced by the inclusion of the 1-dimensional subspace $\mathbb{R}v \hookrightarrow V$.

The explicit construction of the geometric Hopf invariant is readily extended to the fibrewise theory. Suppose now that $X \rightarrow B$ and $Y \rightarrow B$ are fibrewise pointed spaces over an ENR B . (We shall assume that both are locally fibre homotopy trivial and that the fibres have the homotopy type of CW complexes, finite complexes in the case of X .) Consider a fibrewise pointed map $F : (B \times V^+) \wedge_B X \rightarrow (B \times V^+) \wedge_B Y$. If B is compact, we have a fibrewise geometric Hopf invariant

$$h_V(F) \in \mathbb{Z}/2 \omega_B^0 \{ (B \times \Sigma S(LV)_+) \wedge_B X; (B \times LV^+) \wedge_B (Y \wedge_B Y) \},$$

and corresponding variants $h'_V(F)$ and $h''_V(F)$. See, for example, [5, Part II, Section 14]. This fibrewise Hopf invariant is an obstruction to fibrewise desuspension. Indeed, suppose that the restriction of F to a closed sub-ENR $A \subseteq B$ is the fibrewise suspension of a map $X_A \rightarrow Y_A$ over A . Then Lemma 1.5 (iii) shows how to define a relative fibrewise Hopf invariant in

$$\mathbb{Z}/2 \omega_{(B,A)}^0 \{ (B \times \Sigma S(LV)_+) \wedge_B X; (B \times LV^+) \wedge_B (Y \wedge_B Y) \},$$

which lifts $h_V(F)$. (One uses the fact that the inclusion of A in B is a cofibration.) When B is not (necessarily) compact and F is a fibrewise suspension outside some compact subspace of B , the same method gives a fibrewise Hopf invariant with compact supports:

$$h_V(F) \in \mathbb{Z}/2 \omega_B^0 \{ (B \times \Sigma S(LV)_+) \wedge_B X; (B \times LV^+) \wedge_B (Y \wedge_B Y) \}.$$

2. The double point theorem

Let $f : M \looparrowright N$ be a (smooth) immersion of a closed manifold M in a connected manifold N (without boundary) of dimension n , with normal bundle $\nu(f)$, usually abbreviated to ν . We do not require M to be connected, nor that all the components should have the same dimension; the maximum dimension of a component is denoted by m . Let $e : M \rightarrow V$ be a smooth map to a finite-dimensional Euclidean space V such that $e(x) \neq e(y)$ whenever $f(x) = f(y)$ for $x \neq y$. This gives a (smooth) embedding $(e, f) : M \hookrightarrow V \times N$ with normal bundle $V \oplus \nu$. (To be precise, there is a short exact sequence $0 \rightarrow V \rightarrow \nu(e, f) \rightarrow \nu \rightarrow 0$ which is split by a choice of metrics.)

We have an associated Pontryagin-Thom map (defined up to homotopy)

$$F : V^+ \wedge N^+ \rightarrow V^+ \wedge M^\nu.$$

Its geometric Hopf invariant is a stable $\mathbb{Z}/2$ -homotopy class

$$h_V(F) \in \mathbb{Z}/2 \omega^0 \{ \Sigma S(LV)_+ \wedge N^+; LV^+ \wedge (M^\nu \wedge M^\nu) \},$$

where $\mathbb{Z}/2$ interchanges the factors of $M^\nu \wedge M^\nu$.

Suppose that the immersion f is self-transverse and that there are no k -tuple points for $k > 2$. (This is the case for a generic immersion if $3m < 2n$.) The *double point set*

$$\mathfrak{D}(f) = \{ (x, y) \in M \times M - \Delta(M) \mid f(x) = f(y) \}$$

is then a smooth $\mathbb{Z}/2$ -submanifold of $M \times M$ (of constant dimension $2m - n$ if M is connected), on which $\mathbb{Z}/2$ acts freely, and its normal bundle may be identified with the pullback $j^*\tau N$ of the tangent bundle of N by the map $j : \mathfrak{D}(f) \rightarrow N$ mapping (x, y) to $f(x) = f(y) \in N$. We also have a $\mathbb{Z}/2$ -map $d : \mathfrak{D}(f) \rightarrow LV - \{0\}$ given by $d(x, y) = e(x) - e(y)$, and thus an embedding

$$(d, j) : \mathfrak{D}(f) \hookrightarrow (LV - \{0\}) \times N,$$

with normal bundle $LV \oplus k^*(\nu \times \nu)$, where $k : \mathfrak{D}(f) \rightarrow M \times M$ is the inclusion. The Pontryagin-Thom construction gives a $\mathbb{Z}/2$ -map

$$(LV - \{0\})^+ \wedge N^+ \rightarrow LV^+ \wedge \mathfrak{D}(f)^{k^*(\nu \times \nu)}.$$

Composing with the map induced by k , we get a $\mathbb{Z}/2$ -homotopy class

$$\phi : \Sigma S(LV)_+ \wedge N^+ \rightarrow LV^+ \wedge (M^\nu \wedge M^\nu).$$

Theorem 2.1. (The double point theorem). *The geometric Hopf invariant $h_V(F)$ of the Pontryagin-Thom map F is equal to the $\mathbb{Z}/2$ -equivariant stable homotopy class of the map ϕ determined, as described above, by the double point manifold $\mathfrak{D}(f)$.*

We may also consider the second version of the Hopf invariant

$$h'_V(F) \in \mathbb{Z}/2 \omega^0 \{N^+; S(LV)_+ \wedge (M^\nu \wedge M^\nu)\}.$$

This, too, may be described directly in terms of the double points. The embedding $\mathfrak{D}(f) \hookrightarrow LV \times N$ with normal bundle $LV \oplus k^*(\nu \times \nu)$ provides a Pontryagin-Thom map

$$LV^+ \wedge N^+ \rightarrow LV^+ \wedge \mathfrak{D}(f)^{k^*(\nu \times \nu)}$$

which we compose with the map $\mathfrak{D}(f) \rightarrow S(LV) \times (M \times M)$ given by e and the inclusion k to get

$$\phi' : LV^+ \wedge N^+ \rightarrow LV^+ \wedge S(LV)_+ \wedge (M^\nu \wedge M^\nu).$$

Corollary 2.2. *The second version $h'_V(F)$ of the geometric Hopf invariant is represented by the $\mathbb{Z}/2$ -map ϕ' defined in the text.*

We also have the non-equivariant stable Hopf invariant

$$h''_V(F) \in \omega^0 \{N^+; (S(LV) \times_{\mathbb{Z}/2} (M \times M))^{\nu \times \nu}\}.$$

The free $\mathbb{Z}/2$ -manifold $\mathfrak{D}(f)$ is a double cover of the set $\overline{\mathfrak{D}}(f) \subseteq N$ of double points of the immersion f . We have an induced map

$$\overline{\mathfrak{D}}(f) = \mathfrak{D}(f)/\mathbb{Z}/2 \rightarrow S(LV) \times_{\mathbb{Z}/2} (M \times M)$$

and an embedding of $\overline{\mathfrak{D}}(f)$ in N with normal bundle the pullback of $\nu \times \nu$. The Pontryagin-Thom construction gives a map

$$\phi'' : N^+ \rightarrow \overline{\mathfrak{D}}(f)^{\nu \times \nu} \rightarrow (S(LV) \times_{\mathbb{Z}/2} (M \times M))^{\nu \times \nu}.$$

Corollary 2.3. *The stable geometric Hopf invariant $h''_V(F)$ is equal to the stabilization $[\phi'']$ of the class determined by the double point manifold $\overline{\mathfrak{D}}(f) \subseteq N$.*

These results will be obtained as consequences of a more precise fibre-wise theorem, which we describe next.

Let $\mathcal{C} \rightarrow N$ be the space of pairs (x, α) where $x \in M$ and $\alpha : [0, 1] \rightarrow N$ is a continuous path such that $\alpha(0) = f(x)$, fibred over N by projection to the other endpoint $\alpha(1)$. We have a homotopy equivalence $\pi : \mathcal{C} \rightarrow M$ given by $\pi(x, \alpha) = x$.

The homotopy Pontryagin-Thom map defined by the embedding $(e, f) : M \hookrightarrow V \times N$ as described in [4, Section 6] (and in [5]) is a pointed map

$$\tilde{F} : N \times V^+ \rightarrow (N \times V^+) \wedge_N \mathcal{C}_N^{\pi^* \nu}$$

with compact supports over N . (To be exact, the space \mathcal{C} in [4] is the space of pairs (x, β) , where $x \in M$ and $\beta : [0, 1] \rightarrow V \times N$ is a path starting at $\beta(0) = (e(x), f(x))$. We omit here the redundant component in the contractible space V .) We may then form the fibrewise geometric Hopf invariant

$$h_V(\tilde{F}) \in {}^{\mathbb{Z}/2}_c \omega_N^0 \{N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N (\mathcal{C}_N^{\pi^* \nu} \wedge_N \mathcal{C}_N^{\pi^* \nu})\},$$

again as a stable $\mathbb{Z}/2$ -equivariant map with compact supports over N .

Now the fibre product $\mathcal{C} \times_N \mathcal{C}$ of pairs $((x, \alpha), (y, \beta))$ such that $\alpha(1) = \beta(1)$ may be identified, by splicing α to the reversed path β , with the space $\text{h-}\widehat{\mathfrak{D}}(f)$ of triples (x, y, γ) with $x, y \in M$ and $\gamma : [-1, 1] \rightarrow N$ a continuous path from $\gamma(-1) = f(x)$ to $\gamma(1) = f(y)$ projecting to $\gamma(0) \in N$. (Thus $\gamma(t)$ is $\alpha(1+t)$ if $-1 \leq t \leq 0$, $\beta(1-t)$ if $0 \leq t \leq 1$.) It has an action of $\mathbb{Z}/2$ in which the involution interchanges x and y and reverses the path γ . The double point set $\mathfrak{D}(f)$ is included in $\text{h-}\widehat{\mathfrak{D}}(f)$ as the space of constant paths.

Let \mathcal{D} be the space of pairs $((x, y), \alpha)$, where $(x, y) \in \mathfrak{D}(f)$ and $\alpha : [0, 1] \rightarrow N$ is a path such that $\alpha(0) = f(x) = f(y)$. This corresponds to the subspace of points $(x, y, \gamma) \in \text{h-}\widehat{\mathfrak{D}}(f)$ with $\gamma(-t) = \gamma(t)$. The fibrewise Pontryagin-Thom construction on $\mathfrak{D}(f) \hookrightarrow (LV - \{0\}) \times N$ gives a fibrewise map

$$N \times \Sigma S(LV)_+ \rightarrow (N \times LV^+) \wedge_N \mathcal{D}_N^{\nu \times \nu},$$

which we compose with the inclusion

$$\mathcal{D}_N^{\nu \times \nu} \hookrightarrow \text{h-}\widehat{\mathfrak{D}}(f)_N^{\nu \times \nu},$$

to get an equivariant fibrewise map

$$\tilde{\phi} : N \times \Sigma S(LV)_+ \rightarrow (N \times LV^+) \wedge_N \text{h-}\widehat{\mathfrak{D}}(f)_N^{\nu \times \nu}.$$

Theorem 2.4. (Homotopy double point theorem). *The fibrewise geometric Hopf invariant*

$$h_V(\tilde{F}) \in {}^{\mathbb{Z}/2}_c \omega_N^0 \{N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N \text{h-}\widehat{\mathfrak{D}}(f)_N^{\nu \times \nu}\}$$

of the homotopy Pontryagin-Thom map \tilde{F} is equal to the fibrewise stable class of the map $\tilde{\phi}$ determined by the double points of f .

The dual version is a class

$$h'_V(\tilde{F}) \in {}^{\mathbb{Z}/2}_c \omega_N^0 \{N \times S^0; (N \times S(LV)_+) \wedge_N h\text{-}\widehat{\mathfrak{D}}(f)_N^{\nu \times \nu}\}.$$

There is also a non-equivariant form. The stable Hopf invariant $h''_V(\tilde{F})$ lies in

$${}_c \omega_N^0 \{N \times S^0; (S(LV) \times_{\mathbb{Z}/2} h\text{-}\widehat{\mathfrak{D}}(f))_N^{\nu \times \nu}\},$$

and this group can be identified with

$$\tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} h\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N})$$

by fibrewise Poincaré-Atiyah duality. (For a general treatment of fibrewise duality see, for example, [5, Part II, Section 12]. The duality theorem required here is stated in [4] as Proposition 4.1.)

The Pontryagin-Thom construction applied to the double point manifold $\widehat{\mathfrak{D}}(f) = \mathfrak{D}(f)/(\mathbb{Z}/2)$ equipped with the map $\widehat{\mathfrak{D}}(f) \rightarrow S(LV) \times_{\mathbb{Z}/2} h\text{-}\widehat{\mathfrak{D}}(f)$ given by the inclusion $\mathfrak{D}(f) \rightarrow h\text{-}\widehat{\mathfrak{D}}(f)$ and the map $\mathfrak{D}(f) \rightarrow S(LV)$ given by e (via d) produces a stable homotopy class

$$\tilde{\phi}'' : S^0 \rightarrow (S(LV) \times_{\mathbb{Z}/2} h\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N}.$$

Corollary 2.5. *We have*

$$h''_V(\tilde{F}) = \tilde{\phi}'' \in \tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} h\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N}).$$

Before turning to the proof of Theorem 2.4, we explain how the fibrewise Hopf invariant $h_V(\tilde{F})$ determines the simpler invariant $h_V(F)$.

Lemma 2.6. *The Hopf invariant $h_V(F)$ is the image of $h_V(\tilde{F})$ under the composition:*

$$\begin{aligned} & {}^{\mathbb{Z}/2}_c \omega^0 \{N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N h\text{-}\widehat{\mathfrak{D}}(f)_N^{\nu \times \nu}\} \\ & \rightarrow {}^{\mathbb{Z}/2}_c \omega^0 \{N^+ \wedge \Sigma S(LV)_+; LV^+ \wedge h\text{-}\widehat{\mathfrak{D}}(f)^{\nu \times \nu}\} \\ & \rightarrow {}^{\mathbb{Z}/2}_c \omega^0 \{N^+ \wedge \Sigma S(LV)_+; LV^+ \wedge (M^\nu \wedge M^\nu)\} \end{aligned}$$

of the homomorphism defined by collapsing the basepoint sections over N to a point and that induced by the projection $\pi \times \pi : h\text{-}\widehat{\mathfrak{D}}(f) = \mathcal{C} \times_N \mathcal{C} \rightarrow M \times M$.

Proof. This is easily seen from the explicit construction of the geometric Hopf invariant. \square

In a similar manner, $h'_V(F)$ and $h''_V(F)$ are the images of the refined invariants $h'_V(\tilde{F})$ and $h''_V(\tilde{F})$ under homomorphisms defined by collapsing basepoint sections over N and projecting from $h\text{-}\widehat{\mathfrak{D}}(f)$ to $M \times M$.

Remark 2.7. This construction also provides the π -equivariant Hopf invariant considered, in greater detail, in [6].

(i) Suppose that π is a discrete group and that $q : \tilde{N} \rightarrow N$ is a principal π -bundle (for example, a universal covering space with π the fundamental group of N). We let $\widetilde{M} = f^* \tilde{N}$ be the induced bundle over M : thus

$$\widetilde{M} = \{(x, z) \in M \times \tilde{N} \mid f(x) = q(z)\}.$$

We may define a map $\mathrm{h}\text{-}\widehat{\mathfrak{D}}(f) \rightarrow (\widetilde{M} \times \widetilde{M})/\pi$ as follows. Given $(x, y, \gamma) \in \mathrm{h}\text{-}\widehat{\mathfrak{D}}(f)$ (so $x, y \in M$, $\gamma : [-1, 1] \rightarrow N$, $\gamma(-1) = f(x)$, $\gamma(1) = f(y)$), lift γ to a path $\tilde{\gamma}$ in \widetilde{N} , determined up to multiplication by an element of π . The π -orbit of $((x, \tilde{\gamma}(-1)), (y, \tilde{\gamma}(1))) \in \widetilde{M} \times \widetilde{M}$ is independent of the choice of the lift. The Hopf invariant $h_V''(\tilde{F})$ thus gives us an element of

$$\omega^0\{N^+; (S(LV) \times_{\mathbb{Z}/2} ((\widetilde{M} \times \widetilde{M})/\pi))^{\nu \times \nu}\},$$

where $\nu \times \nu$ is lifted from $M \times M$ to $(\widetilde{M} \times \widetilde{M})/\pi$ by the obvious projection.

(ii) Let $(f, b) : N \rightarrow X$ be a normal map from an n -dimensional manifold N to an n -dimensional geometric Poincaré complex X . Let \tilde{X} be the universal cover of X , let $\tilde{N} = f^*\tilde{X}$ be the pullback cover of N , and let $\pi = \pi_1(X)$. The Umkehr $\mathbb{Z}[\pi]$ -module chain map

$$f^! : C(\tilde{X}) \simeq C(\tilde{X})^{n-*} \xrightarrow{\tilde{f}^*} C(\tilde{N})^{n-*} \simeq C(\tilde{N})$$

is induced by a π -equivariant geometric Umkehr map $F : \Sigma^\infty \tilde{X}_+ \rightarrow \Sigma^\infty \tilde{N}_+$ π -equivariant S -dual to $T(\tilde{b}) : T(\tilde{\nu}_N) \rightarrow T(\tilde{\nu}_X)$. The Wall surgery obstruction $\sigma_*(f, b) \in L_n(\mathbb{Z}[\pi])$ was identified in [19] with the cobordism class of an n -dimensional quadratic Poincaré complex $(C(f^!), \psi)$ over $\mathbb{Z}[\pi]$, with ψ the evaluation of the ‘quadratic construction’ ψ_F on the fundamental class $[X] \in H_n(X)$. In [6] we shall express ψ_F in terms of the π -equivariant geometric Hopf invariant of F .

(iii) See [12] for an application of the π -equivariant Hopf invariant to diagonals in geometric Poincaré complexes.

In view of the relations between the various Hopf invariants, it will suffice to prove Theorem 2.4 in order to establish Theorem 2.1 and their sundry corollaries.

Proof of Theorem 2.4. Writing $\bar{\nu}$ for the normal bundle of $\overline{\mathfrak{D}}(f)$ in N , choose a tubular neighbourhood $D(\bar{\nu}) \hookrightarrow N$ of $\overline{\mathfrak{D}}(f)$. For $(x, y) \in \mathfrak{D}(f)$, the fibre of $\bar{\nu}$ at $f(x) = f(y)$ is $\nu_x \oplus \nu_y$. We identify $f^{-1}(\overline{\mathfrak{D}}(f))$ with $\mathfrak{D}(f)$ by projecting to the first factor. Then the inverse image of the tubular neighbourhood $D(\bar{\nu})$ is a tubular neighbourhood $D(\nu')$ of $\mathfrak{D}(f)$ in M , where $\nu'_x = \nu_y$. The normal bundle ν restricted to $D(\nu')$ is then identified with the restriction of ν to $\mathfrak{D}(f)$.

We may assume that e vanishes outside the tubular neighbourhood $D(\nu')$. The homotopy Pontryagin-Thom map is then a V -fold suspension outside $D(\bar{\nu})$. By Proposition 1.5, the fibrewise Hopf invariant is canonically null-homotopic outside $D(\bar{\nu})$. The Hopf invariant $h_V(\tilde{F})$ is thus represented by a fibrewise map which is zero outside the tubular neighbourhood. In this way we localize $h_V(\tilde{F})$ to an element of

$$\mathbb{Z}/2 \omega_{(D(\bar{\nu}), S(\bar{\nu}))}^0 \{N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N \mathrm{h}\text{-}\widehat{\mathfrak{D}}(f)_N^{\nu \times \nu}\}$$

constructed from the immersion data in a neighbourhood of the double points.

The local data consists simply of the double cover $\mathfrak{D}(f) \rightarrow \overline{\mathfrak{D}}(f)$ and the vector bundle ν over $\mathfrak{D}(f)$. The bundle ν' is the pullback of ν under the covering involution and $\bar{\nu}$ over $\overline{\mathfrak{D}}(f)$ is the push-forward of ν . The ‘local M ’ is the total space of ν' over $\mathfrak{D}(f)$, and the ‘local N ’ is the total space of $\bar{\nu}$ over $\overline{\mathfrak{D}}(f)$. The immersion f is given by the projection $\mathfrak{D}(f) \rightarrow \overline{\mathfrak{D}}(f)$. We also need the map e and, without loss of generality, we may assume that it is determined by a $\mathbb{Z}/2$ -map $\mathfrak{D}(f) \rightarrow S(LV)$ (extended radially on $D(\nu')$ to taper to zero).

In the fibre over $\{x, y\} \in \overline{\mathfrak{D}}(f)$, the manifold M is the disjoint union

$$(\{0\} \times \nu_y) \sqcup (\nu_x \times \{0\}) \looparrowright \nu_x \oplus \nu_y$$

immersed in the fibre of N by projection, with the double point at $(0, 0)$. Write $e(x) = v$, $e(y) = -v$. The Hopf invariant is calculated by the sum formula of Proposition 1.7. In the notation used there, we take $X = \nu_x^+ \times \nu_y^+$, $Y = \nu_x^+ \vee \nu_y^+$, and the maps F_+ and F_- are suspensions of the compositions of the projection to ν_x^+ or ν_y^+ , respectively, and the inclusion of the wedge summand. The Hopf invariants of the suspensions F_+ and F_- vanish and the Hopf invariant of the sum is determined by the product $X \rightarrow Y \times Y$, so by the projection $\nu_x^+ \times \nu_y^+ \rightarrow \nu_x^+ \wedge \nu_y^+$.

The same computation performed fibrewise over $\overline{\mathfrak{D}}(f)$ gives the localized fibrewise Hopf invariant as the image of the element in

$$\omega_{(D(\bar{\nu}), S(\bar{\nu}))}^0 \{D(\bar{\nu}) \times S^0; \bar{\nu}_{D(\bar{\nu})}^+\}$$

given by the inclusion of $D(\bar{\nu})$ in $\bar{\nu}$. Unravelling the definitions, one sees that this produces $[\phi]$. \square

In the remainder of the paper we shall be concerned with the behaviour of the geometric Hopf invariant $h_V(\tilde{F})$ as one deforms the map $f : M \rightarrow N$. Suppose, to begin, that we have smooth homotopies $f_t : M \rightarrow N$ and $e_t : M \rightarrow V$ such that each f_t is an immersion and such that $(e_t, f_t) : M \rightarrow V \times N$ is an embedding for each $t \in [0, 1]$. We write $f = f_0$, $f' = f_1$, and $e = e_0$, $e' = e_1$. Then we have, up to homotopy, an isomorphism $\nu' = \nu(f_1) \rightarrow \nu(f_0) = \nu$ between the normal bundles of the immersions and a fibre homotopy equivalence $\mathbf{h}\widehat{\mathfrak{D}}(f') \rightarrow \mathbf{h}\widehat{\mathfrak{D}}(f)$ over N . In the standard terminology, the homotopy f_t is a *regular homotopy* from f to f' and the homotopy (e_t, f_t) is an *isotopy* from (e, f) to (e', f') . Let F' be the map obtained from the homotopy Pontryagin-Thom construction on (e', f') .

Proposition 2.8. (Regular homotopy/isotopy invariance). *The fibrewise Hopf invariants $h_V(\tilde{F}')$ and $h_V(\tilde{F})$ correspond under the induced isomorphism:*

$$\begin{aligned} & \mathbb{Z}/2 \omega_N^0 \{N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N \mathbf{h}\widehat{\mathfrak{D}}(f')^{\nu' \times \nu'}\} \\ & \rightarrow \mathbb{Z}/2 \omega_N^0 \{N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N \mathbf{h}\widehat{\mathfrak{D}}(f)^{\nu \times \nu}\}. \end{aligned}$$

Proof. This follows from the homotopy invariance of the geometric Hopf invariant, which in turn follows easily from the explicit construction described in the Introduction. \square

3. Immersions and embeddings

The vanishing of the stable Hopf invariant $h_V(\tilde{F})$ is a necessary condition for the existence of a regular homotopy f_t and isotopy (e_t, f_t) such that $f' = f_1$ is an embedding. In the opposite direction, we record first:

Proposition 3.1. *Suppose that $3m < 2(n - 1)$ and that $h_V(\tilde{F}) = 0$. Then \tilde{F} is homotopic (through maps with compact support over N) to the V -fold suspension of a map determined by a section $N \rightarrow \mathcal{C}_N^{\pi^* \nu}$.*

Proof. This is a consequence of the fibrewise EHP-sequence (see [5, Part II, Proposition 14.39]), applicable because $0 \leq 3(n - m - 1) - (n + 0)$. \square

To proceed further, we shall assume that the dimension of V is large: $\dim V > 2m$. This guarantees that M can be embedded in V , and we may take the map $e : M \rightarrow V$ to be an embedding. In a metastable range, the fibrewise Hopf invariant of \tilde{F} is the precise obstruction to the existence of a regular homotopy from the immersion $f : M \looparrowright N$ to an embedding.

Theorem 3.2. (Haefliger [8], Hatcher-Quinn [10]). *Suppose that $3m < 2(n - 1)$ and $\dim V > 2m$. Then*

$$h_V''(\tilde{F}) \in \tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N}) = \tilde{\omega}_0((E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N})$$

vanishes if and only if f is homotopic through immersions to an embedding of M into N .

We define $\mathbf{h}\text{-}\mathfrak{D}(f)$ to be the subspace of $\mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)$ consisting of the triples (x, y, γ) such that $x \neq y$ and call $\mathbf{h}\text{-}\mathfrak{D}(f)$ the space of *homotopy double points* of the immersion f . The Hopf bundle $E\mathbb{Z}/2 \times_{\mathbb{Z}/2} L$ over $B\mathbb{Z}/2$ will be denoted by H .

Lemma 3.3. *Let $\mathfrak{L}(f)$ be the complement of $\mathbf{h}\text{-}\mathfrak{D}(f)$ in $\mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)$. Then one has a homotopy cofibration sequence:*

$$\begin{aligned} (E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\mathfrak{D}(f))^{\nu \times \nu - \tau N} &\rightarrow (E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N} \\ &\rightarrow (E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathfrak{L}(f))^{H \otimes \tau M + \nu \times \nu - \tau N}, \end{aligned}$$

in which the first map is given by the inclusion of $\mathbf{h}\text{-}\mathfrak{D}(f)$ in $\mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)$ and the second by the homotopy Pontryagin-Thom construction on the diagonal submanifold M in $M \times M$.

As a space over M , $\mathfrak{L}(f)$ is the pullback by $f : M \rightarrow N$ of the free loop space of N , $\text{map}(S^1, N) \rightarrow N$, fibred over N by evaluation at $1 \in S^1 (\subseteq \mathbb{C})$.

Proof. The vector bundle $L \otimes \tau M$, corresponding to $H \otimes \tau M$ over $B\mathbb{Z}/2$, is the normal bundle of the diagonal inclusion of M in $M \times M$. Choose a $\mathbb{Z}/2$ -equivariant tubular neighbourhood $D(L \otimes \tau M) \hookrightarrow M \times M$ of the diagonal in $M \times M$. The inclusion of the subspace of $\mathbf{h}\text{-}\mathfrak{D}(f)$ consisting of the triples (x, y, γ) such that $(x, y) \notin B(L \otimes \tau M)$ into $\mathbf{h}\text{-}\mathfrak{D}(f)$ is a homotopy equivalence.

The argument used in the proof of Lemma 6.1 in [4] then shows that we have a homotopy cofibre sequence. \square

Corollary 3.4. *The inclusion induces an isomorphism*

$$\tilde{\omega}_0((E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\mathfrak{D}(f))^{\nu \times \nu - \tau N}) \rightarrow \tilde{\omega}_0((E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N})$$

provided that $m < n - 1$.

Proof. This follows at once from the long exact sequence of the cofibration. \square

The involution on $\mathbf{h}\text{-}\mathfrak{D}(f)$ is free, and hence, writing $\mathbf{h}\text{-}\overline{\mathfrak{D}}(f)$ for the quotient $\mathbf{h}\text{-}\mathfrak{D}(f)/(\mathbb{Z}/2)$, we have an isomorphism

$$\tilde{\omega}_0((E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\mathfrak{D}(f))^{\nu \times \nu - \tau N}) \rightarrow \tilde{\omega}_0(\mathbf{h}\text{-}\overline{\mathfrak{D}}(f)^{\nu \times \nu - \tau N}).$$

Until now, we have thought of $\mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)$, which arose as the fibre product $\mathcal{C} \times_N \mathcal{C}$, as a space over N . It also fibres over $M \times M$ and the fibrewise space $\mathbf{h}\text{-}\widehat{\mathfrak{D}}(f) \rightarrow M \times M$ is $\mathbb{Z}/2$ -equivariantly locally fibre homotopy trivial. (Compare [4, Definition 2.3].) In the same way, $\mathbf{h}\text{-}\mathfrak{D}(f)$ is fibred over the complement $M \times M - \Delta(M)$ and $\mathbf{h}\text{-}\overline{\mathfrak{D}}(f)$ is fibred over $(M \times M - \Delta(M))/\mathbb{Z}/2$.

Proof of Theorem 3.2. We shall use results and terminology from [4, Section 7], which derive from work of Koschorke [16] and Klein and Williams [13].

Suppose, first, that M is connected. Put $\tilde{B} = M \times M - B(L \otimes \tau M)$; it is a manifold with a free $\mathbb{Z}/2$ -action. Let B be the manifold $\tilde{B}/\mathbb{Z}/2$ with boundary the projective bundle, $\partial B = P(\tau M)$. Let E be the bundle $(\tilde{B} \times (N \times N))/\mathbb{Z}/2$ over B and $Z \subseteq E$ the diagonal sub-bundle $B \times N = (\tilde{B} \times N)/\mathbb{Z}/2$. The fibrewise normal bundle of the inclusion $Z \hookrightarrow E$ is $H \otimes \tau N$, where H is the line bundle over B associated to the double cover. The $\mathbb{Z}/2$ -equivariant square $f \times f : \tilde{B} \rightarrow N \times N$ defines a section s of $E \rightarrow B$, which, if the tubular neighbourhood is chosen to be sufficiently small, has the property that $s(x) \notin Z_x$ for $x \in \partial B$. Together, the fibre bundle $E \rightarrow B$, the sub-bundle $Z \rightarrow B$ and the section s constitute what is called in [4] an intersection problem. In the language used there, s is nowhere null on the boundary ∂B and the homotopy null-set $\mathbf{h}\text{-Null}(s)$, fibred over B , is easily identified with the restriction of $\mathbf{h}\text{-}\mathfrak{D}(f) \rightarrow (M \times M - \Delta(M))/\mathbb{Z}/2$. The inclusion $\mathbf{h}\text{-Null}(s) \hookrightarrow \mathbf{h}\text{-}\overline{\mathfrak{D}}(f)$ is, as we have already noted in the proof of Lemma 3.3, a homotopy equivalence.

If f is an embedding, then the section s is nowhere null. Now the homotopy Euler index ([4, Definition 7.3])

$$\mathbf{h}\text{-}\gamma(s; \partial B) \in \tilde{\omega}_0(\mathbf{h}\text{-Null}(s)^{H \otimes \tau N - \tau B})$$

is an obstruction to deforming s , through a homotopy constant on ∂B , to a section that is nowhere null. By Proposition 7.4 of [4] it is the precise obstruction if $\dim B < 2(\dim N - 1)$. We conclude that if $\mathbf{h}\text{-}\gamma(s; \partial B) = 0$ and $m < n - 1$, then

$$f \times f : M \times M - B(L \otimes \tau M) \rightarrow N \times N$$

is $\mathbb{Z}/2$ -equivariantly homotopic, by a homotopy that is constant on the boundary $SL \otimes \tau M$, to a map into $N \times N - \Delta(N)$. According to Haefliger [8,

Théorème 2], if further $3m < 2(n - 1)$, then this is a sufficient condition for $f : M \looparrowright N$ to be homotopic through immersions to an embedding of M into N .

Thus far, the argument is taken from [13, Theorem A.4 and Corollary A.5]. We now relate the index $\mathrm{h}\text{-}\gamma(s; \partial B)$ to the geometric Hopf invariant.

Lemma 3.5. *The homotopy Euler index*

$$\mathrm{h}\text{-}\gamma(s; \partial B) \in \tilde{\omega}_0(\mathrm{h}\text{-}\mathrm{Null}(s)^{\nu \times \nu - \tau N})$$

corresponds to the fibrewise Hopf invariant

$$h_V''(\tilde{F}) \in \tilde{\omega}_0((E\mathbb{Z}/2 \otimes_{\mathbb{Z}/2} \mathrm{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N}).$$

Proof. The correspondence is made via the homotopy equivalence $\mathrm{h}\text{-}\mathrm{Null}(s) \hookrightarrow \mathrm{h}\text{-}\widehat{\mathfrak{D}}(f)$ and the isomorphism from Corollary 3.4.

We can assume that f satisfies the conditions for the homotopy double point theorem. Then $h_V''(\tilde{F})$ is represented by the double point manifold $\widehat{\mathfrak{D}}(f)$. By Proposition 7.8 of [4], the homotopy Euler index is represented by the null-set $\mathrm{Null}(s)$, which is exactly $\widehat{\mathfrak{D}}(f)$. \square

Hence the vanishing of $h_V''(\tilde{F})$ implies, in the metastable range, that f is regularly homotopic to an embedding.

This has dealt with the case in which M is connected. Now suppose that M is a disjoint union $M^{(1)} \sqcup M^{(2)}$ and write $f^{(i)}$ for the restriction of f to $M^{(i)}$. Then $\mathrm{h}\text{-}\widehat{\mathfrak{D}}(f)$ splits equivariantly over

$$M \times M = (M^{(1)} \times M^{(1)}) \sqcup (M^{(2)} \times M^{(2)}) \sqcup (M^{(1)} \times M^{(2)} \sqcup M^{(2)} \times M^{(1)})$$

as a disjoint union

$$\mathrm{h}\text{-}\widehat{\mathfrak{D}}(f^{(1)}) \sqcup \mathrm{h}\text{-}\widehat{\mathfrak{D}}(f^{(2)}) \sqcup (\mathbb{Z}/2 \times \mathrm{h}\text{-}\mathfrak{I}(f^{(1)}, f^{(2)})),$$

where $\mathrm{h}\text{-}\mathfrak{I}(f^{(1)}, f^{(2)})$ is the space of triples (x, y, γ) with $(x, y) \in M^{(1)} \times M^{(2)}$ and $\gamma(-1) = f^{(1)}(x)$, $\gamma(1) = f^{(2)}(y)$. The fibrewise Hopf invariant decomposes, according to Proposition 1.7, as a sum of three terms. The first two are the fibrewise Hopf invariants of $f^{(1)}$ and $f^{(2)}$; the third, (12)-component, is a more elementary product obstruction.

Arguing by induction, we may suppose that both $f^{(1)}$ and $f^{(2)}$ are embeddings, intersecting transversely, and that $M^{(2)}$ is connected. We consider a new intersection problem with $B = M^{(2)}$, $E = B \times N$ and $Z = B \times f(M^{(1)})$. Let $s : B \rightarrow E$ be the section $s(x) = (x, f^{(2)}(x))$. This time $\mathrm{h}\text{-}\mathrm{Null}(s)$ is $\mathrm{h}\text{-}\mathfrak{I}(f^{(1)}, f^{(2)})$, and we may identify the homotopy Euler index $\mathrm{h}\text{-}\gamma(s)$ in the same way with the (12)-component of the fibrewise Hopf invariant, both being represented by the manifold $f(M^{(1)}) \cap f(M^{(2)})$. By Proposition 7.4 of [4] again, the vanishing of the homotopy Euler index implies that $f^{(2)}$ is homotopic to a map into $N - f(M^{(1)})$, because $\dim M^{(2)} < 2(n - \dim M^{(1)} - 1)$. Finally, we may apply [10, Theorem 1.1] to deduce that $f^{(2)}$ is isotopic to an embedding of $M^{(2)}$ into the complement of $f(M^{(1)})$. This inductive step is enough to conclude the proof of Theorem 3.2. \square

Remark 3.6. Suppose that M (as well as N) is connected. Choose basepoints $*$ $\in M$ and $* = f(*) \in N$. Then we can include the loop-space ΩN in $\mathbf{h}\widehat{\mathfrak{D}}(f)$ by mapping a loop $\gamma : [-1, 1] \rightarrow N$, with $\gamma(-1) = * = \gamma(1)$, to $(*, *, \gamma) \in \mathbf{h}\widehat{\mathfrak{D}}(f)$, and the set of path components of $\mathbf{h}\widehat{\mathfrak{D}}(f)$ is in this way identified with the set of double cosets

$$f_*\pi_1(M) \backslash \pi_1(N) / f_*\pi_1(M) = \pi_0(\mathbf{h}\widehat{\mathfrak{D}}(f))$$

with the $\mathbb{Z}/2$ -action given by the group-theoretic inverse. Thus we may identify the set of path components of $S(LV) \times_{\mathbb{Z}/2} \mathbf{h}\widehat{\mathfrak{D}}(f)$, if $\dim V > 1$, with the orbit space of the involution.

Example 3.7. Suppose that $n = 2m$. Then $\tilde{\omega}_0((E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathbf{h}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N})$ is a direct sum of groups indexed by $\pi_0(\mathbf{h}\widehat{\mathfrak{D}}(f))/\mathbb{Z}/2$, each component being isomorphic to \mathbb{Z} or $\mathbb{Z}/2\mathbb{Z}$. When M is connected we may label the summands as in Remark 3.6 by equivalence classes of group elements $g \in \pi_1(N)$. Let $w_M : \pi_1(M) \rightarrow \{\pm 1\}$ and $w_N : \pi_1(N) \rightarrow \{\pm 1\}$ be the orientation maps (corresponding to w_1M and w_1N). The g -summand is isomorphic to \mathbb{Z} if and only if for all $a, b \in \pi_1(M)$: (o) if $f_*(a)g = gf_*(b)$, then $w_M(ab) = w_N(f_*(a)) (= w_N(f_*(b)))$, and (i) if $f_*(a)g = g^{-1}f_*(b)$, then $(-1)^m w_M(ab) = w_N(f_*(a)g)$. In particular, if M is orientable, N is oriented and $f_*\pi_1(M)$ is trivial, then the obstruction group is $\mathbb{Z}[\pi_1(N)]/\langle g - (-1)^m g^{-1} \mid g \in \pi_1(N) \rangle$ and the Hopf invariant $h_V''(\tilde{F})$ is Wall's invariant $\mu(f)$ in the form mentioned in the Introduction.

4. Homotopic immersions

We next investigate immersions homotopic, as maps, to the given immersion f . Consider smooth homotopies $f_t : M \rightarrow N$ and $e_t : M \rightarrow V$ such that f_0 and f_1 are immersions and each map $(e_t, f_t) : M \rightarrow V \times N$, for $0 \leq t \leq 1$, is an embedding with normal bundle μ_t . We again write $f = f_0$, $f' = f_1$, and $\nu = \nu(f)$, $\nu' = \nu(f')$. The homotopies determine, up to a homotopy, a bundle isomorphism $a : V \oplus \nu' = \mu_1 \rightarrow \mu_0 = V \oplus \nu$ and a $\mathbb{Z}/2$ -equivariant fibre homotopy equivalence $\mathbf{h}\widehat{\mathfrak{D}}(f') \rightarrow \mathbf{h}\widehat{\mathfrak{D}}(f)$ over $M \times M$.

There is an associated class

$$\theta(e_t, f_t) \in \tilde{\omega}_0((P(V) \times M)^{H \otimes \nu - \tau M})$$

which vanishes if each f_t is an immersion. It is constructed as follows. Let $v_0 : V \rightarrow V \oplus \nu$ be the inclusion and let $v_1 : V \rightarrow V \oplus \nu' = \mu_1 \rightarrow \mu_0 = V \oplus \nu$ be the composition of the inclusion with the isomorphism a . Now we have a stable cohomotopy difference class

$$\delta(v_0, v_1) \in \tilde{\omega}^{-1}((M \times P(V))^{-H \otimes (V \oplus \nu)}),$$

which is the metastable obstruction to deforming v_0 to v_1 through vector bundle monomorphisms; see Section 6 for details. We define $\theta(e_t, f_t)$ to be the dual class in stable homotopy. (Recall that $\tau P(V) \oplus \mathbb{R} = H \otimes V$.)

Using the inclusion $M \hookrightarrow \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f) : x \mapsto (x, x, \gamma)$, where γ is the constant loop at $f(x)$, we get a map

$$i : \tilde{\omega}_0((P(V) \times M)^{H \otimes \nu - \tau M}) \rightarrow \tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N}).$$

Remark 4.1. The map $M \rightarrow \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)$ picks out (under the assumption that M is connected) a component $\mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)_0$ of $\mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)$, which is preserved by $\mathbb{Z}/2$. Hence i maps into the summand

$$\tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f)_0)^{\nu \times \nu - \tau N}).$$

The link between the difference class θ and the Hopf invariant is forged by a generalization of the classical description of the Hopf invariant on the image of the J -homomorphism.

Proposition 4.2. *Let $a : V \oplus \nu' \rightarrow V \oplus \nu$ be a vector bundle isomorphism over M . The fibrewise one-point compactification of a is a map of sphere bundles*

$$A : (M \times V^+) \wedge_M (\nu')_M^+ \rightarrow (M \times V)^+ \wedge \nu_M^+$$

over M . Then the fibrewise Hopf invariant

$h_V(A) \in \mathbb{Z}/2 \omega_M^0 \{(M \times \Sigma S(LV)_+) \wedge_M (\nu')_M^+; (M \times LV^+) \wedge_M (\nu_M^+ \wedge_M \nu_M^+)\}$ of A coincides up to sign, under the identifications described below, with the difference class

$$\delta(v_0, v_1) \in \tilde{\omega}^{-1}((M \times P(V))^{-H \otimes (V \oplus \nu)})$$

of the monomorphisms $v_0, v_1 : V \rightarrow V \oplus \nu$ given, respectively, by the inclusion of the first factor and the composition of the inclusion $V \rightarrow V \oplus \nu'$ with a .

Proof. The fibrewise smash product $\nu_M^+ \wedge_M \nu_M^+ = (\nu \oplus \nu)_M^+$ with the action of $\mathbb{Z}/2$ which interchanges the factors is equivariantly homeomorphic (by the construction κ_V in Section 1) to $(\nu \oplus L\nu)_M^+ = \nu_M^+ \wedge (L\nu)_M^+$. The isomorphism $a : V \oplus \nu' \rightarrow V \oplus \nu$ gives a stable fibre homotopy equivalence $(\nu')_M^+ \rightarrow \nu_M^+$. Taken together, these equivalences allow us to think of $h_V(A)$ as an element of the group

$$\mathbb{Z}/2 \omega_M^0 \{M \times \Sigma S(LV)_+; (LV \oplus L\nu)_M^+\},$$

which is then identified with

$$\mathbb{Z}/2 \tilde{\omega}^{-1}((M \times S(LV))^{-(LV \oplus L\nu)}) = \tilde{\omega}^{-1}((M \times P(V))^{-H \otimes (V \oplus \nu)}).$$

Both $h_V(A)$ and $\delta(v_0, v_1)$ are defined by difference constructions. The proof that they coincide follows from a direct comparison of the definitions. \square

Theorem 4.3. (Homotopy/isotopy variation). *The Hopf invariant*

$$h_V''(\tilde{F}') \in \tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f'))^{\nu' \times \nu' - \tau N})$$

corresponds to

$$h_V''(\tilde{F}) + i\theta(e_t, f_t) \in \tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N}).$$

Proof. Recollect that \tilde{F} is a fibrewise map with compact supports over N :

$$N \times V^+ \rightarrow (N \times V^+) \wedge_N \mathcal{C}_N^{\pi^* \nu}.$$

Up to homotopy, the map \tilde{F}' associated with (e', f') is the composition of \tilde{F} with the map

$$(N \times V^+) \wedge_N \mathcal{C}_N^{\pi^* \nu} = \mathcal{C}_N^{\pi^*(V \oplus \nu)} \rightarrow \mathcal{C}_N^{\pi^*(V \oplus \nu')} = (N \times V^+) \wedge_N \mathcal{C}_N^{\pi^* \nu'}$$

obtained by lifting the bundle isomorphism $a^{-1} : V \oplus \nu \rightarrow V \oplus \nu'$ over M via $\pi : \mathcal{C} \rightarrow M$. In the notation of Proposition 4.2 we have $\tilde{F} \simeq \pi^* A \circ \tilde{F}'$.

The fibrewise version of the composition formula (Proposition 1.6) expresses the difference $\alpha = h_V(\tilde{F}) - h_V(\tilde{F}')$ in terms of A . Using Proposition 4.2 and the explicit form of the geometric Hopf invariant one sees that α lies in the image of the diagonal map

$$\begin{aligned} \Delta_* : \quad & \mathbb{Z}/2 \omega_N^0 \{ N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N \mathcal{C}_N^{\pi^*(\nu \oplus LV)} \} \\ & \rightarrow \mathbb{Z}/2 \omega_N^0 \{ N \times \Sigma S(LV)_+; (N \times LV^+) \wedge_N (\mathcal{C}_N^{\pi^* \nu} \wedge_N \mathcal{C}_N^{\pi^* \nu'}) \}. \end{aligned}$$

This may be rewritten in dual form as:

$$\tilde{\omega}_0((P(V) \times \mathcal{C})^{\pi^*(\nu \oplus H \otimes \nu - \tau N)}) \rightarrow \tilde{\omega}_0((S(LV) \times_{\mathbb{Z}/2} \mathbf{h}\text{-}\widehat{\mathfrak{D}}(f))^{\nu \times \nu - \tau N}).$$

But $\pi : \mathcal{C} \rightarrow M$ is a homotopy equivalence. Hence Δ_* is just another manifestation of the map i in the statement of the theorem.

Now the tubular neighbourhood of M in $V \times N$ gives a proper map $B(\nu) \rightarrow N$. To calculate α , which is concentrated on the diagonal, we can thus lift from N to $B(\nu)$. Here we have a fibrewise problem over M , and the identification of α is achieved by Proposition 4.2. \square

Remark 4.4. In the stable range $\dim V > 2m$, where the maps e_t are redundant, we can use the methods of the previous section to give an alternative proof of Theorem 4.3.

Theorem 4.5. (Hirsch [11]). *Suppose that $3m < 2n - 1$ and $\dim V > 2m$. Then two immersions f and f' are homotopic through immersions if and only if the associated difference class θ in $\tilde{\omega}_0((P(V) \times M)^{H \otimes \nu - \tau M})$ is zero.*

Proof. The derivative of the immersion f gives a vector bundle monomorphism $df : \tau M \rightarrow f^* \tau N$ over M . According to Hirsch [11], for $m < n$ immersions $f' : M \rightarrow N$ together with a homotopy f_t from $f = f_0$ to $f' = f_1$ are classified by homotopy classes of vector bundle monomorphisms $\tau M \rightarrow f^* \tau N$. In the metastable range $m + 1 < 2(n - m)$, that is, $3m < 2n - 1$, immersions with a homotopy to f are thus classified by

$$\tilde{\omega}^{-1}(P(\tau M)^{-H \otimes f^* \tau N}) = \tilde{\omega}_0(P(\tau M)^{H \otimes (f^* \tau N - \tau M) - \tau M}) = \tilde{\omega}_0(P(\tau M)^{H \otimes \nu - \tau M}).$$

Assuming that $\dim V > 2m$ we may fix an embedding $e : M \hookrightarrow V$. The derivative of e includes τM in the trivial bundle $M \times V$ and gives an isomorphism

$$\tilde{\omega}_0(P(\tau M)^{H \otimes \nu - \tau M}) \rightarrow \tilde{\omega}_0((M \times P(V))^{H \otimes \nu - \tau M}). \quad \square$$

5. Immersions close to an embedding

Consider the special case of a closed manifold M of (constant) dimension m and a real vector bundle ν of dimension $n - m$ over M . Working in the metastable range $3m < 2n - 1$, we take N to be the total space of ν and $f : M \rightarrow N$ to be the embedding given by the zero section of the vector bundle. As $e : M \rightarrow V$ we may take the constant map 0.

Let $v_0 : V \hookrightarrow V \oplus \nu$ be the inclusion of the first factor. Suppose that $v_1 : V \hookrightarrow V \oplus \nu$ is another inclusion, which we may assume to be isometric. Thus v_0 and v_1 give sections of the bundle $O(V, V \oplus \nu)$ whose fibre at $x \in M$ is the Stiefel manifold of orthogonal linear maps $v : V \rightarrow V \oplus \nu_x$. Let $X_1(V, \nu)$ be the sub-bundle with fibre consisting of those linear maps v such that $v + i_x$ has kernel of dimension 1, where $i_x : V \rightarrow V \oplus \nu_x$ is the inclusion of the first factor. In the range of dimensions that we are considering, for a generic smooth section v_1 of $O(V, V \oplus \nu) \rightarrow M$ the kernel of $(v_1)_x + i_x$ is nowhere of dimension > 1 , v_1 is transverse to the sub-bundle $X_1(V, \nu)$ and $v_1^{-1}(X_1(V, \nu))$ is a submanifold Z of M of dimension $2m - n$, equipped with a map $Z \rightarrow P(V)$ classifying the 1-dimensional kernel. The normal bundle of Z in M is identified with $H \otimes \nu$ and Z represents the element $\delta \in \tilde{\omega}_0((M \times P(V))^{H \otimes \nu - \tau M})$ dual to $\delta(v_0, v_1)$. More details are provided in an appendix (Section 6).

We want to construct an immersion f' close to the zero section f with double point set Z . More precisely, we shall construct homotopies f_t and e_t , with $f_0 = f$, $f_1 = f'$ and each (e_t, f_t) an embedding, such that $\overline{\mathfrak{D}}(f') = Z$ and $\theta(e_t, f_t) = \delta$.

Choose an open tubular neighbourhood $\Omega = H \otimes \nu \hookrightarrow M$ of Z . Since $\dim \nu = n - m > 2m - n = \dim Z$, we can split off a trivial line from the restriction of ν to Z as: $\nu|_Z = \mathbb{R} \oplus \zeta$.

Whitney gave in [24], for a Euclidean space U , an explicit ‘punctured figure-of-eight’ immersion:

$$w : \mathbb{R} \oplus U \rightarrow (\mathbb{R} \oplus U) \times (\mathbb{R} \oplus U)$$

with double points at $(\pm 1, 0)$. In slightly modified form it may be written as

$$w(s, y) = ((1 - \lambda(s, y))s, y, -\lambda(s, y), \lambda(s, y)sy),$$

where $\lambda(s, y) = \psi(s^2 + \|y\|^2)$ and $\psi : [0, \infty) \rightarrow \mathbb{R}$ is a smooth, non-negative, monotonic decreasing function, with $\psi(1) = 1$, $\psi'(1) = -1/2$ and $\psi(r) = 0$ for $r \geq 2$. (In [24], $\psi(r) = 2/(1 + r)$.) The derivative at the double point $(\pm 1, 0)$ is

$$\frac{\partial w}{\partial s} = (1, 0, \pm 1, 0), \quad \frac{\partial w}{\partial y} = (0, 1, 0, \pm 1).$$

Writing $w(s, y)$ in the form $(s, y, 0, 0) + \lambda(s, y)(-s, 0, -1, sy)$, we see that $w(s, y) = (s, y, 0, 0)$ for $\|(s, y)\|$ large. The immersion w has $\mathbb{Z}/2$ -symmetry as an equivariant map

$$w : L \oplus LU \rightarrow (L \oplus LU) \times (\mathbb{R} \oplus U).$$

The two double points are distinguished by the $\mathbb{Z}/2$ -map

$$c : L \oplus LU \rightarrow L$$

given by $c(s, y) = \lambda(s, y)s$.

Now Whitney's construction, applied on the fibres of ζ , gives an immersion

$$\begin{aligned} f' : \Omega = H \otimes \nu = H \oplus (H \otimes \zeta) \\ \rightarrow (H \oplus (H \otimes \zeta)) \times (\mathbb{R} \oplus \zeta) = (H \otimes \nu) \times \nu = \nu|_{\Omega} \subseteq N \end{aligned}$$

of the open subset Ω of M into the total space of the restriction of ν to Ω . We extend f' to the whole of M to coincide with the zero section f outside a compact subspace of Ω . Its double point set $\mathfrak{D}(f')$ is the double-cover $S(H|Z)$ of Z in Ω . The map c composed with the inclusion of H into the trivial bundle $\Omega \times V$ and the projection to V gives a map

$$e' : \Omega = H \oplus (H \otimes \zeta) \rightarrow V,$$

which is zero outside a compact subset of Ω and can be extended by 0 to a map $e' : M \rightarrow V$ which distinguishes the double point pairs. The required homotopies e_t and f_t , $t \in [0, 1]$, joining e to e' and f to f' are defined by replacing λ in the definition of e' and f' by $t\lambda$. One checks that (e_t, f_t) is an embedding for all t , but that f_t fails to be an immersion when $t = 1/\psi(0)$.

Theorem 5.1. *Suppose that $3m < 2n - 1$. Then Whitney's construction described above produces, for any given element $\delta \in \tilde{\omega}_0((M \times P(V))^{H \otimes \nu - \tau M})$, a homotopy (e_t, f_t) with $\theta(e_t, f_t) = \delta$.*

Proof. By construction, the double point manifold $\mathfrak{D}(f') = S(H|Z)$ represents δ . The assertion thus follows from the Double Point Theorem 2.4 for (e', f') in conjunction with the Homotopy Variation Theorem 4.3. \square

Remark 5.2. The same construction may be used to modify a general immersion $f : M \looparrowright N$, and map $e : M \rightarrow V$, in the complement of $\mathfrak{D}(f)$. We can insert Z in the complement, because $2(2m - n) < m$.

Example 5.3. Whitney's construction gives the classical immersions of the sphere $S^m \looparrowright S^{2m}$ close to the equatorial inclusion and $S^m \looparrowright S^m \times S^m$ close to the diagonal.

The early work of Smale [21, 22] has been followed by a vast literature on the homotopy-theoretic properties of immersions, including [7], [9], [10], [11], [14], [18] and [20].

6. Appendix: Monomorphisms of vector bundles

Let ξ and η be smooth real vector bundles, of dimension n and r respectively, over a closed m -manifold M . We shall describe the differential-topological classification of homotopy classes of vector bundle monomorphisms $\eta \rightarrow \xi$ in the metastable range $m + 1 < 2(n - r)$.

Suppose that $v_0, v_1 : \eta \hookrightarrow \xi$ are two vector bundle monomorphisms. Doing homotopy theory, we may assume that ξ and η have positive-definite inner products and that the monomorphisms are isometric embeddings. Then v_0 and v_1 are sections of the bundle $O(\eta, \xi)$ whose fibre at $x \in M$ is the Stiefel manifold of orthogonal linear maps $v : \eta_x \rightarrow \xi_x$. Topological obstruction theory gives a difference class

$$\delta(v_0, v_1) \in \tilde{\omega}^{-1}(P(\eta)^{-H \otimes \xi}),$$

where $P(\eta)$ is the projective bundle of η and H is the Hopf line bundle. This arises as follows. A section of $O(\eta, \xi)$ determines a nowhere zero section of $H \otimes \xi$ over $P(\eta)$, as follows. An element $\ell \in P(\eta_x)$ is a line $\ell \subseteq \eta_x$. A monomorphism $v : \eta_x \rightarrow \xi_x$ gives an embedding of ℓ in ξ_x and so a non-zero vector in $\ell^* \otimes \xi_x$, which is the fibre of $H \otimes \xi$. Then $\delta(v_0, v_1)$ is defined as the difference class $\delta(s_0, s_1)$ of the two nowhere zero sections s_0 and s_1 of the vector bundle ξ over $P(\eta)$ constructed in this way from v_0 and v_1 . We may assume that s_0 and s_1 are sections of the sphere bundle $S(H \otimes \xi)$. Write $s_t = (1 - t)s_0 + ts_1$ for $0 \leq t \leq 1$. Then $\delta(s_0, s_1)$ is represented explicitly by the map, over $P(\eta)$,

$$\bar{s} : ([0, 1], \partial[0, 1]) \times P(\eta) \rightarrow (D(H \otimes \xi), S(H \otimes \xi))$$

given by the homotopy s_t . In the metastable range, the vector bundle monomorphisms v_0 and v_1 are homotopic if and only if $\delta(v_0, v_1) = 0$. (See, for example, [3, 5, 4].)

Thus far, the theory is topological. We now use Poincaré duality for the manifold $P(\eta)$ to identify $\tilde{\omega}^{-1}(P(\eta)^{-H \otimes \xi})$ with $\tilde{\omega}_0(P(\eta)^{H \otimes (\xi - \eta) - \tau M})$. (Up to homotopy, the stable tangent bundle is given by an isomorphism $\mathbb{R} \oplus \tau P(\eta) \cong (H \otimes \eta) \oplus \tau M$.) Assuming that the monomorphisms v_0 and v_1 are smooth we shall represent the dual obstruction class by a submanifold Z of M together with a map $Z \rightarrow P(\eta)$ and appropriate normal bundle information. The monomorphism v_0 will play a special rôle in the description; to emphasize this, we write $i = v_0$ for the preferred embedding $i : \eta \hookrightarrow \xi$ and write ν for the orthogonal complement of $i(\eta)$ in ξ . Let $X_k(\eta, \nu)$, for $k \geq 1$, be the sub-bundle of $O(\eta, \xi) = O(\eta, \eta \oplus \nu)$ with fibre consisting of those linear maps v such that $i_x + v$ has kernel of dimension k . By Lemma 6.3 below, we may assume, if $m + 1 < 2(n - r)$, that v_1 never meets $X_k(\eta, \nu)$ for $k > 1$ and is transverse to the sub-bundle $X_1(\eta, \nu)$. The inverse image $v_1^{-1}(X_1(\eta, \nu))$ is, therefore, a submanifold Z of M of dimension $m + r - n$, equipped with a section $Z \rightarrow P(\eta)$ classifying the 1-dimensional kernel. The normal bundle of Z in M is identified, by Lemma 6.3, with $H \otimes \nu$.

Proposition 6.1. *The submanifold Z described above, with the line bundle H classified by the section of $P(\eta)$ over Z and the isomorphism between the normal bundle and $H \otimes \nu$, represents the dual of $\delta(v_0, v_1)$ in $\tilde{\omega}_0(P(\eta)^{H \otimes \nu - \tau M})$.*

Proof. Consider the sections s_0 and s_1 of $S(H \otimes \xi)$ over $P(\eta)$ associated with v_0 and v_1 as in the definition of $\delta(v_0, v_1)$. The section \bar{s} of $D(H \otimes \xi)$ over $[0, 1] \times P(\eta)$ given by the homotopy $s_t = (1 - t)s_0 + ts_1$ is transverse to

the zero section and its zero-set is precisely $\{\frac{1}{2}\} \times Z$. The normal bundle is $\mathbb{R} \oplus \tau_M P(\eta) \oplus (H \otimes \nu)$, where $\tau_M P(\eta)$ is the tangent bundle along the fibres of $P(\eta) \rightarrow M$, and this is identified with $(H \otimes \eta) \oplus (H \otimes \nu) = H \otimes \xi$. Hence, Z with the normal bundle data represents the stable homotopy class dual to $\delta(s_0, s_1) = \delta(v_0, v_1)$. (This is the classical representation of the dual Euler class of a vector bundle by the zero-set of a generic smooth section.) \square

Remark 6.2. A more symmetric treatment may be given by looking at sections of the fibre product $O(\eta, \xi) \times_M O(\eta, \xi)$ and the sub-bundles with fibre consisting of the pairs (u, v) such that $\dim \ker(u + v) = k$.

The properties of v_1 required in the proof of Proposition 6.1 follow from the next lemma, in which the Lie algebra of the orthogonal group $O(V)$ of a Euclidean vector space V is written as $\mathfrak{o}(V)$.

Lemma 6.3. *Let V and W be finite-dimensional orthogonal vector spaces. For $0 \leq k \leq \dim V$, let $X_k(V, W)$ be the subspace of the Stiefel manifold $O(V, V \oplus W)$ consisting of the maps v such that $i + v$, where i is the inclusion of the first summand $V \hookrightarrow V \oplus W$, has kernel of dimension k . Then $X_k(V, W)$ is a submanifold diffeomorphic to the total space of the vector bundle $\mathfrak{o}(\zeta^\perp) \oplus \text{Hom}(\zeta^\perp, W)$ over the Grassmann manifold $G_k(V)$ of k -planes in V , where ζ^\perp is the orthogonal complement in V of the canonical k -dimensional vector bundle ζ over $G_k(V)$. Its normal bundle is naturally identified with $\mathfrak{o}(\zeta) \oplus \text{Hom}(\zeta, W)$.*

Proof. This can be established by using the (generalized) Cayley transform, which is written down explicitly in [5, Part II, Lemma 13.13]. The restriction of the normal bundle of the embedded submanifold to the subspace $G_k(V)$ is naturally identified with $\mathfrak{o}(\zeta) \oplus \text{Hom}(\zeta, W)$. The normal bundle itself is naturally identified with the pullback by parallel translation. \square

In particular, the submanifold $X_k(\eta, \nu)$ considered above has codimension $(n - r)k + k(k - 1)/2$ in $O(\eta, \eta \oplus \nu)$. So, if $k \geq 2$, the codimension is at least $2(n - r) + 1$. The condition $m < 2(n - r) + 1$ ensures that a generic section of $O(\eta, \eta \oplus \nu)$ is transverse to $X_1(\eta, \nu)$ and disjoint from $X_k(\eta, \nu)$ for $k > 1$.

Remark 6.4. In [15] Koschorke gave an intermediate representation of the dual of $\delta(v_0, v_1)$ by a submanifold of $(0, 1) \times M$. Consider the section \bar{v} of $\text{Hom}(\eta, \xi)$ over $[0, 1] \times M$ given by the homotopy $v_t = (1 - t)v_0 + tv_1$. Let $Y_k(\eta, \xi)$ be the sub-bundle of $\text{Hom}(\eta, \xi)$ with fibre consisting of the linear maps with kernel of dimension k ; it has codimension $(n - r)k + k^2$, which is $\geq 2(n - r) + 4 > m + 1$. Suppose that \bar{v} can be deformed, by a homotopy through maps coinciding with v_0 and v_1 at the endpoints, to a smooth section \bar{v}' , that never meets $Y_k(\eta, \xi)$ for $k > 1$ and meets $Y_1(\eta, \xi)$ transversely. This is always possible if $m < 2(n - r) + 3$. The inverse image of $Y_1(\eta, \xi)$ is then a submanifold Z of $(0, 1) \times M$ of dimension $m + r - n$ equipped with a map $Z \rightarrow P(\eta)$ given by the 1-dimensional kernel. This data, too, represents the dual of $\delta(v_0, v_1)$.

References

- [1] J.F. Adams, *Prerequisites (on equivariant stable homotopy) for Carlsson's lecture*. Proc. 1982 Aarhus Topology Conf., Springer Lecture Notes 1051, (1984), 483–532.
- [2] M. Boardman and B. Steer, *Hopf invariants*. Comm. Math. Helv. **42** (1967), 217–224.
- [3] M.C. Crabb, *$\mathbb{Z}/2$ -homotopy theory*. LMS Lecture Notes 44, Cambridge University Press, 1980.
- [4] M.C. Crabb, *The homotopy coincidence index*. J. Fixed Point Theory and Appl. **7** (2010), 1–32.
- [5] M.C. Crabb and I.M. James, *Fibrewise Homotopy Theory*. Springer, Berlin, 1998.
- [6] M.C. Crabb and A.A. Ranicki, *The geometric Hopf invariant and surgery theory*. In preparation.
Preliminary announcement <http://www.maths.ed.ac.uk/~aar/slides/geohopf.pdf>
- [7] J.P. Dax, *Étude homotopique des espaces de plongements*. Ann. Scient. École Norm. Sup. **5** (1972), 303–377.
- [8] A. Haefliger, *Plongements différentiables dans le domaine stable*. Comment. Math. Helv. **37** (1962/1963), 155–176.
- [9] A. Haefliger and M.W. Hirsch, *Immersions in the stable range*. Ann. of Math. **75** (1962), 231–241.
- [10] A. Hatcher and F. Quinn, *Bordism invariants of intersections of submanifolds*. Trans. Amer. Math. Soc. **200** (1974), 327–344.
- [11] M.W. Hirsch, *Immersions of manifolds*. Trans. Amer. Math. Soc. **93** (1959), 242–276.
- [12] J.R. Klein, *Poincaré complex diagonals*. Math. Zeit. **258** (2008), 587–607.
- [13] J.R. Klein and E.B. Williams, *Homotopical intersection theory, I*. Geometry & Topology **11** (2007), 939–977.
- [14] J.R. Klein and E.B. Williams, *Homotopical intersection theory, II*. Math. Zeit. **264** (2010), 849–880.
- [15] U. Koschorke, *Vector fields and other vector bundle morphisms – a singularity approach*. Lecture Notes in Mathematics 847, Springer, Berlin, 1981.
- [16] U. Koschorke, *Nielsen coincidence theory in arbitrary codimensions*. J. reine angew. Math **598** (2006), 211–236.
- [17] U. Koschorke and B. Sanderson, *Geometric interpretations of the generalized Hopf invariant*. Math. Scand. **41** (1977), 199–217.
- [18] B. Li, R. Liu and P. Zhang, *A simplified approach to embedding problem in normal bordism framework*. Systems Sci. Math. Sci. **5** (1992), 180–192.
- [19] A.A. Ranicki, *The algebraic theory of surgery II. Applications*. Proc. London Math. Soc. **40** (1980), 193–283.
- [20] H.A. Salomonsen, *Bordism and geometric dimension*. Math. Scand. **32** (1973), 87–111.
- [21] S. Smale, *Regular curves on riemannian manifolds*. Trans. Amer. Math. Soc. **87** (1958), 492–512.

- [22] S. Smale, *The classification of immersions of spheres in Euclidean spaces*. Ann. of Math. **69** (1959), 327–344.
- [23] C.T.C. Wall, *Surgery on compact manifolds*. Academic Press, 1971, 2nd Edition A.M.S., 1999.
- [24] H. Whitney, *The self-intersections of a smooth n -manifold in $2n$ -space*. Ann. of Math. **45** (1944), 220–246.

Michael Crabb
Institute of Mathematics,
University of Aberdeen,
Aberdeen AB24 3UE
Scotland, UK
e-mail: `m.crabb@maths.abdn.ac.uk`

Andrew Ranicki
School of Mathematics,
University of Edinburgh,
Edinburgh, EH9 3JZ
Scotland, UK
e-mail: `a.ranicki@ed.ac.uk`

References

1. J. F. Adams, *On the non-existence of elements of Hopf invariant one*, Ann. of Maths. 72, 20–104 (1960)
2. ———, *Infinite loop spaces*, Ann. of Maths. Study 90, Princeton (1978)
3. ———, *Prerequisites (on equivariant stable homotopy) for Carlsson’s lecture*, Proc. 1982 Aarhus Topology Conf., Springer Lecture Notes 1051, 483–532 (1984)
4. M. Boardman and B. Steer, *Hopf invariants*, Comm. Math. Helv. 42, 217–224 (1967)
5. T. Bröcker and E. C. Hook, *Stable equivariant bordism*, Math. Z. 129, 269–277 (1972)
6. W. Browder, *The Kervaire invariant of framed manifolds and its generalization*, Ann. of Maths. 90, 157–186 (1969)
7. ———, *Surgery on simply-connected manifolds*, Springer (1972)
8. P. E. Conner and E. E. Floyd, *Differentiable periodic maps*, Springer (1964)
9. S. Costenoble, *The structure of some equivariant Thom spectra*, Trans. A.M.S. 315, 231–254 (1989)
10. ———, *An introduction to equivariant cobordism*, in ‘Equivariant homotopy and cohomology theory’ CBMS 91, A.M.S., 153–162 (1996)
11. ——— and S. Waner, *G-transversality revisited*, Proc. A.M.S. 116, 535–546 (1992)
12. M. Crabb, *\mathbb{Z}_2 -homotopy theory*, L.M.S. Lecture Notes 44, Cambridge (1980)
13. ——— and I. M. James, *Fibrewise homotopy theory*, Springer Monographs in Mathematics, Springer (1998)
14. ——— and A. A. Ranicki, *The geometric Hopf invariant and double points*, J. Fixed Point Th. Appl. 7, The Steve Smale Festschrift, 325–350 (2010).
15. J.-P. Dax, *Étude homotopique des espaces de plongements*, Ann. Sci. École Norm. Sup. (4) 5, 303–377 (1972)
16. P. Eccles, *Double points of immersions of spheres in Euclidean space*, In: Prospects in Topology: Proceedings of a conference in honor of William Browder, Ann. of Maths. Studies 138, Princeton, 125–137 (1995)
17. T. tom Dieck, *Bordism of G-manifolds and integrality theorems*, Topology 9, 345–358 (1970)
18. E. Dyer and R. Lashof, *Homology of iterated loop spaces*, Amer. J. Math. 84, 35–88 (1962)
19. A. Haefliger, *Plongements différentiables dans le domaine stable*, Comment. Math. Helv. 37, 153–176 (1962/1963)
20. ——— and B. Steer, *Symmetry of linking coefficients*, Comm. Math. Helv. 39, 259–270 (1965)
21. A. Hatcher and F. Quinn, *Bordism invariants of intersections of submanifolds*, Trans. Amer. Math. Soc. 200, 327–344 (1974)

22. H. Hauschild, *Zerspaltung äquivarianter Homotopiemengen*, Math. Ann. 230, 279–292 (1977)
23. P. Hilton, *On the homotopy groups of the union of spheres*, J. Lond. Math. Soc. 30, 154–172 (1955)
24. M. Hirsch, *Immersions of manifolds*, Trans. A.M.S. 93, 242–276 (1959)
25. H. Hopf, *Über die Curvatura integra geschlossener Hyperflächen*, Math. Ann. 95, 340–367 (1925)
26. ———, *Über die Abbildung von Sphären auf Sphären niedriger Dimension*, Fund. Math. 25, 427–440 (1938)
27. I. M. James, *On the iterated suspension*, Quart. J. Math. Oxford Ser. (2) 5, 1–10 (1954)
28. ———, *Reduced product spaces*, Ann. of Maths. 62, 170–197 (1955)
29. ———, *On spaces with a multiplication*, Pacific J. Math. 7, 1083–1100 (1957)
30. ———, *The topology of Stiefel manifolds*, L.M.S. Lecture Notes 24, Cambridge (1976)
31. M. Kervaire, *Courbure intégrale généralisée et homotopie*, Math. Ann. 131, 219–252 (1956)
32. ———, *Relative characteristic classes*, Amer. J. Math. 79, 517–558 (1957)
33. ———, *An interpretation of G. Whitehead's generalization of H. Hopf's invariant*, Ann. of Maths. 69, 345–365 (1959)
34. ———, *Sur le fibré normal à une sphère immergée dans un espace euclidien*, Comm. Math. Helv. 33, 121–131 (1959)
35. ——— and J. Milnor, *Groups of homotopy spheres*, Ann. of Math. (2) 77, 504–537 (1963)
36. J. Klein, *Poincaré complex diagonals*, Math. Z. 258, 587–607 (2008)
37. U. Koschorke and B. Sanderson, *Geometric interpretations of the generalized Hopf invariant*, Math. Scand. 41, 199–217 (1977)
38. ——— and ———, *Self intersections and higher Hopf invariants*, Topology 17, 283–290 (1978)
39. P. Kühn, T. Macko and A. Mole, *The total surgery obstruction revisited*, Münster J. Math. 6, 181–269 (2013)
40. J. Levine, *An algebraic classification of some knots of codimension two*, Comm. Math. Helv. 45, 185–198 (1970)
41. J. P. May et al., *Equivariant homotopy and cohomology theory*, C.B.M.S. Regional Conference Series in Math. 91, A.M.S. (1996)
42. R. J. Milgram, *Unstable homotopy from the stable point of view*, Springer Lecture Notes 368, Springer (1974)
43. J. Milnor and J. Stasheff, *Characteristic classes*, Ann. of Maths. Study 76, Princeton (1974)
44. A. S. Mishchenko, *Homotopy invariants of non-simply connected manifolds. III. Higher signatures*, Izv. Akad. Nauk SSSR, ser. mat. 35, 1316–1355 (1971)
45. G. D. Mostow, *Equivariant embeddings in Euclidean space*, Ann. of Math. 65, 432–445 (1957)
46. L. Pontryagin, *Smooth manifolds and their applications in homotopy theory*, AMS Translations, Ser. 2, Vol. 11, 1–114 (1959)
47. F. Quinn, *Surgery on Poincaré and normal spaces*, Bull. Am. Math. Soc. 78, 262–267 (1972)
48. A. A. Ranicki, *The total surgery obstruction*, Proc. 1978 Aarhus Topology conference, Springer Lecture Notes 763, 275–316 (1979)
49. ———, *Algebraic theory of surgery I. Foundations*, Proc. Lond. Math. Soc. 40, 87–192 (1980)
50. ———, *Algebraic theory of surgery II. Applications to topology*, Proc. Lond. Math. Soc. 40, 193–283 (1980)
51. ———, *Exact sequences in the algebraic theory of surgery*, Mathematical Notes 26, Princeton (1981)

52. ———, *Algebraic L-theory and Topological Manifolds*, Cambridge Tract 102, CUP (1992)
53. ——— and M. Weiss, *Chain complex and assembly*, Math. Z. 204, 157–185 (1990)
54. W. Richter, *High-dimensional knots with $\pi_1 \cong \mathbb{Z}$ are determined by their complements in one more dimension than Farber's range*, Proc. A.M.S. 120, 285–294 (1994)
55. H. Samelson, $\pi_3(S^2)$, *H.Hopf, W.K.Clifford, F.Klein*, in History of Topology, 575–578, North-Holland, Amsterdam (1999)
56. G. B. Segal, *Equivariant stable homotopy theory*, Proc. 1970 Nice I.C.M., Vol. 2, 59–63 (1971)
57. ———, *Operations in stable homotopy theory*, New Developments in Topology, L.M.S. Lecture Notes 11, Cambridge (1974)
58. ———, private correspondence.
59. D. Sinha, *Real equivariant bordism and stable transversality obstructions for $\mathbb{Z}/2$* , Proc. A.M.S. 130, 271–281 (2001)
60. S. Smale, *Classification of immersions of spheres in Euclidean space*, Ann. of Maths. 69, 327–344 (1959)
61. V. P. Snaith, *A stable decomposition of $\Omega^n S^n X$* , J. Lond. Math. Soc. 7, 577–583 (1974)
62. N. Steenrod, *Cohomology invariants of mappings*, Ann. of Maths. 50, 954–988 (1949)
63. E. Stiefel, *Richtungsfelder und Fernparallelismus in n -dimensionalen Mannigfaltigkeiten*, Comment. Math. Helv. 8, 305–353 (1935)
64. R. E. Stong, *Bordism and involutions*, Ann. of Maths. 90, 47–74 (1969)
65. T. tom Dieck, *Orbittypen und äquivariante Homologie, II.*, Arch. Math. 26, 650–662 (1975)
66. P. Vogel, *Cobordisme d'immersions*, Ann. Sci. École Norm. Sup. (4) 7, 317–358 (1974)
67. C. T. C. Wall, *Classification of $(n - 1)$ -connected $2n$ -manifolds*, Ann. of Ma. 75, 163–189 (1962)
68. ———, *Surgery on compact manifolds*, Academic Press (1970); 2nd edition, A.M.S. (1999)
69. A. G. Wasserman, *Equivariant differential topology*, Topology 8, 127–150 (1969)
70. H. Weyl, *The classical groups*, Princeton (1939)
71. G. W. Whitehead, *A generalization of the Hopf invariant*, Ann. of Maths. 51, 192–237 (1950)
72. H. Whitney, *Differentiable manifolds*, Ann. of Maths. 37, 645–680 (1936)
73. ———, *On regular closed curves in the plane*, Comp. Math. 4, 276–284 (1937)
74. ———, *The self-intersections of a smooth n -manifold in $2n$ -space*, Ann. of Maths. 45, 220–246 (1944)
75. K. Wirthmüller, *Equivariant S-duality*, Archiv der Math. 26, 427–431 (1975)
76. R. M. W. Wood, *Framing the exceptional Lie group G_2* , Topology 16, 303–320 (1976)

Index

- Q -groups
 - V -coefficient hyperquadratic, $\widehat{Q}_V^n(C)$, 166
 - V -coefficient quadratic, $Q_n^V(C)$, 166
 - V -coefficient symmetric, $Q_V^n(C)$, 166
 - $[i, j]$ -quadratic, $Q_n^{[i, j]}(C)$, 163
 - $[i, j]$ -symmetric, $Q_{[i, j]}^n(C)$, 163
- \mathbb{Z}_2 -equivariant Pontryagin-Thom, 258
- \mathbb{Z}_2 -framed
 - \mathbb{Z}_2 -bordism, 258
 - \mathbb{Z}_2 -manifold, 258
- \mathbb{Z}_2 -isovariant
 - V -coefficient chain homotopy, 165
 - V -coefficient chain map, 164
 - composite chain map, 165
 - cup product, 165
- α_V , 26
- β_V , 26
- $\gamma(f, g)$, 7
- γ_V , 26
- π -equivariant
 - geometric Hopf invariant, $h_V(F)$, 352
 - geometric Hopf invariant, stable, $h'_V(F)$, 352
 - quadratic construction, $\psi_V(F)$, 352
 - spectral Hopf invariant, $sh_V(F)$, 353
 - symmetric construction, $\phi_V(X)$, 351
- s_V , 26
- z_V , 26
- absolute stable \mathbb{Z}_2 -equivariant
 - cohomotopy, $\omega_{\mathbb{Z}_2}^n(X)$, 98
 - homotopy, $\omega_{\mathbb{Z}_2}^{\mathbb{Z}_2}(X)$, 98
- algebraic mapping cone, $\mathcal{C}(f)$, 11
- assembly map, A , 264, 358
- bi-degree, 89
- cellular diagonal chain approximation, Δ , 165
- cone, cX , 2
- curvatura integra, 247
- degree, 85
- difference map
 - $\overline{\nabla}_V$, 23
- difference map of stable maps
 - $p - q$, 30
- double point set
 - $X \times_Y X$, 264
 - ordered, $D_2(f)$, 264
 - unordered, $D_2[f]$, 265
- Dyer-Lashof map, d_V , 294
- embedded map, 265
- embedding of map, 265
- embedding-immersion pair, 302
- fibrewise
 - map, 357
 - pointed space, 358
 - product, $X \sqcup_B Y$, 358
 - product, $X \times_B Y$, 357
 - smash product, $X \wedge_B Y$, 358
 - space, 357
 - suspension, $\Sigma_B X$, 359
 - Thom space, $T_B(\xi)$, 359
- fixed point set, 77
- framed
 - bordism, $\Omega_m^{fr}(X)$, 246
 - manifold, 245
- framed manifold, (M, b) , 245
- free π -space, 78
- generalized Gauss map, 246
- geometric Hopf invariant

- $h_V(F)$, 191
 - stable, $h'_V(F)$, 204
- Grassmann manifold, $G(V, W)$, 50
- homotopy pushout, $\mathcal{C}(f_1, f_2)$, 293
- Hopf invariant
 - framed manifold, 247
- inner product \mathbb{Z}_2 -space, 79
- isotopy, 17
- join, $X * Y$, 2
- local geometric Hopf map, $h_V(F)_Y$, 266
- mapping
 - cone, $\mathcal{C}(F)$, 2
 - cone, reduced $\mathcal{C}(F)$, 2
 - cylinder, $\mathcal{M}(F)$, 2
 - cylinder, reduced $\mathcal{M}(F)$, 2
- negative
 - circle, $S^-(V)$, 215
 - difference, $\delta^-(p, q)$, 215
 - line, V^- , 215
- null-homotopy, 2
- one-point
 - adjunction, X^+ , 1
 - compactification, X^∞ , 15
- open
 - embedding, 16
 - isotopy, 17
- orientation-preserving
 - embedding, 23
 - homotopy equivalence, 22
- orientation-reversing
 - embedding, 23
 - homotopy equivalence, 22
- orthogonal group, $O(V)$, 50
- Pontryagin-Thom, 303
 - map, α_M , 245
- positive
 - circle, $S^+(V)$, 215
 - difference, $\delta^+(p, q)$, 215
 - embedding, 274
 - line, V^∞ , 215
 - linear automorphism, $GL^+(V)$, 273
- positive orthogonal automorphism, $O^+(V)$, 273
- projective space $P(V)$, 18
- quadratic construction, $\psi_V(F)$, 207
- reduced cone, CX , 2
- reduced stable \mathbb{Z}_2 -equivariant
 - cohomotopy, $\tilde{\omega}_{\mathbb{Z}_2}^n(X)$, 98
 - homotopy, $\tilde{\omega}_{\mathbb{Z}_2}^{\mathbb{Z}_2}(X)$, 98
- reduced suspension, ΣX , 2
- relative difference
 - chain homotopies, $\delta(p, q)$, 11
 - maps, $\delta(f, g)$, 5
 - stable maps, $\delta(p, q)$, 30
- reverse of a homotopy, $-f$, 4
- second extended power, $e_2(\alpha)$, 305
- semicharacteristic, $\chi_{1/2}(M)$, 247
- semidegree
 - chain map, 87
 - pointed map, 87
- semifree pointed π -space, 78
- separation element, $\delta(f, g)$, 6
- smash product, $X \wedge Y$, 1
- spectral Hopf invariant, $sh_V(F)$, 221
- spectral quadratic construction, $s\psi_V(F)$, 224
- square
 - map $f \times f$, 264
- square embedding-immersion, 304
- stable \mathbb{Z}_2 -equivariant homotopy
 - group, $\{X; Y\}_{\mathbb{Z}_2}$, 93
- stable \mathbb{Z}_2 -equivariant homotopy group
 - of sphere, $\omega_{\mathbb{Z}_2}^n$, 98
- Stiefel manifold, $O(V, W)$, 50
- sum map, ∇_V , 23
- sum of homotopies, $f + g$, 4
- suspension
 - chain complex, SC , 11, 162
 - space, sX , 2
- Thom space, $T(\xi)$, 47
- transposition
 - T , 104
 - κ_V , 104
- trivial π -action, 78
- ultraquadratic construction, $\hat{\psi}(F)$, 214, 353
- Umkehr
 - adjunction, 16, 303
 - compactification, 16, 265
 - geometric, 333
 - stable, 44
- unit
 - disc, $D(V)$, 18
 - sphere, $S(V)$, 18